Vol. 92, Sect. 9, 1955

MOLLUSCA

COMPILED BY

A. E. SALISBURY, A.M.I.E.E.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE TEN SHILLINGS
December, 1957

SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The following are the Sections and their prices, viz .:-

-	20 20220					T	,	-		
		1						s.	d.	
1	. Comprehen	sive Z	oology					3	0	
2	. Protozoa							10	0	
3	. Porifera							2	0	
4	. Coelenterat	a						4	0	
5	. Echinodern	na						4	0	
6	. Vermes							10	0	
7	. Brachiopod	a						4	0	
8	Bryozoa		V					2	6	
9	. Mollusca	\						10	0	
10	. Crustacea							6	6	
11	. Trilobita							5	0	
12	. Arachnida							10	0	
* 13	. Insecta							50	0	
14	. Protochord	ata						2	0	
15	. Pisces							8	6	
16	A. Amphibia							6	0	
16	в. Reptilia							7	6	
17	. Aves							10	0	
18	. Mammalia							10	0	
19	. List of Nev	w Gen	era and	Subg	genera			4	0	

^{*} Obtainable only from the Commonwealth Institute of Entomology, 56, Queen's Gate, London, S.W.7.

9. MOLLUSCA

COMPILED BY

A. E. SALISBURY, A.M.I.E.E.

CONTENTS

							1	PAGE
I.	TITLES	•••	•••	•••		•••	•••	2
II.	SUBJECT INDEX					***		47
III.	Systematic Index			•••		•••		71
	Polyplacophora	• • •		•••	• • •		• • •	71
	Gastropoda							72
	Prosobranchiata					•••		73
	Docoglossa							74
	Opisthobranchiata							90
	Nudibranchiata							91
	Pulmonata							93
	Pelecypoda or Lan	nellibra	anchia	·				101
	Cephalopoda							112
	Incertae Sedis							122

I.—TITLES

Direction 14. Addition to the "Official List of Family-Group Names in Zoology," or, as the case may be, to the "Official Index of Rejected and Invalid Family-Group Names in Zoology" of the family-group names involved in the applications relating to the names of ammonites published in volume 2 of the "Bulletin of Zoological Nomenclature." Opin. int. Comm. zool. Nom. 10 1955: 463–480.

Direction 27. Addition to the "Official List of Family-Group Names in Zoology" of family group names based upon the names of certain genera of non-marine Mollusca placed on the "Official List of Generic Names in Zoology" by the ruling given in Opinion 335. Opin. int. Comm. zool. Nom. 10 1955: 481–492.

Opinion 215. Emendation to Palaeoneilo of the generic name Palaeaneilo Hall, 1869. Opin. int. Comm. zool. Nom. 4 1954: 51-62.

Opinion 217. Designation under the plenary powers, of a type species for the genus *Erycina* Lamarck, 1805 (Class Pelecypoda) in harmony with accustomed usage. Opin. int. Comm. zool. Nom. 4 1954: 73–82.

Opinion 221. Emendation to macfarlandi of the specific name published as mcfarlandi Cockerell, 1902 in the combination Chromodoris mcfarlandi. Opin. int. Comm. zool. Nom. 4 1954: 115-124.

Opinion 225. Validation, under the plenary powers, of *Belone* Cuvier, 1817 (Class Pisces) and *Raphistoma* Hall, 1847 (Class Gastropoda). Opin. int. Comm. zool. Nom. 4 1954: 161–176.

Opinion 233. Suppression, under the plenary powers, of the name Octopodia Schneider, 1784 (Class Cephalopoda), and of certain reputed names published by the same author in 1784. Opin. int. Comm. zool. Nom. 4 1954: 275–296.

Opinion 235. Designation, under the plenary powers, of a lectotype for the nominal species *Ammonites cordatus* Sowerby (J.) 1813. Opin. int. Comm. zool. Nom. 4 1954: 313–328.

Opinion 237. Addition of the generic name Tonna Brünnich, 1771 (Class Gastropoda) to the Official List of Generic Names in Zoology. Opin. int. Comm. zool. Nom. 4 1954: 343–350.

Opinion 287. Validation, under the plenary powers, of the generic name Scaphander Montfort, 1810. Opin. int. Comm. zool. Nom. 8 1954: 49-62.

Opinion 292. Validation, under the plenary powers, of the generic name Cardinia (Class Lamellibranchiata) as from Agassiz (J.L.R.) [1841] for use in its accustomed sense. Opin. int. Comm. zool. Nom. 8 1954: 115–130.

Opinion 296. Suppression, under the plenary powers, for nomenclatorial purposes, of volume 3 (Regnum lapideum) of the twelfth edition of the Systema Naturae of Linnaeus published in 1768 and of the corresponding volume in the Houttuyn (1785), Gmelin (1793), and Turton (1806) editions of the above work. Opin. int. Comm. zool. Nom. 8 1954: 167–178.

Opinion 300. Validation, under the plenary powers, of the generic name Sphaeroceras Bayle, 1878. Opin. int. Comm. zool. Nom. 8 1954: 237-248.

Opinion 301. Designation, under the plenary powers, of a type species for *Procerites* Siemiradzki 1898 (Class Cephalopoda, Order Ammonoidea) in harmony with accustomed nomenclatorial usage. Opin. int. Comm. zool. Nom. 8 1954: 249–260.

Opinion 302. Designation, under the plenary powers, of a type species, for the genus Aulacostephanus Tornquist, 1896 (Class Cephalopoda, Order Ammonoidea) in harmony with accustomed nomenclatorial usage. Opin. int. Comm. zool. Nom. 8 1954: 261–272.

Opinion 303. Designation, under the plenary powers, of type species in harmony with current usage for the nominal genera Kosmoceras, Perisphinctes and Harpoceras (Class Cephalopoda, Order Ammonoidea) established by Waagen in 1869. Opin. int. Comm. zool. Nom. 8 1954: 273–284.

Opinion 304. Suppression, under the plenary powers, of five early generic names for Ammonites (Class Cephalopoda, Order Ammonoidea) now fallen into desuetude. Opin. int. Comm. zool. Nom. 8 1954: 285–296.

Opinion 335. Addition to the Official list of Generic names in Zoology of the names of thirty-four non-marine genera of the Phylum Mollusca. Opin. int. Comm. zool. Nom. 10 1955: 45-76.

Opinion 336. Addition to the Official List of specific names in Zoology of the specific names of one hundred and twenty-two non-marine species of the Phylum Mollusca. Opin. int. Comm. zool. Nom. 10 1955: 77–108.

Opinion 337. Rejection of the generic name "Arieticeras" Quenstedt, 1883 (Class Cephalopoda, Order Ammonoidea) and addition of the name "Arieticeras" Seguenza, 1885 to the Official List of Generic names in Zoology. Opin. int. Comm. zool. Nom. 10 1955: 109–124.

Opinion 338. Acceptance of the mesozoic fossil species "Gryphaea arcuata" Lamarck, 1801, as the type species of the nominal genus "Gryphaea" Lamarck, 1801 and addition of the generic name "Gryphaea" Lamarck, 1801 (Class Pelecypoda) to the Official List of Generic names in Zoology. Opin. int. Comm. zool. Nom. 10 1955: 125–180, 1 pl.

Opinion 344. Validation under the plenary powers of the generic name 'Truncatella' Risso, 1826, and addition of that name and the names 'Acmaea' Eschscholtz, 1833, and 'Acicula' Hartmann, 1821 (Class kastropoda) to the "Official List of keneric Names in Zoology." Opin. int. 'Jomm. 2001. Nom. 10 1955: 313–352.

Opinion 351. Acceptance of the mendation to *Dreissena* of the generic ame *Driessena* van Beneden 1835. Class Pelecypoda.) Opin. int. Comm. ool. Nom. 11 1955: 1–24.

Opinion 353. Validation under the lenary powers of the generic name oplites Neumayr, 1875 (Class Cephaloda, Order Ammonoidea). Opin. int. omm. zool. Nom. 11 1955: 47–78.

Opinion 354. Addition to the Official List of Specific Names in Zoology of the specific name fasciata Poiret, 1789, as published in the combination Laplysia [recte Aplysia] fasciata (Class Gastropoda). Opin. int. Comm. zool. Nom. 11 1955: 79-90.

Opinion 355. Validation, under the Plenary Powers, of the specific name "punctata" Cuvier, 1803, as published in the combination Laplysia [recte "Aphysia"] punctata (Class Gastropoda). Opin. int. Comm. zool. Nom. 11 1955: 91-102.

Opinion 356. Addition to the Official List of Family-Group Names in Zoology of the names of the family-group taxa having Ostrea Linnaeus, 1758, and Gryphaea Lamarck, 1801 (Class Pelecypoda) as their respective type genera (Opinion supplementary to Opinion 338). Opin. int. Comm. zool. Nom. 11 1955: 103-112.

Opinion 360. Validation under the plenary powers of the specific name "knorrii" Voltz, 1828 as published in the combination Ostrea knorrii. Opin. int. Comm. zool. Nom. 11 1955: 151–160.

Opinion 361. Validation under the plenary powers of the generic name "Antalis" Adams (H.) & Adams (A.) [1854]. [Class Scaphopoda]. Opin. int. Comm. zool. Nom. 11 1955: 161-172.

Opinion 362. Rejection for nomenclatorial purposes of Geoffroy (E. L.) 1767, "Traité Sommaire des Coquilles, tant fluviatiles que terrestres, qui se trouvent aux environs de Paris." Opin. int. Comm. zool. Nom. 11 1955: 173– 182.

Opinion 363. Designation, under the plenary powers, of a type species in harmony with accustomed usage for the nominal genus "Ancylus" Müller (O. F.) 1774. (Class Gastropoda.) Opin. int. Comm. zool. Nom. 11 1955: 183–202.

Opinion 364. Addition of the name "Ancylastrum" Bourguignat, February 1853 (Class Gastropoda) to the Official List of Generic Names in Zoology with "Ancylus (Ancylastrum) cumingianus" Bourguignat, May 1853, as the type species of the genus named. Opin. int. Comm. zool. Nom. 11 1955: 203-214.

- Anon. (1). Merrill E. Champion. Occ. Pprs. on Mollusks Harvard Cambridge 1945–1954 pp. iii with portrait.
- **Anon.** (2). Louisa Fischer (1871–1954). J. Conchyliol., Paris **95** 1955: 3–5.
- Anon. (3). Conus litteratus L. Mins. Conch. Club S. Calif. No. 147 1953: 2-3.
- Anon. (4). List of Japanese species of the Conidae. Venus, Fukuyama 18 4 1955: 288-302.
- Abbott, R. T. Shells of superstition. Frontiers 20 1 1955: 12-13, 33.
- Abbott, R. T. The Titian R. Peale shell collection. Nautilus, Philad. 68 1955: 123-127 pl. 4 figs. 1-6.
- Abbott, R. T. (1). Anatomy of the Venezuelan gastropod *Doryssa kappleri*. Nautilus, Philad. **69** 1955 : 44–46 pl. 4.
- Abdel-Malek, E. T. Anatomy of Biomphalaria boissyi as related to its infection with Schistosoma mansoni. Amer. Midl. Nat. 54 1955: 394-404 4 pls.
- Abdussalam, M. & Sarwar, M. M. Intermediate host of *Schistosoma turkestanicum* in Pakistan. Proc. Pakist. sci. Conf. 6th (3): 228.
- Accordi, B. Interessanti rinvenimenti di *Lingula* e di *Myophoria* nel Werfeniano inferiore del Bellunese e del Bergamasco. Riv. ital. Paleont. **61** 2 1955: 47–52 pl. 7.
- Adam, W. & Knudsen, J. Note sur quelques espèces de mollusques marins nouveaux ou peu connus de l'Afrique Occidentale. Bull. Inst. Sci. nat. Belg. 31 61 1955: 1–25 2 pls. 4 figs.
- Aguayo, C. G. & Jaume, M. L. Indice al catalogo de los moluscos de Cuba. Circ. Mus. biblio. zool. Habana 1954: 1359-1374.
- Ahders, C. Shells of Fire Island [near Long Island, U.S.A.]. Circ. Mus. biblio. zool. Habana 1954: 1083-1084.
- Airan, J. W. & Joshi, J. V. Distribution of sulphur and nitrogen in *Sepia* and *Rhabdocynthia*. J. Univ. Bombay N.S. 21A 3 1955: 72-73.
- Aldrich, F. A. A new symbiotic relationship involving *Molgula manhattensis* (De Kay) and *Mya arenaria* L. Notul. nat. Acad. Philad. No. 274 1955: 1–3 1 fig.

- Alexander, F. E. S. Nassarius thersites bored through the apertural callus by a predator. Sarawak Mus. J. 6 1955: 321.
- Allan, J. The Kraken—legendary terror of the seas. Aust. Mus. Mag. 11 1955: 275–278 2 figs.
- Allen, F. E. Identity of breeding temperatures in Southern and Northern hemisphere species of *Mytilus* (Lamellibranchia), Pacif. Sci. 9 2 1955: 107–109 2 figs.
- Allen, M. F. & J. A. On the habits of Pandora inaequivalvis (Linné). Proc. Malac. Soc. London 31 1955: 175–185 5 text-figs. & graphs.
- Allison, E. C. Middle Cretaceous gastropoda from Punta China, Baja California, Mexico. J. Paleont. 29 3 1955: 400-432 pls. 40-444 text-figs. 1-3.
 - Almela, A. see Rios, J. M.
- Altena, C. O. v. R. Faunistische aantikeningen, 4, Over het vermeende voorkomen von *Epitonium trevelyanum* in Nederland. Basteria 19 2/3 1955: 26. [Summary in English.]
- Altena, C. O. v. R. (1). Notes sur les limaces. 2. Description d'une espèce nouvelle de *Gigantomilax* provenant des environs de Kouldja (Chine). Beaufortia 4 36 1954: 47-50 3 figs.
- Altena, C. O. v. R., Bloklander, A. & Pouderoyen. De fossiele schelpen van de Nederlandse stranden en zeegaten. Basteria 19 2/3 1955: 27-34 pls. 5-8.
- Altena, C. O. v. R., Aten, R. & Schouten, A. R. Notes sur les Limaces. 3. [Arion lusitanicus Mab., anatomy, figs.] J. Conchyliol. Paris 95 1955: 89-99 6 text-figs.
- Amano, M. Occurrence of a new species of *Pleuronectites* in the Triassic of Japan. Trans. paleont. Soc. Japan N.S. 17 1955: 23–28 pl. 5.
- Amio, M. On the egg masses and larvae of seven species of Naticidae (Gastropoda) from the coastal region of Japan. J. Shimonoseki Coll. Fish. 5 1955: 137-158 9 figs.
- Andjelković, M. Ž. Existence des sédiments fossilifères neritiques et bathyals du Crétacé inférieur au sud du village Rusanj (environs de Beograd). Zborn. Radova geol. Inst. Beograd. 7 1954: 69-87 pls. 1-2 text-figs. 1-3. [French résumé p. 86.]

- Andjelković, M. ž. L'Urgonien et l'Albien dans la zone crétacée de Topola-Drača (Sumadija-Serbie). Ann. géol. Pén. balkan 22 1954: 27–73 pls. 1–8 text-figs. 1–14. [French résume p. 73.]
- Andreu, B. & Arte, P. Experiencias previas sobre fijacíon de larvas y crecimiento invernal de la ostra (O. edulis) joven en las rías gallegas (N.W. de España). Publ. Inst. Biol. apl. Barcelona 19 1955: 115–129 figs. 1–3. [English summary.]
- Andrusov, D. & Kováčik, J. The Carpathian Mesozoic fossils: Part II. Triassic Cephalopods of Slovakia and the stratigraphic sequence of the Triassic in Slovakia. Geol. sbornik. Slovensk. Akad. 6 3–4 1955: 258–301 pls. 9–10 text-figs. 7–8. [Russian and English resumé.]
- Angel, L. M. Parorchis acanthus var. australis, n. var., with an account of the life cycle in South Australia. Trans. roy. Soc. S. Aust. 77 1954: 164-174 4 figs.
 - Angelescu, V. see Popovici, Z.
- Anteunis, A. Succinea arenaria Bouchard - Chantereaux een nieuwe soort voor België. Basteria 19 2/3 1955 : 23–25 text-figs.
- Aoki, S. Mollusca from the Miocene Kabeya Formation, Jôban coal-field, Fukushima Prefecture, Japan. Sci. Rep. Tokyo Kyoika Daig. 3C 17-21 1954: 23-41 2 pls. map.
- Arias C., S. Los Pupillidae (Pulmonata, Stylommatophora: Orthurethra) colestados en Venezuela septentrional. Mem. Soc. Cienc. nat. La Salle 15 41 1955: 140–169 6 figs. [English summary.]
- Arkell, W. J. Designation of a type genus for the nominal order "Ammonoidea" Zittel, 1884, and proposed addition of that name to the "Official list of names in the Order/Class group in Zoology." Bull. zool. Nom. 11 1955: 199-200.
- Arkell, W. J. (1). Proposed use of the plenary powers to validate and protect the family-group name "Mayaitidae" Spath (L. F.), 1928 (Class Cephalopoda, Order Ammonoidea). Bull. zool. Nom. 11 1955: 229–231.

- Arkell, W. J. (2). Proposed suppression under the plenary powers of the family-group name Seguenziceratidae Spath (L. F.) 1924. Bull. zool. Nom. 11 1955: 297–298.
- Arkell, W. J. (3). Monograph of the English Bathonian ammonites. Palaeontogr. Soc. [Monogr.] 5 1955: 129–140 pls. 16–17 text-figs. 47–52.
- Arkell, W. J. (4). Proposed adoption of a declaration that a generic or specific name based solely upon the aptychus of an ammonite (Class Cephalopoda, Order Ammonoidea) be excluded from availability under article 27 of the "Règles" and proposed suppression of certain such names under the plenary powers. Bull. zool. Nom. 9 1954: 266–269.
- Arkell, W. J. & Donovan, D. T. Proposed use of the plenary powers to suppress the generic name *Caenisites* Buckman, 1925 founded upon a monstrosity. Bull. zool. Nomencl. 6 1954: 364–366.
- Arnould, M. La collection de mollusques testaces de la Station Océanographique de Salammbo. Bull. Sta. océanogr. Salammbo No. 51 1955 : 3–21.
 - Arte, P. see Andreu, B.
- Asano, M. Occurrence of choline in the shellfish, Callista brevisiphonata Carpenter. Tohoku J. agric. Res. 43-41954: 239-250 2 figs.
- Asano, M. & Sato, H. Biochemical studies on octopus. I. Trimithylamine and trimithylamine oxide contents of octopus. Tohoku J. agric. Res. 5 3 1954: 191–195.
- Astre, G. Listes stratigraphiques des faunes de Radiolitidés nord-pyrénées. J. Conchyliol., Paris 95 1955: 6–10.
- Aten, D. see Altena, C. O. v. R. and others.
- **Attardo, C.** Effetti dell'azide sodico sulle uova di *Bithynia codiella*. R.C. Accad. Lincei (8) **19** 1–2 1955: 83–84 3 figs.
- Augusta, J. & Pokorný, V. Critical remarks on the Stegocephalia tracks in the lower Namurian at Kyjovice, Silesia, Č.S.R. Mém. Soc. Sci. Bohème 1949 [1950] 6: 1-6. [English summary.]

Avias, J. Contribution à l'Etude Stratigraphique et Paléontologique des formations antécrétacées de la Nouvelle-Calédonie Central. Sciences de la Terre, Nancy 1 1–2 1953: 1–276 pls. 14–20, 23–25 text-figs. 1–117 maps & tables.

Azevedo, J. F. de. see Fraga de Azevedo.

Baba, K. Notes on the species of the genus *Notarchus* (Subg. *Stylocheilus*) from Japan. Zool. Mag. Tokyo **63** 1954: 218–221 2 figs.

Baden-Powell, D. F. W. Report on the marine fauna of the Clacton channels in Warren, S. H. Quart. J. geol. Soc. Lond. 111 3 1955: Appendix B 301-304.

Bader, R. G. The role of organic matter in determining the distribution of pelecypods in marine sediments. J. Mar. Res. 13 1 1954: 32-47 4 figs.

Baily, J. L. Support for the validation under the plenary powers of *Bithynia* Leach, 1818. Bull. zool. Nom. 11 1955: 279-280.

Baker, A. A. see Gill, E. D.

B[aker], H. B. Dates of the Nautilus. Nautilus, Philad. 69 1955: 24.

Baker, H. B. (1). Heterurethrous and aulacopod. Nautilus, Philad. 68 1955: 109-112.

Bányai, M. see Kiss-Kocsisné Bányai.

Barkman, J. J. On the distribution and ecology of *Littorina obtusata* (L.) and its subspecific units. Arch. néerl. Zool. 11 1955: 22–86 9 figs.

Barnard, K. H. South African shore life, Cape Town (Maskew Miller Ltd.) 1954 pp. 135 25 pls.

Baroni, C., Incitti, L., Oliveri, A. & Viola, V. Revisione della fauna Neocretacica della Libia: Fam. Pinnidae, Mytilidae, Limidae, Nuculidae, Ledidae, Arcidae, Cyprinidae, Astartidae, Crassatellidae, Veneridae, Cardiidae, Lucinidae, Aloididae, Thraciidae, Chamidae. Ann. Mus. libico Stor. nat. 4 1953: 11–109 6 pls.

Bartha, F. & Soós, L. Die pliozäne Molluskenfauna von Balatonszentgyörgy. Ann. hist.-nat. Mus. hung. N.S. 6 1955: 51–72 2 pls.

Bartsch, P. The pyramidellid mollusks of the Pliocene deposits of North St. Petersburg, Florida. Smithson. misc. Coll. 125 2 1955: 1-102 18 pls.

Basse, E. Description d'une nouvelle espèce d'ammonite tunisien appartenant au genre *Neolobites* Fischer 1882: *Neolobites medeninensis* nov. sp. et considérations sur ce genre. Bull. Soc. Sci. nat. Tunis 6 1954: 197–204 1 pl.

Basse, E. (1). Fossiles du Jurassique supérieur et des "Grès de Nubie" de la région de Sana (Yémen). Bull. Soc. géol. Fr. 6 4 7-9 1954 : 663-673 pl. 27 text-fig. 1.

Baugh, S. C. see Lal, M. B.

Baughman, J. L. From oyster shell to ice cream. Circ. Mus. biblio. zool. Habana 1954: 1264-1266.

Beenakkers, A. M. T. see Raven, C. P.

Beetle, D. E. The terrestrial and aquatic mollusks of Albany County, Wyoming. J. Colo.-Wyo. Acad. Sci. 4 6 1954: 59.

Benitez, F. C. La Estrella de Mar y la disminución de erizos y algunos moluscos en el Golfo de Arauco. Rev. Biol. mar. 5 1-3 1955: 219-224.

Bennison, G. M. A Myalina fauna from the Namurian of north Ayrshire. Geol. Mag. 92 1955: 448–456 text-figs. 1–2.

Bernhauser, A. Zur Kenntnis der Retzer Saude. S.B. öst. Akad. Wiss. Abt. I 164 3 1955: 163–192 1 pl. 12 figs.

Bernstein, J. The unorthodox seahare. Nat. Hist. N.Y. 64 1955: 518-521 figs.

Berry, S. S. A new Sierran pulmonate of the genus *Monadenia*. Bull. S. Calif. Acad. Sci. **54** 1955: 14–16 1 pl.

Berry, S. S. (1). On recent Californian occurrences of the rare octopod *Ocythoe*. Calif. Fish Game 41 1955: 177–181 1 fig.

Berry, S. S. (2). An important new land snail from the Mission Range, Montana. Bull. S. Calif. Acad. Sci. 54 1955: 17-19 2 pls.

Berry, S. S. (3). On the supposed stenobathic habitat of the California sea-mussel. [Mytilus californianus.] Calif. Fish Game 40 1954: 69-73 1 fig.

Bertholdt, W. Are snails useful in aquaria? Aquarium, Philad. 23 1954: 204–207 7 figs.

Beth, K. see Nümann, W.

- Beurlen, K. Horizontes fossiliferos das Camadas Serra Alta do Paraná. Bol. Geol. Min. Caracas 152 1954: 1–30 1 pl. 1 text-fig. & sections.
- Biberson, P. Nouvelles observations sur la faune de mollusques de Quaternaire marin de la région de Casablanca. C.R. Soc. Sci. nat. Maroc No. 2 1955 : 60-62.
- Binder, E. Mollusques nouveaux de Cote d'Ivoire. Prosobranches d'éau douce. Rev. suisse Zool. 62 Suppl. 1955: 73-82 18 figs.
- Birkenmajer, K. & Znosko, J. Contribution to the stratigraphy of the Dogger and Malm in the Pieniny Klippen-Belt (Central Carpathians). Ann. Soc. géol. Pologne 23 1955: 3-36 pls. 1-2. [In English, Russian and Polish.]
- Blažka, P. Temperaturadaptation des Gesamtmetabolismus bei der Weinbergschnecke *Helix pomatia* L. Zool. Jb. (Allg. Zool.) 65 1955: 430–438 6 figs.
- Bloklander, A. see Altena, C. O. v. R. and others.
- Blume, W. Systrophia stenogyra zischkai, eine neue Lokalrasse aus Bolivien. Arch. Molluskenk. 84 1955: 179–180 text-fig.
- Boekelman, H. J. Archaeo- and Ethno-Conchology. Circ. Mus. biblio. zool. Habana 1954: 1496–1499.
- Boettger, C. R. Zoogeographische Betrachtungen über die europäischen Süsswasser schnecken der Gattung Viviparus Montfort, Arch, Molluskenk. 84 1955: 87–95.
- Boettger, C. R. (1). Die Systematik der euthyneuren Schnecken. Verh. dtsch. zool. Ges. 1954 1955: 253–280 1 fig.
- Boettger, C. R. (2). Flussperlmuschel und Perlenfischerei in der Lüneberger Heide. Abh. braunschw. Wiss. Ges. 6 1954: 1–40 3 figs. [English summary.]
- Boettger, C. R. (3). Zur Frage der Verteilung bestimmter Varianten bei der Landschneckengattung Cepaea Held. Biol. Zbl. 73 5-6 1954: 318-333.
- Bolognari, A. Aspetti dell'ovogenesi di Aplysia depilans L. (Moll. Gast. Opis.). Boll. Zool. 21 1954: 185–198 2 pls.

- Bolognari, A. (1). Ricerche sulla sessualità di *Haliotis lamellosa* Lam. Arch. 2001. ital. Napoli 38 1954 : 361–402 3 pls.
- Bolognari, A. (2). Alcuni aspetti dell' ovogenesi di *Haliotis lamellosa* Lam. (Moll. Gas. Pros.). Boll. Accad. Gioenia (4) 2 7 1954: 420–435 1 pl. [English, French, German summaries.]
- Borrello, A. V. Sobre el Valanginense del arroyo de las Yeseras, norte del rio Atuel, Mendoza. Notas Mus. Eva Perón 16 1953: 127–136.
- Bouillon, J. Sur l'anatomie de Caelatura aegyptiaca Caillaud 1827 f. calathus Bourguignat 1885. Rev. Zool. Bot. afr. 51 1955: 58-64 3 figs.
- Bouillon, J. (1). Sur l'anatomie et la position systématique du Gastéropode thalassoïde *Martelia tanganyicensis* Dautzenberg 1908. Rev. Zool. Bot. afr. 52 1955: 233–240 3 figs.
- Bousfield, E. L. Viviparus viviparus L. in Eastern Canada. Canad. Fld Nat. 69 1 1955; 27–28.
- Bowsher, A. L. Origin and adaption of platyceratid gastropods. Paleont. Contr. Univ. Kans. Moll. art. 5 1955: 1–11 pls. 1–2 text-fig. 1.
- Boycott, B. B. Learning in Octopus vulgaris and other Cephalopods. Pubbl. Staz. zool. Napoli 25 1954: 67–93 8 figs.
- Boycott, B. B. & Young, J. Z. Memories controlling attacks on food objects by *Octopus vulgaris* Lamarck. Pubbl. Staz. zool. Napoli 27 1955: 232–249.
- Brabenec, J. Vodní měkkýši Československa a jejich systematické zařazení. [Czechoslovakian aquatic molluscs.] Přírod. Sborn. Ostravsk. Kraje 13 1–2 1952: 135–165 12 pls.
- Brabenec, J. (1). Malakozoologický rýzkum Sleszka a některých částé Západnich Karpat. Přírod. Sborn. Ostravsk. Kraje 14 3–4 1953 : 428–469 4 pls. See also p. 53–4–.
- Brady, L. F. Possible nautiloid mandibles from the Permian of Arizona. J. Paleont. 29 1 1955: 102–104 pl. 21.
- Braga, J. M. Materias para o estudo da fauna Malacológica de Moçambique. An. Jta. Miss. geogr., Lisboa 7 Tom. 3 1952: 63–127 14 pls. [English and French summaries.]

Brand, T. von. Anaerobiosis in Australorbis glabratus: Temperature effects of tissue hydration. J. Wash. Acad. Sci. 45 1955: 373–377 3 figs. tables & graph.

Brand, T. von. see Newton, W. L.

Brander, T. Über die Bisamratte, Ondatra z. zibethica (L.), als Vernichter von Najaden. Arch. Hydrobiol. 50 1955: 92–103 1 fig.

Brander, T. (1). Weitere Beobachtungen zur Muscheldiät der Bisamratte. Arch. Hydrobiol. 51 1955: 261–266 2 pls.

Brandtner, F. Kamegg, eine Freilandstation des späteren Paläolithikums in Niederösterreich. Mitt. prähist. Komm. Wien 7 1955: 1–93 20 pls.

Braucourt, J. de H. Deuxième série d'observations stratigraphiques au Kruisschans. Coupes de l'Écluse Baudouin. 2. Conclusions. Bull. Inst. Sci. nat. Belg. 31 67 1955: 1-14.

Braucourt, J. de H. (1). Deuxième série d'observations stratigraphiques au Kruisschans. Coupes de l'Ecluse Baudouin. 1. Analyse stratigraphique. Bull. Inst. Sci. nat. Belg. 31 66 1955: 1–29 3 pls. 2 figs.

Braucourt, J. de H. see Glibert, M.

Braun, A. F. New light on the biolog Δ of *Spirula*, a mesopelagic cephalopod. Essays in the nat. Sciences in Honor of Capt. A. Hancock, Los Angeles, 1955: 61-72 2 pls.

Braun, R. Zum Lichtsinn facettenaugentragender Muscheln. Zool. Jb. (Allg. Zool.) 65 1954: 91–125 4 figs.

Bricchi, C. Revisione della fauna Neocretacica della Libia: Fam. Pleurotomariidae, Calyptraeidae, Turritellidae, Omalaxidae, Cerithiidae, Aporrhaidae, Strombidae, Volutidae, Ringiculidae. Ann. Mus. libico Stor. nat. 4 1953: 169–194 2 pls.

Brookes, A. E. Distributional notes on some of our small terrestrial mollusca. Bull. Auckland Mus. Conch. Sect. No. 9 1953: 1–2.

Brown, M. J. Some notes on the activities of the spotted golden slug. J. Tenn. Acad. Sci. 30 2 1955: 133.

Bruggen, A. C. v. Lymnaea columella Say und Zonitoides arboreus (Say) in den Gewächshäusern das Basler Botanischen Gartens (Moll., Gastr., Pulm.). Verh. naturf. Ges. Basel 66 1 1955: 1-4 2 figs. Brunner, G. Die Höhle am Butzmannsacker bei Auerbach (Opf.). Geol. Bl. NO-Bayern 5 3 1955: 109–120 textfigs. 1–4.

Brunnschweiler, R. O. Mesozoic stratigraphy and history of the Canning Desert and Fitzroy Valley, Western Australia. J. geol. Soc. Aust. 1 1954: 35–53 1 map.

Brunson, R. B. see Pilsbry, H. A.

Burch, J. B. Some ecological factors of the soil affecting land snails in Virginia. Nautilus Philad. 69 1955: 62–69 graphs.

Burch, J. B. (1). The land snails of Hanover, Henrico, and Chesterfield Counties, Virginia. Virginia J. Sci. N.S. 6 3 1955: 154-162 map.

Burch, J. Q. A systematic outline of the Muricacea in the Eastern Pacific. Mins. Conch. Club S. Calif. **149** 1955: 3-13.

[Burch, J. Q.] (1). Conus ermineus Born. and Conus lithoglyphus Reeve. Mins. Conch. Club S. Calif. 152 1955: 1-2.

Burch, J. Q. (2). Resurrection of Ancient Names. Mins. Conch. Club S. Calif. 152 1955: 2-3.

Bushinskii, G. I. Lithology of the Cretaceous deposits of the mouth of the Dnieper-Don rivers. Trav. Inst. géol. S.S.S.R. 156 (G.S. 67) 1954: 1–306 text-figs. 121–131.

Butler, P. A. Summary of our knowledge of the oyster in the Gulf of Mexico. Fish. Bull. U.S. 55 1954: 479-489.

Butot, L. J. M. Shells from Komodo Island. Penggemar Alam 34 3-4 1954: 105-107 3 figs.

Butot, L. J. M. (1). Over een zoetwater roofslak. Anantome helena (v.d. B.) Gastr.-Prosobr. Penggemar Alam 34 3-4 1954: 108-115 2 figs.

Butot, L. J. M. (2). The mollusc fauna of Pulau Panaitan (Prinseneiland). Land and freshwater molluscs. Treubia 23 1955: 69–135 33 figs. & map.

Büttner, K. Die Molluskenfauna von Südwestsachsen. Abh. Mus. Tierk. Dresden 22 1954: 47–87.

- Cabo, F. L. Notas anatómicas sobre especies españolas de moluscos de interés pesquero. Bol. Soc. esp. Hist. nat. 50 2 Biol. 1952: 255–278 6 pls.
- Cabo, F. L. (1). Una campana de prospeccion pesquera en Mar Menor (Murcia). Bol. Inst. esp. Oceanogr. 66 1954: 1-34.
 - Colaço, F. see Fraga de Azevedo, J. Cameron, C. see Trussell, P.
- Campbell, J. D. The Oretian stage of the New Zealand Triassic system. Trans. roy. Soc. N.Z. 82 5 1955: 1033– 1047 1 pl. 4 figs.
- Carcelles, A. R. Nuevas especies de gastropodos marinos de las Republicas oriental del Uruguay y Argentina. Comun. zool. Mus. Montevideo 4 70 1953: 1-16 5 pls.
- Carcelles, A. R. (1). Especies Sudamericanos de *Argobuccinum* Bruguière 1792. Comun. Inst. Invest. Mus. argent. Cienc. nat. Zool. 2 15 1954: 243-254 1 pl.
- Carcelles, A. R. (2). Observaciones sobre los generos Nucella Chorus y Concholepas. Comun. Inst. Invest. Mus. argent. Cienc. nat. Zool. 2 16 1954: 255-275 5 pls.
- Casey, R. The Pelecypod family Corbiculidae in the Mesozoic of Europe and the Near East. J. Wash. Acad. Sci. 45 1955: 366-372 6 figs.
- Casey, R. (1). The Neomiodontidae. A new family of the Arcticacea (Pelecypoda). Proc. Malac. Soc. London 31 1955: 208-222 3 text-figs. & pl. 11.
- Casey, R. (2). Proposed use of the plenary powers to designate (i) a neotype for the nominal species Ammonites mammillatus Schlotheim, 1813 and (ii) a type species for the genus Douvilliceras De Grossouvre, 1893 (Class Cephalopoda, Order Ammonoidea). Bull. zool. Nom. 9 1954: 250–254.
- Casey, R. & Wright, C. W. Proposed use of the plenary powers to vary the type species of the genus *Cheloniceras* Hyatt, 1903 (Class Cephalopoda, Order Ammonoidea), in order to validate existing nomenclatorial practice. Bull. zool. Nom. 9 1954: 278–280.
- Cassie, R. M. Population studies on the toheroa, Amphidesma ventricosum Gray (Eulamellibranchiata). Aust. J. mar. freshw. Res. 6 1955: 348-391 2 pls. 18 figs.

- Castellanos, Z. J. A. de. Los quitones del Museo de la Ciudad Eva Perón. Notas Mus. Eva Perón 16 1953: 173– 183 6 pls. figs.
- Castro, J. O. de. Glossario de nomes des peixes, Lisboa (Gabinete de Estudos das Pescas) 1954 pp. 1–249. [Mollusca pp. 39–109.]
- Cate, J. & C. Some *Phyllonotus* of the Gulf of California. Mins. Conch. Club S. Calif. **145** 1955: 2-6.
- Cate, J. Ten. Contribution à la question de l'innervation cholinergique du coeur de l'*Anodonta cygnea*. Pubbl. Staz. zool. Napoli 27 1955: 199–203 figs. 1, 2.
- Cate, J. T. & Raesinck, M. J. The action of acetylcholine and eserine on the heart and the intestine of *Anodonta cygnea* L. Physiol. comp. 3 1954: 337–342 6 figs.
- Cather, J. N. A serological technique for gastropod taxonomy. Field & Lab. 23 1 1955: 5-11 7 figs.
 - Chandler, A. C. see Ray, S. M.
- Chang, Hsi, Ch'i Chung-yen & Li, Chien-min. Marine molluscs of North China, Peking (Acad. Sinica, Marine Biol. Section) 1955, pp. ii, 5, iii, 93, 35 pls., text illust. [In Japanese.]
- Chapman, G. Aspects of the fauna and flora of the Azores: The density of animal life in the coralline algal zone. Ann. Mag. nat. Hist. (12) 8 1955: 801-805.
- Charalambakis, S. Contribution à la connaissance du Néogène de l'Attique. Ann. géol. hellen. 1 4 1952 : 1-156 pls. 1-11 text-figs. 1-4. [French résumé.]
 - Ch'i Chung-yen see Chang, Hsi.
- Chiba, T., Touruta, A. & Inque, Y. Studies on the food bivalves in Setonaikai, Inland Sea of Japan. 1. Fulvia mutica Reeve. J. Shimonoseki Coll. Fish. 4 1955: 321-326 1 fig.
- Chipman, W. A. & Hopkins, I. G. Water filtration by the bay scallop, *Pecten irradians*, as observed with the use of radioactive plankton. Biol. Bull. Woods Hole 107 1954: 80-91 1 fig.
 - Chopra, H. C. see Nath, V.
- Christiansen, M. E. The life history of Lepidopleurus asellus (Spengler). Nytt Mag. Zool. 2 1954: 52-72 31 figs.

- Chubb, L. J. A revision of Whitfield's type specimens of the Rudist Mollusks from the Cretaceous of Jamaica, British West Indies. Amer. Mus. Novit. No. 1713 1955: 1–15.
- Chubb, L. J. (1). The Cretaceous succession in Jamaica. Geol. Mag. 92 3 1955: 177-195 text-fig. 1.
- Clench, W. J. A new Murex from Matanzas, Cuba. Breviora 44 1955: 1-2 3 figs.
- Clench, W. J. (1). Setaepoma, a new genus in the Synceridae from the Solomon Islands. Nautilus, Philad. 68 1955: 134.
- Clench, W. J. (2). Supplement to the Catalogue of the genus *Liguus*. Occ. Pprs. on Mollusks, Cambridge, Mass. 1 18 1954: 442-444.
- Clench, W. J. & Turner, R. D. Supplement to the Catalogue of the Family Truncatellidae. Occ. Pprs. on Mollusks, Cambridge, Mass. 1 18 1954: 445.
- Clench, W. J. & Turner, R. D. (1). Supplement to Western Atlantic marine mollusks described by C. B. Adams (Occ. Pprs. No. 15). Occ. Pprs. on Mollusks, Cambridge, Mass. 1 18 1954: 447.
 - Colaço, A. T. F. see Feijó Colaço, A. T.
- Cole, H. A. Purification of Oysters in simple pits. Fishery Investigations (2) 18 No. 5 1954 12 pp. pls. i-iv.
- Collinson, C. A new species of Endolobus (Nautiloidea) from the Clore limestone (Chester) of western Kentucky. J. Paleont. 29 1 1955: 178-181 text-figs. 1-3.
- Collinson, C. W. (1). Mississippian prolecanitid goniatites from Illinois and adjacent states. J. Paleont. 29 3 1955: 433-438 pl. 45 text-figs. 1-2.
 - Colman, J. S. see Mason, J.
- Conde, J. A. Una carta inedita del Doctor Carlos de la Torre sobre el caracol pintado. Mem. Soc. cubana Hist. nat. 22 2 1955: 97–109 4 figs.
 - Cotton, B. C. see Crocker, R. L.
- Cox, L. R. Proposed determination of the nominal species to be accepted as the type species of the genus "Inoceramus" Sowerby (J.), 1814 (Class Pelecypoda) and proposed addition of that name to the "Official List of Generic Names in Zoology." Bull. Zool. Nom. 11 1955: 239-245.

- Cox, L. R. (1). On the affinities of Trigonioides and Hoffetrigonia. Geol. Mag. 92 4 1955: 345-349 text-figs. 1 A-D.
- Cox, L. R. (2). Observations on gastropod descriptive terminology. Proc. Malac. Soc. London 31 1955: 190-202 9 text-figs.
- Cox, L. R. (3). Supplementary application concerning the suppression of works by S. A. Renier (1804 and 1807). Bull, zool. Nom. 9 1954: 265.
- Crampton, H. E. New species of land snails of the genus *Partula* from Raiatea, Society Islands. Amer. Mus. Novit. No. 1761 1956: 1–17 3 figs.
- Crocker, R. L. Notes on a recent raised beach at Point Brown, Yorke Peninsula, South Australia. Trans. roy. Soc. S. Aust. 70 1946: 108-109 2 pls. 1 fig.
- Crocker, R. L. & Cotton, B. C. Some raised beaches of the Lower South-east of South Australia and their significance. Trans. roy. Soc. S. Aust. 70 1946: 64–82 8 pls. 2 figs. map.
- Crusafont Pairó, M. & Comella, J. F. de V. El Burdigaliense continental de la cuenca de Vallés-Penedés. Parte paleontologica. Mem. Comun. Inst. geol. Barcelona 12 1955: 73-272. [French summary.]
- Cumano, H. B. Que animal está aqui enterrado? Naturália 5 1 1955: 22–27 7 figs.
- Curry, D. The occurrence of the dibranchiate Cephalopods Vasseuria and Belosepiella in the English Eocene, with notes on their structure. Proc. Malac. Soc. London 31 1955: 111-122 text-figs.
- Curtis, Jr., N. M. Paleoecology of the Viesca member of the Weches formation at Smithville, Texas. J. Paleont. 29 2 1955: 263–282 pls. 30–31 text-figs. 1–5.
- Dalloni, M. Sur la faune néoconienne du Tell algérien et la présence en Afrique du Nord des Ammonites de la famille des Oostérellidés. C.R. Soc. géol. Fr. 4 1954 : 88–89.
- Dan, J. C. & Wada, S. K. Studies on the acrosome. IV. The acrosome reaction in some bivalve spermatozoa. Biol. Bull. Woods Hole 109 1955: 40– 55 33 figs.

- Dartevelle, E. & Roger, J. Contribution a la connaissance de la faune du Miocène de l'Angola. Comun. Serv. geol. Portug. 35 1954: 227-312 pls. 1-5 text-figs. 3-4 1 map.
- David, L. & Jodot, P. Quelques précisions sur la transgression miocène dans les environs de Gambetta (Est-Constantinois). Bull. Soc. Hist. nat. Afr. N. 46 3-4 1955: 121-131 1 pl.
- Davis, A. G. Truncatellina cylindrica britannica (Pilsbry) in Dorset and Isle of Wight. J. Conch. London 24 2 1955: 61-62.
- Davis, A. G. (1). Acanthinula lamellata (Jeffreys) in a holocene tufa at Totland, Isle of Wight. J. Conch. London 24 3 1955: 40.
- Davis, H. C. Mortality of Olympia oysters at low temperature. Biol. Bull. Woods Hole 109 1955: 404-406.
- Dawley, C. Minnesota land snails. Nautilus, Philad. 69 1955: 56-62.
 - Dawood, M. M. see Dias, E.
- Dawson, C. E. A study of the oyster biology and hydrography at Crystal River, Florida. Publ. Inst. Mar. Sci. Univ. Tex. 4 No. 1 1955: 279-302 2 pls. 6 figs.
- Deevey, E. S., Jr. Paleolimnology of the upper swamp deposit, Pyramid Valley. Rec. Canterbury [N.Z.] Mus. 6 1955: 291-344 figs.
- Dehnel, P. A. Rates of growth of gastropods as a function of latitude. Physiol. Zoöl. 28 1955: 115–144 5 pls. 6 figs.
- Delga, M. D. A propos de "Bochianites" superstes Perv.: remarques sur les Ammonites droites du Crétacé inférieur. C.R. Soc. géol. Fr. 7 1954: 134-137 text-figs. A-C.
- Dell, R. K. The Suter Collection of land and fresh-water Mollusca, Bull. Auckland Mus. Conch. Sect. No. 10 1954: 6-8.
- **Dell, R. K.** (1). Deep-water rock faunas in New Zealand. N.Z. J. Sci. **Tech.** 36B 2 1954: 123–128.
- Dell, R. K. (2). The land Mollusca of the Treasury Islands, Solomon Islands, Pacif. Sci. 9 4 1955: 423–429 2 figs.
- Dell, R. K. (3). The land Mollusca of Nissan Island, Solomon Islands. Pacif. Sci. 9 3 1955: 324–331 2 figs.

- Dell, R. K. (4). A new tornatellinid land mollusk from the Solomon Islands. Pacif. Sci. 9 3 1955: 357-358 1 fig.
- Dell, R. K. (5). The land Mollusca of Fiordland, South-west Otago. Trans. roy. Soc. N.Z. 82 1955: 1135-1148 15 figs.
- Dell, R. K. (6). A synopsis of the Nuculanidae with check lists of the Australian Tertiary and recent species. Rec. Dom. Mus., Wellington 2 3 1956: 123-134.
- Dell, R. K. (7). Some New Zealand subfossil land Mollusca. Rec. Dom. Mus., Wellington 2 3 1956: 135–144 1 map 1 fig.
- Dell, R. K. (8). A contribution to the study of rates of growth in *Paryphanta busbyi* (Gray), (Mollusca, Pulmonata). Rec. Dom. Mus., Wellington 2 3 1956: 145–146 2 figs.
- Dell, R. K. (9). Nature in New Zealand. Native shells. Wellington, N.Z. (A. H. & A. W. Reed) 1955 pp. 64 text-figs.
 - Desai, B. N. see George, J. C.
 - Deslandes, N. see Paraense, W. L.
- De Villafranca, G. W. Adenosine-triphosphatase activity of squid mantle muscle (*Loligo pealii*). Biol. Bull. Woods Hole 108 1955: 113–119 3 figs.
 - De Witt, see Witt, de.
- Dew, B. & Wood, E. J. F. Observations on periodicity in marine invertebrates. Aust. J. mar. freshw. Res. 6 3 1955: 469–478 8 figs.
- Dexter, R. W. Extension of known range of the slug *Pallifera fosteri* F. C. Baker. Nautilus, Philad. **69** 1955: 35–36.
- Dias, E. Nota prévia sôbre ensaios de combate aos planorbideos por métodos biológicos e bioquímicos. Mem. Inst. Osw. Cruz 52 1954: 247-252 11 figs.
- Dias, E. (1). Guerra bacteriológica contra os hospedeiras intermediários da esquistossomose humana. Mem. Inst. Osw. Cruz 52 1955: 315-327 7 figs. [English summary.]
- Dias, E. & Dawood, M. M. Preliminary trials on the biological snail control with *Bacillus pinottii* in Egypt. Mem. Inst. Osw. Cruz 53 1955: 13-29 10 figs.

12 [1955]

Dickie, L. M. Fluctuations in abundance of the giant scallop, Placopecten magellanicus (Gmelin) in the Digby area of the Bay of Fundy. J. Fish. Res. Bd. Can. 12 1955: 797-857 16 figs.

Dickinson, P. Field notes for a freshwater naturalist. Dom. Mus. Handb. No. 3 1951: 1-32 figs.

Dieuzeide, R. Recherches sur les fonds chalutables de la région d'Alger. I.—Introduction. Draguages et chalutages. Notes faunistiques sur la zone meso-abyssale. [Moll. pp. 58-72.] Bull. Sta. Aquic. Pêche Castiglione (N.S.) 7 1955 : 7-86 15 pls.

Dingle, A. N. & Nash, C. B. Acclimatization? Science 121 1955: 110.

Dittmar, H. Ein Sauerlandbach. Untersuchungen an einen Wiesen-Mittelgebirgsbach. Arch. Hydrobiol. **50** 1955 : 305–552 4 pls. 24 figs.

Dodge, H. A historical review of the mollusks of Linnaeus. Part 3. Bull. Amer. Mus. nat. Hist. 107 1955: 1-158.

Dopita, M. & Růžička, B. Proposal for a new method of describing fossil Lamellibranchiata. Přírod. Sborn. Ostravsk. Kraje 14 1-2 1953: 32-41 7 figs. [English summary.]

Donovan, D. T. see Arkell, W. J.

Dow, R. L. & Wallace, D. E. Natural redistribution of a quahog [Venus mercenaria population. Science 122 1955:641-642.

Drăghindă, I. see Filipescu, M. G.

Dral, A. D. G. see Tammes, P. M. L.

Draycot, W. McK. The fossil Trigonia. Victoria Nat. 11 6 1954: 62-63 1 fig.

Dubar, G. Gisements liasiques de la Dorsale tunisienne (Djebel Bent-Saïdane et Dj. Zaghouan). C.R. Soc. geol. Fr. **15** 1953 : 354-356.

Duke, B. O. L. see McCullough, F. S.

Dundee, D. S. Additional localities for Land Mollusca in Oklahoma. Nautilus, Philad. 69 1955: 16-18.

Dundee, D. S. see Schalie, H. v. d.

Dulzetto, F. & Labruto, G. Ricerche biochimiche sulle capsule ovifere di *Murex trunculus* L. Atti Accad. Gioenia (6) **7** 1951 : 211–219 1 fig. [English summary.]

Dvořák, J. A contribution to the knowledge of the Cretaceous in the vicinity of Velký Dřevíč (North of Hronov). Sborn. geol. Ust. čsl. 21 1 1954 369-393 (1-25) pl. 1-4. [English summary.]

Eichman, C. J. A new Cretaceous Emarginula. Nautilus, Philad. 68 1955: 133–134 pl. 4 figs. 7–8.

Einsele, G. & Mosebach, R. Zur Petrographie, Fossilerhaltung und Entstehung der Gesteine des Posidonienschiefers im Schwäbischen Jura. Neues Jb. Min. Geol. Paläont. 101 3 1955: 319-430 pls. 36-40 text-figs. & tables.

Elgood, J. H. see Webb, J. E.

Ellis, A. E. Proposed validation under the plenary powers of the generic name Bithynia Leach 1818. Bull. zool. Nom. **11** 1955 : 275–278.

[Ellis, A. E., editor.] Mollusca in Hertfordshire. List by L. W. Stratton. [Note on issue in Trans. Herts Nat. Hist.] J. Conch. London 24 1955: 58.

Emberger, J. (1). Présence d'Agria marticensis d'Orb. et de Praeradiolites pulchellus Vidal dans le terrains crétacés du bassin du Hodna (Algérie). C.R. Soc. géol. Fr. 1 1954: 48-50 figs. 1-2.

Emberger, J. (2). Découverte de Jurassique supérieur à Ammonites dans l'Atlas saharien (Algérie). C.R. Soc. géol. Fr. 15 1954 : 383-385.

Emerson, W. K. Notes on the Scaphopod Mollusks: rectifications of Notes on the nomenclature. [Eleven new names.] Proc. biol. Soc. Wash. 67 1954: 183-

Emerson, W. K. see Puffer, E. L.

Emura, S. see Komai, T.

Engle, J. see Loosanoff, V.

Eyles, J. P. Field notes on the excavations [at Pyramid Swamp, Canterbury, New Zealand]. Rec. Canterbury [N.Z.] Mus. 6 4 1955: 257-260 4 pls. 1 fig.

Fänge, R. Use of the isolated heart of a freshwater mussel (Anodonta cygnea L.) for biological estimation of 5-hydroxytryptamine. Experientia 11 4 1955 : 156.

Faro, M. da C. see Fraga de Azevedo.

Feijó Colaço, A. T. see Fraga de Azevedo.

- Ferreira, O. da V. Os pectinídeos do Miocénico da Ilha de Santa Maria (Açores). Rev. Fac. Cienc. Lisboa (2C) 2 2 1952: 243-258 4 pls. map.
- Ferreira, O. da V. (1). Pectinídeos do Miocénico do Vale do Sado e da Serra da Arrábida. Comun. Serv. geol. Portug. 35 1954: 155-192 pls. 1-8 2 tables.

Ferreri, E. see Guardabassi, A.

- Feugueur, L. Essai de synchronisation entre les assises saumâtres du Thanétien-Landénien (Gand-Ostende) et du Sparnacien (Ile-de-France). Bull. Soc. belge Géol. Pal. Hydr. 64 1 1955: 67–92 pls. 1–4 1 geol. section.
- Filipescu, M. G., Dräghindä, I. & Mutihac, V. Contributions à l'établissement des horizons et de l'âge des Schistes Noirs de la zone moyenne du Flysch des Carpathes orientales. Rev. Biol., Bucureşti 2 1955: 101-107 13 figs.
- Finks, R. M. Conularia in a sponge from the west Texas Permian. J. Paleont. 29 5 1955: 831-836 pl. 82.
- Fischer, P.-H. Valor de las conchas en las primeras civilaciones. Circ. Mus. biblio. zool. Habana 1954: 1097–1098
- Fischer, P.-H. (1). Gastéropodes excavateurs de roches littorales. J. Conchyliol. Paris 95 1955: 23-27.
- Fischer, P.-H. (2). Remarques sur la famille des Calyptraeidae. J. Conchyliol. Paris 95 1955: 38-42.
- Fischer, P.-H. (3). Fluorescence des oothèques chez quelques Gasteropodes. J. Conchyliol. Paris 95 1955: 86-88.
- Fischer, P.-H. (4). Gastéropodes littoraux fossilles. J. Conchyliol. 95 1955: 108-112.
- Fischer, P.-H. (6). Opercules de Gastéropodes examinés en lumière de Wood, J. Conchyliol. Paris 95 1955: 113-114.
- Fitch, J. E. Results of the 1954 Pismo clam census. Calif. Fish Game 41 1955: 209-211.
- Fitch, J. E. (1). The Pismo clam in 1952 and 1953. Calif. Fish Game 40 1954: 199-201.
- Fleming, C. A. A new subspecies of scallop from Byron Bay, New South Wales. Aust. Zool. 12 1955: 108-109 l fig.

- Fleming, C. A. (1). The fossil fauna of the Campbell Island breceias. Cape Exped. Ser. Bull. No. 3 1950: 47-62.
- Fleming, C. A. (2). Collections of Mollusca in the New Zealand geological survey. Bull. Auckland Mus. Conch. Sect. No. 9 1953: 6-8.
- Fleming, C. A. (3). Notes on the Suter collection of New Zealand Mollusca—1. Bull. Auckland Mus. Conch. Sect. No. 10 1954: 3–4.
- Fleming, C. A. (4). Kapitean (Upper Miocene) Mollusca from the Te Waewae Bay, Southland, New Zealand. Trans. roy. Soc. N.Z. 82 5 1955: 1049–1059 2 pls. 1 fig.
 - Fleming, C. A. see Wellman, H. H.
- Flower, R. H. Saltations in nautiloid coiling. Evolution 9 1955: 244-260 3 figs.
- Flower, R. H. (1). Status of endoceroid classification. J. Paleont. 29 3 1955: 329-371 pls. 32-35 text-figs. 1-6.
- **Flower, R. H.** (2). New Chazyan Orthocones. J. Paleont. **29** 1955: 806–830 pls. 77–81 1 text-fig.
- Flower, R. H. (3). Trails and tentacular impressions of orthoconic cephalopods. J. Paleont. 29 5 1955: 857–867 text-figs. 1–4.
- Fonseca Sacarrão, G. da see Sacarrão, G. da F.
- Forcart, L. Revision von Vitrina kochi Andreae und Vitrina nivalis (Dumont & Mortillet). (Moll., Pulm.). Verh. naturf. Ges. Basel 65 2 1954: 264–269.
- Forcart, L. (1). Veronicellidae de l'ile Fernando Po. Ann. Mus. Congo belg. N.S. 4to Zool. 1 1954: 193-.
- Forcart, L. (2). Die nordischen Arten der gattung *Vitrina*. Arch. Molluskenk. 84 1955: 155–166 pl. 12.
- Forest, J. Beautés du fond des mers, Paris (Larousse) [Moll., pls. 85–97] 1955 pp. 100, frontis., text illust. col.
- Fox, H. M. The effect of oxygen on the concentration of haem in invertebrates. Proc. roy. Soc. 143B 911 1955; 203-214.
- Fox, H. M. & Taylor, A. E. R. The tolerance of oxygen by aquatic invertebrates. Proc. roy. Soc. 143B 911 1954: 214-225.

Fraga de Azevedo, J., Costa, M. M. da & Gomes, F. A. C. A susceptibilidadae do Planorbis metidjensis ao Schistosoma haematobium da Guiné Portuguesa e ao Schistosoma mansoni de Moçambique. An. Inst. Med. Trop. Lisboa 11 2 1954: 251-260. [English & French summaries.]

Fraga de Azevedo, J., Feijó Colaço, A. T. & Faro, M. da C. As bilharzioses humanas no sul do Save (Moçambique). An. Inst. Med. Trop. Lisboa 11 1 1954: 5-120 110 figs. [English & French summaries.]

Fraga de Azevedo, J. & Medeiros, L. C. M. de. A identificação do *Planorbis metidjensis*, Forbes, do Algarve, pela morfologia dos órgãos genitais e rádula, sua comparação com os mesmos elementos do *Planorbis corneus*. An. Inst. Med. Trop. Lisboa 11 2 1954: 231-240 9 figs. [English & French summaries.]

Franciscolo, M. E. Fauna cavernicola del Savonese. [Mollusca pp. 169–170.] Ann. Mus. Stor. nat. Genova 67 1955: 1–223.

Franzén, A. Investigations into spermiogenesis and sperm morphology among invertebrates. Uppsala (Almqvist & Wiksells) 1956 pp. 13.

Fretter, V. Observations on Balcis devians (Monterosato) and Balcis alba (Da Costa). Proc. Malac. Soc. London 31 1955: 137-144 3 text-figs.

Fretter, V. (1). Some observations on Tricolia pullus (L.) and Margarites helicinus (Fabricius). Proc. Malac. Soc. London 31 1955: 159-162 text-fig.

Fretter, V. & Graham, A. Observations on the opisthobranch molluse Acteon tornatilis (L.). J. Mar. biol. Ass. U.K. 33 1954: 565-585 9 figs.

Freyberg, B. v. Eine unterpliocäne Fauna von Elaea (Lakonien, Peloppones). Ann. géol. hellen. 1 1 1947: 120-125 1 table.

Frömming, E. Ein Beitrag zur Frage des Kalkstoffwechsels unserer Süsswasserschnecken. Z. Naturf. 8B 5 1953: 258–263 2 figs.

Frömming, E. (1). Gehören die ätherischen Öle zu den Pflanzeninhaltsstoffen, welche auf Tiere eine abschreckende Wirkung ausüben. Mitt. Berl. Malak. Nr. 6 1954; 56–60.

Frömming, E. (2). Die Rolle unserer Schnecken bei der Ausbreitung von infektiösen Pflanzenkrankheiten. I. Faule-Krankheiten bei Erntegut. Z. angew. Zool. 1955: 433–439.

Frömming, E. (3). Okologische Einflüsse auf die postembryonale Entwicklung der gehäusetragenden Landlungenschnecke Eobania vermiculata (O. F. Müll.). Zool. Jb. Syst. 83 1955: 503-510 4 figs.

Frömming, E. & Plate, H. P. Über das Verhalten pflanzenfressender Schnecken zu unseren wirtschaftlich wichtigen Gräsern. Z. angew. Zool. 1954: 71–75.

Frömming, E. see Plate, H. P.

Fuhrmann, A. Petrographie, Fauna und stratigraphische Stellung einiger Aufschlüsse im Oberharger Oberdevon, Blatt Zellerfeld und Riefensbeek. Geol. Jb. 69 1955: 629-652 text-figs. 1-5.

Fuji, A. Some ecological aspects of Zyusan-gata with special reference to the habitats of *Corbicula japonica* group. Bull. Fac. Fish. Hokkaido Univ. 5 1955: 313-319 8 figs.

Fulton, C. see Trussell, P.

Furnish, W. M. (1), Miller, A. K. (1) & Youngquist, W. Discovery of the early Mississippian goniatite *Protocanites* in northern Nevada. J. Paleont. 29 1 1955: 186 fig. 1.

Furnish, W. M. see Miller, A. K.

Furnish, W. M. see Glenister, B. F.

Gabe, M. & Prenant, M. Sur une nouvelle espèce d'*Oncidiella*. Bull. Inst. franç. Afr. noire 17A 1 1955: 84-91 2 figs.

Gabe, M. & Prenant, M. (1). Particularités anatomiques de la gaine radulaire chez quelques Tectibranches. [Abstract.] Bull. Soc. zool. Fr. 78 1954: 338-339.

Galkin, Y. I. Gastropod Mollusca (Trochidae) of the Far-Eastern and Northern Seas of the U.S.S.R. Tabl. anal. Faune URSS 57 1955: 1-131 90 figs. [In Russian.]

Ganss, O. & Schmidt-Thomé, P. Die gefaltete Molasse am Alpenrand zwischen Bodensee und Salzach. L. dtsch. geol. Ges. 105 3 1955: 402–495 text-figs. 1–8.

- Gardner, E. N. & N. W. Some observations on the Fijian snail *Placostylus* (Callistocharis) gracilis (Brod.). Bull. Auckland Mus. Conch. Sect. No. 9 1953: 9-11.
- Gardner, N. W. An extension of the range of *Paryphanta watti* Powell. Bull. Auckland Mus. Conch. Sect. No. 9 1953: 12.
- Gardner, N. W. (1). Liarea turriculata waipoua Powell. Bull. Auckland Mus. Conch. Sect. No. 11 1955: 23.
- Gardner, N. W. (2). Some observations on Serpho matthewsi Suter. Bull. Auckland Mus. Conch. Sect. No. 11 1955: 21-22.
 - Garner, H. F. see Miller, A. K.
- Geib, K. W. Über den Vorgang der Konkretionsbildung bei den Barytkonkretionen des mitteloligozänen Meeressandes von Steinhardt (Kreis Kreuznach). Notizbl. hess. Landesamt, Bodenforsch. 83 1955: 243–245 7 figs.
- Genesi, M. Localizzazione delle fosfatasi acida ed alcalina e di sostanze Schiff-positive in Lamellibranchi e Gasteropodi marini (Mytilus, Patella e Fissurella). Riv. Biol. N.S. 47 1 1955: 97-106 3 pls. [English summary.]
- Genovese, S. Su un particolare caso di inquilinismo riscontrato in *Mytilus gallo provincialis* Lamk. Boll. Zool. 22 1955: 117-120 1 pl.
- George, J. C. & Desai, B. N. On the liver fat in *Pila globosa* (Swainson). J. Anim. Morph. Physiol. 1 1954: 56-58.
- Giglioli, M. E. C. The egg masses of the Naticidae. (Gastropoda.) J. Fish. Res. Bd. Can. 12 1955: 287-327 13 figs.
- Gill, E. D. & Baker, A. A. Tertiary fossils at Aberfeldie, Melbourne. Vict. Nat., Melb. 72 3 1955: 39-43 6 figs.
- Givulescu, R. Contributions à l'étude du crétacé supérieur du Bassin de Borod (la région d'Oradéa). Stud. Cercet. Acad. romîne. Fil. Cluj 5 1-2 1954: 173-218 10 figs. map. [French résumé.]
- Glaessner, M. F. Pelagic fossils (Aturia, Penguins, Whales) from the Tertiary of South Australia. Rec. S. Aust. Mus. 11 4 1955: 353-372 3 pls. 7 figs.

- Glenister, B. F., Miller, A. K. & Furnish, W. M. Upper Cretaceous and early Tertiary nautiloids from Western Australia. J. Paleont. 30 3 1956: 492-503 4 pls. 7 text-figs.
- Glibert, M. Quelques espèces nouvelles ou mal connues de l'Oligocene moyen et supérieur de la Belgique. Bull. Inst. Sci. nat. Belg. 31 86 1955: 1-7.
- Glibert, M. & Heinzelin, J. de B. Le gîte des vertebres tongriens de Hoeleden. Bull. Mus. Hist. nat. Belg. 30 1 1954: 1–14 2 pls. 3 figs.
- Glibert, M. & Heinzelin, J. de B. (1). La faune et l'age Miocène supérieur des Sables de Deurne. Bull. Inst. Sci. nat. Belg. 31 71, 72, 1955: 1-27, 1-12.
- Glinski, A. Über ein Eisen-Ooid-Vorkommen in der Sötenischer Mulde (Devon, Eifel) und seine stratigraphische Bedeutung. Senckenbergiana, Lethaea 36 1955: 235–242 2 figs.
- Goguel, J. A propos de la profondeur de dépôt des marnes à Ammonites pyriteuses. C.R. Soc. géol. Fr. 11 1954 : 222-225.
- Gokhale, S. V. Recovery of a fishery. Curr. Sci. 23 12 1954: 404.
- Goldring, R. The Upper Devonian and Lower Carboniferous trilobites of the Pilton beds in N. Devon, with an appendix on *Goniatites* of the Pilton beds. Senckenbergiana, Lethaea 36 1955: 27-48 2 pls. 7 figs.
- Gomes, F. A. C. see Fraga de Azevedo, J.
- Gonçalves, N. B. & Soares, R. de R. L. Molluscocide action of the cupric ricinoleate. Mem. Inst. Osw. Cruz 53 1955: 397-409 3 figs.
- Gottis, C. Sur un *Tisoa* trés abondant dans le Numidien de Tunisie. Bull. Soc. Sci. nat. Tunis **7** 1954: 183–192 **3** pls.
- Graham, A. Molluscan diets (Presidential address). Proc. Malac. Soc. London 31 1955: 144-159.
 - Graham, A. see Fretter, V.
- Grahle, H.-O. Rezente und holozäne Mollusken aus dem Flussgebiet der Leine bei Hannover. Ber. naturh. Ges. Hannover No. 102 1954: 13-24 1 pl. 2 figs.

Graue, G. A rectification of Peeten nomenclature. Nautilus Philad. 68 1955: 113-115.

Graziadei, P. see Rossi, F.

Greenfield, L. J. Concentration of *Teredo* Cellulase. Proc. Soc. exp. Biol. Med. **89** 2 1955: 241-243.

Greer, B. see Trussell, P.

Grossu, A. V. Gastropoda Pulmonata. Faune Rep. Pop. Romane **3** fasc. 1 1955: 1–518 282 figs.

Guardabassi, A. & Ferreri, E. Istofisiologia dell'apparato digerente di *Helix pomatia*. Arch. 2001. ital. Napoli 38 1954: 61-156 3 pls. [English summary.]

Guérin, S. (1). Contribution á l'étude du Lias de Fay (Ain). Sciences de la Terre, Nancy 2 3 1954 : 3-18 text-figs. 1-2.

Guérin, S. (2). Sur la présence de Lias fossilifère entre Thoirette et Turgon (Jura). C.R. Soc. géol. Fr. 8 1954: 157-159.

Haas, F. Binnenschnecken aus einer Karbon-Landschaft im unteren Amazonasgebiete. Arch. Molluskenk. 84 1955: 101-103 figs. 1-6.

Haas, F. (1). Malakozoologisches aus Otto Hamann's "Europäischer Höhlenfauna." Arch. Molluskenk. 84 1955: 211–212.

Haas, F. (2). Unionella Haas 1913 vorweggenommen durch Unionella Etheridge 1888. Arch. Molluskenk. 84 1955 : 212.

Haas, F. (3). On some small collections of inland shells from South America. Fieldiana, zool. 34 1955: 361-387 figs.

Haas, F. (4). Bivalvia. Bronns Klassen **3** Abt. **3** Teil **2** 1955 : 679–909 **2** figs.

Haas, F. (5). On non-marine shells from northeastern Brazil and Peru. Fieldiana, Zool. 37 1955: 303–337 17 figs.

Haas, F. (6). The Percy Sladen Trust Expedition to Lake Titicaca in 1937. Mollusca: Gastropoda. Trans. Linn. Soc. Lond. (3) 1 1955: 275–308 28 figs.

Haas, O. Eocene Aturia from Cyrenaica. Amer. Mus. Novit. No. 1747 1955: 1-4 3 figs.

Haas, O. (1). Revision of the Jurassic ammonite fauna of Mount Hermon, Syria. Bull. Amer. Mus. nat. Hist. 108 1 1955: 1-210 3 pls. 4 figs.

Haas, 0. (2). A case of parallelism in Upper Cretaceous ammonites. J. Paleont. 29 6 1955: 1070–1071.

Habe, T. Ringiculidae and Retusidae in Japan (Volvulella and Semiretusa not included). Illustrated Catalogues of Japanese Shells No. 2 1950: 7–16 pl. 2.

Habe, T. (1). Hydatinidae, Bullidae and Akeridae in Japan. Illustrated Catalogue of Japanese Shells No. 3 1950: 17-27 2 text-figs. pl. 3.

Habe, T. (2). Myochamidae in Japan. Illustrated Catalogue of Japanese Shells No. 4 1950: 25–30 pl. 4.

Habe, T. (3), Japanese Acmarid Limpets, Mins. Conch. Club S. Calif. 146 1955: 2-3.

Habe, T. (4). Notes on the Systematic position of the genus *Orectospira* Dall, 1925. Mins. Conch. Club S. Calif. 147 1955: 4.

Habe, T. (5). Fauna of Akkeshi Bay. XXI. Pelcypodes and Scaphopoda. Publ. Akkeshi mar. biol. Sta. No. 4 1955: 1-31 7 pls.

Habe, T. (6). Anatomical studies on Japanese land snails (3). Venus, Fukuyama 18 4 1955: 221-234 figs. 1-12.

Habe, T. (7). Notes on three Tellinid species described by V. Bertin. Venus, Fukuyama 18 4 1955: 240-243 text-figs.

Habe, T. (8). Deposit of shell remains in Ofunato Bay. Venus, Fukuyama 18 4 1955: 244-251.

Habe, T. (9). Note on the systematic position of the genus *Orectospira* Dall, 1925. Zool. Mag., Tokyo 64 1955: 259-260. [In Japanese, English résumé.]

Habe, T. (10). Notes on Potamocorbula (gen. nov.) amurensis (Schrenck). Zool. Mag., Tokyo 64 1955: 271–272. [In Japanese & English résumé.]

Habe, T. see Kuroda, T.

Habe, T. see Yamaguchi, N.

Haefelfinger, H. R. Inkretorische Drüsenkomplexe im Gehirn decapoder Cephalopoden. (Untersuchungen an Illex coindeti.) Rev. suisse Zool. 61 1954: 153–162 5 figs. Hand, C. Alderia modesta in Washington. Nautilus, Philad. 69 1955: 72.

Hand, C. & Steinberg, J. On the occurrence of the Nudibranch Alderia modesta (Lovén 1844) on the central Californian Coast. Nautilus, Philad. 69 1955: 22-28 4 text-figs.

Hanna, G. D. see Smith, A. G.

Hano, V. & Seneš; J. Die Untermiozäne fauna bei Rapovce. Geol. sborn. Slovensk. Akad. 3 3-4 1953: 315-354 pls. 43-60 text-figs. 1-0. [Russian and German summaries pp. 354 & 359.]

Hansen, K. Beitrag zur faunistischen Kenntnis des Grunewaldes, Berlin Mitt. Berl. Malak. Nr. 9 1955: 137–139

Harrington, H. J. The Permian Eurydesma fauna of eastern Argentina. J. Paleont. 29 1 1955: 112-128 pls. 23-26.

Haskin, H. H. Age determination in molluses. Trans. N.Y. Acad. Sci. (2) 16 6 1954: 300–304 7 figs. & graphs.

Haskin, H. see Pomeroy, L.

Hasslein, L. Zur Weichtierfauna des Obenburger Mains. Nachr. naturw. Mus. Aschaffenburg No. 45 1954: 1–29 2 pls.

Hayasaka, I. An occurrence of Koninckioceras from the Japanese Permian, Jap. J. Geol. Geogr. 25 1-2 1954: 57-59 1 pl. 1 fig.

Hayasaka, I. (1). On the giant ammonite recently donated to the National Science Museum [Tokyo]. Nat. Sci. & Mus. 22 3-5 1955; 9-11 1 fig. [In Japanese.]

Hayasaka, I. (2). A new Permian species of *Porcellia* from Japan. J. Fac. Sci. Hokkaido Univ. Geol. 4 9 1 1955: 21-24 pl. 1.

Hayasaka, I. & Ozaki, K. A Foordiceras from the Permian formation of Central Japan, Sci. Rep. Kanazaw Univ. 3 1 1955: 183–186 1 pl.

Hedgpeth, J. W. A problem in oyster taxonomy. Syst. Zool. 3 1954: 21-25 4 figs.

Heide, S. van der. Die Frage der praktischen Verwendbarkeit der Süsswasserfauna in der Karbonstratigraphie. Geol. Jb. 69 1955: 625–528.

Heinzelin, J. see Glibert, M.

Heldt, H. Observations sur la ponte chez le Moule (Mytilus galloprovincialis Lmk.) et l'émission des produits sexuels mâles chez le Palourde (Tapes decussatus Forb. et Hly.). Bull. Soc. Sci. nat. Tunis 5 1952: 171–174 1 pl.

Heller, R. Stratigraphy and paleontology of the Roubidoux formation of Missouri. Bienn. Rep. Mo. Bur. Geol. Min. (2) 35 1954: 28–42 pls. 10–13, 16–17.

Hemming, F. On the type species of *Ancilla* Lamarck, 1799 (Class Gastropoda), a genus established with no cited nominal species. Bull. zool. Nom. 9 1954: 219–220.

Hepper, B. T. Environmental factors governing the infection of mussels, *Mytilus edulis*, by *Mytilicola intestinalis*. Fish. Invest. (2) 20 No. 3 1955: 1-21 7 figs.

Herold, H. Potamopyrgus jenkinsi E. A. Smith, in Berlin und Brandenburg. Mitt. Berl. Malak. Nr. 6 1954: 61-65.

Hertlein, L. G. Description of a new pelecypod of the genus Lima from deep water off central California. Proc. Calif. Acad. Sci. (4) 27 12 1952: 377–381 2 figs.

Hertlein, L. G. & Strong, A. M. Marine mollusks collected during the "Askoy" Expedition to Panama, Colombia, and Ecuador in 1941. Bull. Amer. Mus. nat. Hist. 107 2 1955: 159–318 3 pls.

Hertlein, L. G. & Strong, A. M. (1). Marine mollusks collected at the Galapagos Islands during the voyage of the 'Velero III,' 1931-32. Essays in nat. Sciences in Honor of Capt. A. Hancock, Los Angeles 1955; 111-145 l pl.

Hess, O. Der ersten Furchungsteilungen bei *Bithynia tentaculata* L. Naturwissenschaften **42** 12 1955 : 374 1 fig.

Hess, 0. (1). Die entwicklung von Halbkeimen bei *Bithynia tentaculata* L. Naturwissenschaften 42 13 1955; 398 1 fig.

Hess, 0. (2). Exogastrula—Keime bei Bithynia tentaculata. Z. Naturf. 10B 6 1955: 356-357 1 pl. 2 figs.

Hewer, A. M. Snails and reptiles on Betsy Island. Tasm. Nat. 2 2 1951: 7. Hill, H. R. & Tompkins, P. D. Common sea shells of the Los Angeles County Coast. Publ. Los Angeles County Mus. Zool. No. 7 1954: 1–48 text illust.

Hinsch, W. Miozängliederung in den Erdölfelden Heide und Bramstedt. Neues Jb. Geol. Paläont. B. 8 1955: 348–357 1 text-fig.

Hipkins, A. K. Minute gems from seaweeds, under stones and shell sand. Bull. Auckland Mus. Conch. Cl. No. 7 1951: 10-11.

Hipkins, A. K. (1). The genus Scrobs Watson, 1886. Bull. Auckland Mus. Conch. Sect. No. 11 1955: 15-19 9 figs.

Hirase, S. An illustrated handbook of shells in natural colors from the Japanese Islands and adjacent territory. Tokyo (Maruzen Co.) 1954 pp. xxiv [3] & 124, frontis. (port.) 134 pls.

Hirayama, K. Discovery of *Perotrochus* in the Miocene deposits of Tochigi Prefecture, Japan. Venus, Fukuyama 18 4 1955: 266-272 pl. 11.

Hirayama, K. (1). On some Miocene species of *Lucinoma* from Japan, with descriptions of two new species.—Notes on Japanese lucinid molluscs part 1. Jap. J. Geol. Geogr. 25 1-2 1954: 101-115 2 pls.

Hirayama, K. (2). Miocene Mollusca from the Arakawa Group, Tochigi Prefecture, Japan (Part 1). Sci. Rep. Tokyo Kyoika Daig. 3 C 1954: 43-76 3 pls.

Hirayama, K. (3). The Asagai formation and its molluscan fossils in the northern region, Jôban coal-field, Fukushima Prefecture, Japan. Sci. Rep. Tokyo Kyoiku Daig. 4 1955: 49–130 5 pls. 5 figs.

Hiscock, I. D. A note on the life history of the Australian freshwater mussel, *Hyridella australis* Lam. Trans. roy. Soc. S. Aust. **74** 2 1951: 146-148 5 figs.

Hlaváč, V. F. Příspevek k malakologické topografii Č.S.R. Cas. nár. Mus. 123 1 1954: 59-62.

Hoepen, E. C. N. van. New and little-known ammonites from the Albian of Zululand. S. Afr. J. Sci. 51 12 1955: 355-361 6 figs.

Hoepen, E. C. N. van (1). Turonianconiacian ammonites from Zululand. S. Afr. J. Sci. 51 12 1955: 361-377 figs. 1-31. Hoepen, E. C. N. van (2). A new family of keeled ammonites from the Albian of Zululand, S. Afr. J. Sci. 51 12 1955: 377-382 6 figs.

Hoffstetter, R. Moluscos subfosiles de los estanques de sal de Salinas. (Pen. de Santa Elena, Ecuador). Comparacion con la fauna actual del Ecuador. Bol. Inf. cient. nac. Quito 7 64 1954: 303–333 7 figs.

Hoffstetter, R. (1). Moluscos subfosiles de los estanques de sal de Salinas. (Pen. de Santa Elena, Ecuador). Comparacion con la fauna actual del Ecuador. Bol. Inf. cient. nac., Quito 7 65 1954: 399-426 3 figs.

Hoffstetter, R. (2). Moluscos subfosiles de los estanques de sal de Salinas. Rev. Biol. mar. 5 1-3 1955: 132-203.

Hölder, H. Die Ammoniten-Gattung Taramelliceras im Südwestdeutschen Unter- und Mittelmalm. Morphologische und taxionomische Studien an Ammonites flexuosus Buch (Oppeliidae). Palaeontographica A 106 3–6 1955: 37–153 pls. 16–19 text-figs. 1–182.

Hopkins, J. G. see Chipman, W. A. Hopkins, S. H. see Menzel, R. W.

Horibata, T. see Yamazi, I.

Horiguchi, Y. & Miyake, M. Biochemical studies on pearl oyster *Pinctada martensii*—I. Distribution of glucosulfatase, phenolsulfatase and chondrosulfatase in various tissues of pearl oyster. Bull, Jap. Soc. sci. Fish. 19 1954: 957-962.

Hornibrook, N. de B. see Wellman, H. H.

Hornŷ, R. New gastropods from the Silurian of central Bohemia. Sborn. geol. Ust. čsl. 20 1953: 189–212 (1–24) pls. 22–24. [Russian and English summaries.]

Horný, R. (1). Palaeozygopleuridae nov. fam. (Gastropoda) from the Devonian of Central Bohemia. Sborn. geol. Ust. Čsl. 21 ser. palaeont. 1954: 17–160 (1–144) pl. 1–10. [English and Russian summaries.]

Horstmann, H.-J. Untersuchungen zur Physiologie der Begattung und Befruchtung der Schlammschnecke Lymnaea stagnalis L. Z. Morph. Ökol. 44 1955: 222-268 25 figs. Horváth, A. Mollusken-fauna des Fehértó in Szeged. Ann. Biol. Univ. Szeged. 1 1950: 321–326. [German summary.]

Horváth, A. (1). Die Molluskenfauna der Theiss. Acta biol. Szeged N.S. 1 1955: 174–180.

Hotchkiss, N. Wildlife Abstracts 1935–51. An annotated bibliography of the publications abstracted in the Wildlife Review Nos. 1–66. Washington, D.C. (U.S. Fish & Wildlife Service) 1954 pp. iv + 434.

Hubendick, B. On the genus Camptoceras (Moll. Pulm.). Ark, zool. **7** 1955: 451-457 11 figs.

Hubendick, B. (1). A new species of Amerianna from the Philippines with remarks on the spired Planorbidae of the Pacific area. Philipp. J. Sci. 83 1954: 319-323 1 pl. 1 fig.

Hubendick, B. (2). The Percy Sladen Trust Expedition to Lake Titicaca in 1937. The anatomy of the Gastropoda. Trans. Linn. Soc. Lond. (3) 1 1955: 309-327 95 figs.

Hubendick, B. (3). Phylogeny in Planorbidae. Trans. zool. Soc. Lond. 28 1955: 453-542 210 figs.

Hubendick, B. (4). The relationships of the East African Vitrinae with notes on taxonomy of Vitrina. Ark. zool. 6 1954: 83-96 29 figs.

Hubendick, B. (5). On the anatomy of Choanomphalus (Moll. Pulm.). Ark. zool. 6 1954: 503-509 10 figs.

Hubricht, L. Discus macclintocki (F. C. Baker). Nautilus, Philad. 69 1955: 34.

Hudec, V. Dva příspěvky k malakologickému pruzkumu jihovýchodní Moravy. Čas. nár. Mus. 123 2 1954: 223–224 1 fig.

Hunter, W. & M. Russell. Roebuck's papers on Scottish non-marine mollusca. J. Conch. London 24 1955: 59-60.

Huzioka, K. see Nagao, T.

Hynd, J. S. A revision of the Australian pearl-shells, genus *Pinctada* (Lamellibranchia). Aust. J. Mar. Freshw. Res. 6 1955: 98-137 13 pls. 2 figs.

Iacob, D. Contributions à la connaissance de la stratigraphie et de la tectonique de la région Zam-Godinești-Cărmăzănești (Hunedoara). Bul. sti. Acad. Republ. rom. 5 3 1953 : 663–670 5 figs. [French résumé.]

Ichikawa, K. Early Neo-Triassic pelecypods from Iwai, near Itsukaichi, Tokyo Prefecture. Jap. J. Geol. Geogr. 25 1–2 1954: 177–1951 pl.

Ichikawa, K. & Yabe, Y. Eumorphotis multiformis shionosawensis subsp. nov. from the Shinosawa limestone at Shionosawa, north of the Sanchu Graben, Kwanto mountainland, Japan. Trans. palaeont. Soc. Japan n.s. 17 1955: 5–12 pl. 2 1 text-fig.

Ikemoto, N. Action potential in the smooth muscle of the snails and its meaning on the orientation of impulse propagation. Biol. J. Okayama Univ. 2 2-3 1955: 67-74 5 figs.

Ikinaga, K. The metabolism of the pearl oyster *Pinctada martensii* I. On the carbonic anhydrase. Bull. Jap. Soc. sci. Fish. 19 1954: 925-928 1 fig.

Ikenaga, K. see Ojima, Y.

Imai, T. see Ito, S.

Imai, T., Sakai, S., Okada, H. & Yoshida, T. Breeding of the Olympia oyster in tanks and culture experiments in Japanese waters. Tohoku J. agric. Res. 5 1 1954: 13-25 9 figs.

Imlay, R. W. Characteristic Jurassic molluses from northern Alaska. Prof. Pap. U.S. geol. Surv. 274-D 1955: 69-96 pls. 8-13 text-fig. 20 4 tables & 1 chart.

Imlay, R. W. (1). Stratigraphic and geographic range of the late Cretaceous pelecypod *Didynotis*. J. Paleont. 29 3 1955: 548-550.

Incitti, L. see Baroni, C.

Ingua, Y. see Chiba, T. and others.

Iredale, T. Mollusca Part 1. [Names buried in the text are omitted in Record of Great Barrier Reef Expedition, 1928–29.] Sci. Report 5 No. 6 1939.

Iredale, T. (1). Rissoid Sectional Names. [These are termed genera in the text and reference is made to Laseron Rec. Austr. Mus. 22 pp. 257–287 figs. 1–94 1950 for the description of type species.] Proc. Roy. Soc. N.S.W. 1950–1954 Mar. 4 1955 p. 81.

- Iredale, T. (2). Cuttle fish "bones" again. Aust. zool. 12 1954: 63-82 2 pls.
- Iredale, T. (3). Froriep and Lamarck. Aust. zool. 12 1955: 175.
- Iredale, T. (4). On Sepia cultrata Hoyle. Proc. R. zool. Soc. N.S.W. 1953-54 [1955]: 78-79.
- Ishida, S. Metabolic patterns in bivalves. IV. The urea-deaminating activity in *Corbicula*. Sci. Rep. Tôhoku Univ. (4) 20 4 1954: 329–333.
- Ito, S. & Imai, T. Ecology of oyster bed. 1. On the decline of productivity due to repeated cultures. Tohoku J. agric. Res. 5 4 1955: 251-268 7 figs.
- Ito, S. & Leuchtenberger, C. The possible role of the DNA content of spermatozoa for activation process of the eggs of the clam, Spisula solidissima. Chromosoma 7 4 1955: 328-339 6 figs.
- Iyengar, V. K. S. Studies on the reproductive tract of *Ariophanta maheraspatana* (Gray). (Mollusca: Pulmonata.) Proc. nat. Inst. Sci. India 20 6 1954: 683-690 2 pls. 1 fig.
- Jaanusson, V. & Mutvei, H. Stratigraphie und Lithologie der unterordovizischen *Platyurus*-Stufe im Siljan-Gebiet, Dalarna. Bull. geol. Inst. Univ. Upsala 35 1953–55: 7–34 pls. 1–5 text-figs. 1–6.
- Jackiewicz, M. Anatomical comparative investigations on lymnaeids of the genus *Radix* Montfort in Great Poland. Wydz. mat.-przyr. Kom. Biol. 15 3 1954: 1-20 3 figs. [English summary.]
- **Jackson, N.** Albinism in volutes. Proc. R. zool. Soc. N.S.W. **1952–53** 1954: 35–37 2 figs.
- Jacobson, M. K. A new Macroceramus from Puerto Rico. Arch. Molluskenk. 84 1/3 1955: 97-99 figs. 1-3.
- Jaeckel, S. Aculifera Aplacophora und Aculifera Placophora im Gebiet der Nord- und Ostsee. Kieler Meeresforsch. 10 1954: 261–271.
- Jaeckel, S. (1). Zur Systematik und Faunistik der Mollusken der nordlichen Balkanhalbinsel. Mitt. zool. Mus. Berl. 30 1954: 54-95 24 figs.
- Jaeckel, S. (2). Nordamerikanische Mützenschnecken der Gattung Ferrissia (Gastrop. Bas.) in deutschen Gewässern. Zool. Anz. 153 1954: 96–97.

Jaeckel, S. (3). Beiträge zur malakologischen Faunistik Thüringens. Mitt. Berl. Malak. Nr. 3 1953: 24–32.

[1955]

- Jaeckel, S. (4). Mollusken aus mitteldeutschen Gebirgen (Thüringer, Wald, Harz, Sächsische, Schweiz). Mitt. Berl. Malak. Nr. 4 1953: 3-7.
- Jaeckel, S. (5). Limapontia depressa Ald. et Hanc. var. pellucida Kevan, eine für das deutsche Küstengebiet neue Opisthobranchier-Art. Zool. Anz. 155 1955: 143-146 1 fig.
 - Jaeckel, S. see Pfitzner, I.
- Janus, H. Die Kartauserschnecke (Monacha cartusiana) neu in Württemberg. Jh. Ver. vaterl. Naturh. Württemb. 110 1955: 278-
- Jaume, M. L. Catalogo de la fauna Cubana. XI. Catalogo del genero Liguus en Cuba. Circ. Mus. biblio. zool. Habana 1954: 1423–1452.
- Jaume, M. L. (1). Novedades en Liguus de Cuba e Isla de Pinos (Mollusca Gasteropoda Bulimulidae). Circ. Mus. biblio. zool. Habana 1954: 1375-1388.
 - Jaume, M. L. see Aguayo, C. G.
- Jeannet, A. Die Macrocephaliten des Callovien von Herznach (Aargau). Ecl. geol. helv. 47 2 1954: 223–267 pls. 13–27 text-figs. 1–50 1 table.
- Jeffrey, S. W. The metabolism of oyster spermatozoa. 5. The effect of glycine. 6. The effect of metals. Aust. J. exp. Biol. med. Sci. 32 6 1954: 807-818 2 figs.
- Jeffreys, M. D. W. The cowry shell and the lozenge: African decorative art. Bull. S. Afr. Mus. Ass. 6 4 1955: \$3-93 11 figs.
- Jeletzky, J. A. Proposed use of teh plenary powers to preserve the generic name Aucella Keyserling, 1846 (Class Lamellibranchiata, Order Anisomyaria, Family Aviculidae) and the specific name of its type species "Avicula mosquensis" von Buch, 1884, and to suppress the generic name "Buchia" Rouillier, 1845. Bull. zool. Nom. 11 1955: 158–166.
- Jeletzky, J. A. (1). Evolution of Santonian and Campanian Belemnitella and paleontological systematics: exemplified by Belemnitella praecursor Stolley. J. Paleont. 29 3 1955: 478-509 pls. 56-58 1 text-fig.

- Jeletzky, J. A. (2). Belemnitella praecursor, probably from the Niobrara of Kansas, and some stratigraphic implications. J. Paleont. 29 5 1955: 876–885 1 text-fig.
- Jodot, P. Gasteropodes continentaux plaisanciens du Bassi lacustre de Baza (Province de Grenade). Mem. Comun. Inst. geol. Barcelona 10 1953: 43-50 pl. 1.
- Jodot, P. (1). Mollusques de petites dimensions des formations continentales Ludienne, Sannoisienne et Stampienne du Jura méridional et de la Haute-Savoie. Bull. Soc. géol. Fr. 6 4 7-9 1954: 537-555 pl. 23b.

Jodot, P. see David, L.

Jodrey, L. H. & Wilbur, K. M. Studies on shell formation. IV. The respiratory metabolism of the oyster mantle. Biol. Bull. Woods Hole 108 1955: 346-358 4 figs.

Jodrey, L. H. see Wilbur, K. M.

Johansson, J. On the Phylogeny of the Mollusca. Zool. Bidr. Uppsala 29 1952: 277-292.

Johnson, R. I. Supplement to Jesse Wedgwood Mighels (Occ. Pprs. No. 14). Occ. Pap. on Mollusks, Cambridge, Mass. 1 18 1954 p. 447 pl. 58.

Johnson, R. I. (1). Supplement to Joseph Pitty Couthouy. Occ. Pap. on Mollusks, Cambridge, Mass. 1 18 1954: 441-442.

Jones, A. H. The blue storm snail [Janthina]. Bull. Auckland Mus. Conch. Sect. No. 11 1955: 21.

Jones, A. H. (1). Varices on Cassids. Bull. Auckland Mus. Conch. Sect. No. 11 1955: 15.

Jones, A. H. see Thomson, W. P. Joshi, J. V. see Airan, J. W.

Jurado, M. D. Sunetta americana n. sp. (Lamellibranchia—Veneridia). Physis **20** 58 1951: 249–251 6 figs.

Jutting, W. S. S. v. B. The Malayan Streptaxidae of the genera *Discartemon* and *Oophana*. Bull. Raffles Mus. No. 25 [1954] 1955: 71-106 13 figs.

Jutting, W. S. S. v. B. (1). Vondsten van tropische kauri's in Nederland. Basteria 19 1 1955: 1-6. [Summary in English.]

- Jutting, W. S. S. v. B. (2). Süsswassermollusken von Sumba. Verh. naturf. Ges. Basel 66 1 1955: 49-60.
- Kaas, P. Report on a collection of Loricata from Manor Island, Karachi, with descriptions of three new species and a new variety. Zool. Meded. 33 1954: 1-9 14 figs.
- **Kado, Y.** The distribution of alkaline phosphatase in mantle tissue of bivalves. J. Sci. Hiroshima Univ. (Zool.) **15** 1954: 183–188 6 figs.

Kamada, Y. Non-marine mollusca from the Paleogene Uchigo group of the Joban coal-field, northeast Honshu, Japan. Trans. Proc. Palaeont. Soc. Japan, N.S. 17: 19-22 pl. 4.

Kanamaru, T. A History of the Japanese Conchology (34). Venus, Fukuyama 18 4 1955: 272–287. [In Japanese.]

Kanno, S. Non-marine fossil Mollusca from the Ôsawa-Pass, Fukushima Prefecture. Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: 77-86 1 pl. 1 fig.

Kanno, S. (1). Faunal analysis of the molluscan fauna from the raised beach deposits of Kamakura, Kanagawa Prefecture. Sci. Rep. Tokyo Kyoiku Daig. 4 1955: 23–47 10 figs.

Kanno, S. (2). Tertiary mollusca from the Taishu mine, Tsushima, Nagasaki Prefecture, Japan. Trans. palaeont. Soc. Japan N.S. 18 1955: 31-36 pl. 6 text-fig. 1.

Kanno, S. (3). On some Miocene species of *Dosinia* from Japan. Trans. palaeont. Soc. Japan N.S. 19 1955: 81-84 pl. 13.

Kantner, M. Uber subepitheliale Nervenausbreitungen bei Loligo vulgaris, Torpedo ocellata und Amiurus nebulosus. Z. mikr. anat. Forsch. 60 1954: 145-157 6 figs. [English summary.]

Kanungo, M. S. see Mukherji, R. N. Kanwisher, J. W. Freezing in intertidal animals. Biol. Bull. Woods Hole 109 1955: 56-63 3 figs.

Kapler, O. Viviparus viviparus morpha modestus morpha nova. Přírod. Sborn. Ostravsk. Kraje 16 1 1955: 89– 95 2 figs. [German summary.]

Kapp, K. Zwei Curiosa aus der Paläontologie. Boll. Soc. tiein. Sci. nat. 39-40 1945: xxi-xxiv 2 pls. Karandikar, K. R. see Subrahmanyam, T. V.

Kerslake, J. see Rutland, K.

Kaseno, Y. Neogene Argonautinae from Kahoku-gun, Ishikawa Prefecture, Japan. Sci. Rep. Kanazawa Univ. 3 2 1955: 339–344 1 pl. 2 figs.

Kato, K. & Kubomura, K. Calcium oxalate in the crystalline style of some mollusks. Sci. Rep. Saitama Univ. 2B 1 1955: 15-26.

Kato, K. & Kubomura, K. (1). Uric acid and guanine excretion from the crystalline style sac in some gastropods. Sci. Rep. Saitama Univ. 2B 1 1955: 21-34 2 pls. 3 figs.

Kawai, D. K. Carbonic anhydrase in pearl oyster. I. [Pinctada martensii.] Mem. Coll. Sci. Kyoto 21B 1 1954: 39-44.

Kawamoto, N. Y. & Motoki, E. Influence of diluted sea waters on the heart beat of pearl oyster. Bull. Jap. Soc. sci. Fish. 20 1954: 184–188 2 pls. [English summary.]

Kawamoto, N. Y. & Motoki, E. (1). Influence of diluted sea water on the pearl oyster, "Pteria (Pinctada) martensi Dunker." Rep. Fac. Fish. Univ. Mie 1 1954: 346–354 pl. 1, 2, figs. 1, 2.

Keen, A. M. Application for a ruling that works credited to S. A. Renier as of the dates 1804 and 1807 were not published within the meaning of article 25 of the "Règles." Bull. zool. Nom. 9 1954: 257-262.

Kegel, W. Zonar angeordnete Lebensgemeinschaften an Felsküsten bei Rio de Janeiro. Senckenbergiana Lethaea 35 3-4 1954: 231-234.

Keller, B. M. The Ordovician of Kazakstan. 1. Type sections of the Ordovician. Trav. Inst. géol. S.S.S.R. 154 1954: 5-47 text-figs. 1-5.

Kerney, M. P. Planorbis (Gyraulus) laevis Alder in Surrey. J. Conch. London 24 1955: 54.

Kerney, M. P. (1). On the former occurrence of *Vertigo parcedentata* (Al. Braun) in Hertfordshire. J. Conch. London 24 1955: 55-58.

Kershaw, R. C. A systematic list of the Mollusca of Tasmania, Australia. Pap. roy. Soc. Tasm. 89 1955: 289**Kershaw, R. C.** (1). The Victorian Mollusca. Vict. Nat., Melb. **72** 4 1955: 59-63.

Kershaw, R. C. (2). Studies on Australian Charopidae. Part 2. Some genera. Vict. Nat., Melb. 72 2 1955: 28-30.

Kershaw, R. C. (3). Studies on Australian Charopidae, Part 3—Planate genera. Vict. Nat., Melb. 72 9 1955: 137–142.

Kerslake, J. A. Aberrant cowries of the Keppel Islands, Queensland. Proc. R. zool. Soc. N.S.W. 1952-53, 1954: 40-41.

Kerslake, J. A. see Rutland, K.

Keyl, H.-G. Der Formwechsel der Chromosomen in der Spermatogenese von *Bithynia tentaculata* (L.). Chromosoma 7 1954: 387–419 30 figs.

Kira, T. Coloured illustrations of the shells of Japan. Osaka (Hoikusha) 1955 pp. viii + 204, 67 pls. col. text illust. [In Japanese.]

Kishimoto, H. An abnormal tentacle of Euhadra herklotsi hesperia Pilsbry. Venus, Fukuyama 18 4 1955: 282–283 text-fig.

Kiss-Kocsisné Bányai, M. Dunántúli eocén Cerithium-félék (Cerithien aus dem transdanubischen Eozän). Földt. Közl. 85 3 1955: 361–379 pls. 13–16. [Germ. summary p. 380.]

Kleerekoper, H. Limnological observations in Northeastern Rio Grande do Sul, Brazil I. Arch. Hydrobiol. 50 1955: 553-567 1 fig.

Kleinsmid, R. B. v. An appreciation [Capt. G. Allan Hancock, 80th birthday July 26, 1955.] Essays in the Nat. Sciences in honor of Capt. A. Hancock, Los Angeles 1955 pp. vii, viii, frontispiece.

Klinghardt, F. Neue Deutung einer scheinbaren Kriechspur eines Tintenfisches aus dem Solnhofener Plattenkalk. Paläont. Z. 29 1–2 1955: 79–82 pl. 7.

Knight, J. B. New families of Gastropoda. J. Wash. Acad. Sci. 46 2 1956: 41–42.

Knight-Jones, E. W. Notes on invertebrate larvae observed at Naples during May and June. Pubbl. Staz. zool. Napoli 25 1954: 135-144 2 figs. Knudsen, J. Notes on some marine prosobranchs from tropical West Africa. Rev. Zool. Bot. afr. 51 1955: 97-106 11 figs.

Knudsen, J. (1). Geographical variation of *Littorina obtusata* (L.) in the North Atlantic. Vidensk. Medd. Dansk naturh. Foren. 111 1949: 247–255 figs, tables & graphs.

Knudsen, J. see Adam, W.

Kobayashi, H. Cyto-chemical study on the nucleolus in the common Japanese oyster (Ostrea laperousi Schrenck). II. Nucleic acids and nucleoproteins. J. Sci. Hiroshima Univ. (Zool.) 15 1954 167–180 1 pl.

Kobayashi, T. (1). Izummonauta, a new genus and the Argonautinae, with a note on their rare but gregarious fossil occurrence. Jap. J. Geol. Geogr. 25 1-2 1954: 21-34 2 pls. 2 figs.

Kobayashi, T. (2). A contribution toward palaeo-flumenology, science of the oceanic current in the past, with a description of a new Miocene Aturia from Central Japan. Jap. J. Geol. Geogr. 25 1-2 1954: 35-56 1 pl. 2 figs.

Kobayashi, T. (3). The Ordovician fossils from the MacKay group in British Columbia, Western Canada, with a note on the Early Ordovician palaeogeography. J. Fac. Sci. Univ. Tokyo (2) 9 3 1955: 357–493 9 pls.

Kobayashi, T. & Mastani, K. On the occurrences of Aturia in provinces of Etchu and Iwami and their bearing on the palaeoflumenology in the Miocene of Japan. Trans. palaeont. Soc. Japan N.S. 17 1955: 1-4 pl. 1.

Kobayashi, T. & Mori, K. Studies on the Jurassic trigonians in Japan, Part I. Preliminary notes. Part II. Prosogyrotrigonia and the Trigoniinae. Jap. J. Geol. Geogr. 25 1954: 61-80, 155-175 2 pls. 3 figs.

Kobayashi, T. & Mori, K. (1). The Vaugoniinae from the Kitakami Mountains in North Japan. On the Jurassic Trigonians in Japan. Part III. Jap. J. Geol. Geogr. 26 1-2 1955: 73-88 2 pls. 2 figs.

Kobayashi, T. & Tamura, M. The Myophorellinae from North Japan. Studies on the Jurassic Trigonians in Japan. Part IV. Jap. J. Geol. Geogr. 26 1–2 1955: 89–103 2 pls.

Kocsisné, M. B. K. see Kiss-Kocsisné Banyai, M.

Koizumi, S. see Tomila, G.

Kolb, H. Über zwei abnorme Ammonitengehäuse der Gattung Paltopleuroceras. Geol. Bl. NO-Bayern 5 4 1955: 148–150 pl. 5 figs. 5 a, b, 6a, b.

Komai, T. & Emura, S. A study of population genetics on the polymorphic land snail *Bradybaena similaris*. Evolution **9** 1955: 400–418 2 figs.

Kotake, N. & Miyawaki, Y. Experimental studies on the propagation of the pearl oyster, *Pinctada martensii*. On the unusual high mortality of pearl oyster caused by bacteria. Bull. Jap. Soc. sci. Fish. 19 1954: 952-956.

Kováčik, J. see D. Andrusov.

Koyama, T. Studies on the Chromosome Numbers of two species of *Succinea* in Japan. Venus, Fukuyama **18** 4 1955: 259–265 pl. 10.

Krassner, M. Shellfish contamination controls in the New York City area. Circ. Mus. biblio. zool. Habana 1954: 1093-1096.

Kraus, O. Besprechung von: E. Frömming Biologie der mitteleuropäischen Landgastropoden. Arch. Molluskenk. 84 1955: 107–108.

Krausp, C. Die Pupilliden Estlands. Mitt. Berl. Malak. Nr. 3 1953: 10-23.

Krausp, C. (1). Die Clausiliiden Estlands [Estonia]. Mitt. Berl. Malak. Nr. 4 1953: 8-33.

Krausp, C. (2). Die Valloniiden und Eniden Estlands [Estonia] (Gastrop., Pulmon.). Mitt. Berl. Malak. Nr. 5 1954: 1–12.

Krenmayr, J. The Manila Clam, new-comer in our sea-food family (ex Seattle Times Oct. 16, 1955). Mins. Conch. Club, S. Calif. 152 1955: 4-5.

Kretschmann, I. Untersuchungen über Bau und Funktion der Atmungsorgane bei Ampullarien. Wiss. Z. Humboldt Univ. Math.-Nat. R. 4 2 1955: 109-119 16 figs.

Kretzoi, M. Jelentés a kislángi kalabriai (villa-frankai) fauñe feltárásáról. (Bericht über die calabrische (villafranchische) Fauna von Kisláng, Kom. Fejér. J. ber. ung. geol. Anst. 1953 1 1954: 213–238. [German summary p. 239.]

Kubista, V. Flavonfarbstoffe in einigen Stylommatophoren. Mém. Soc. zool. tehéosl. 18 1954: 177–180 2 figs.

Kubomura, K. see Kato, K.

Kubota, S. see Tomita, G.

Kugrin, L. N. Chlamys elegans Andrz. from the Tortonian deposits of Ugo-Zapadnoi on the outskirts of the Russian platform. Gheol. Sborn. 1 1954: 220-223 pl. 1.

Kühn, O. Cephalopoden der Trias in Unterkrain. Razpr. Akad. Ljubljani Cl. IV 2 1954: 387–394 1 fig.

Kühn, 0. (1). Süsswassermiozän von bosnischen Typus in Griechenland. Ann. géol. Hellen. 1 3 1951: 185–192 pl. 17 1 text-fig. [Summary in Greek.]

Kühn, O. (2). Der erste Nachweis von *Psiloceras* in den Pflanzen-schiefern des oberfränkischen Rätolias. Neues Jb. Geol. Paläont. B. 9 1955: 408–411.

Kummel, B. Triassic stratigraphy of south eastern Idaho and adjacent areas. Prof. Pap. U.S. Geol. Surv. 254-H 1954: 165-194 pls. 34-40 (geol. sects.).

Kummel, B. & Lloyd, R. M. Experiments on relative streamlining of coiled cephalopod shells. J. Paleont. 29 1 1955: 159-170 text-figs. 1-5.

Kuroda, T. A new Pleurotomaria from Japan with a note on a specimen of P. rumphii Schepman collected from Taiwan. Venus, Fukuyama 18 4 1955: 211-221, pls. 8, 9.

Kuroda, T. (1). On the Japanese species of Conus. Venus, Fukuyama 18 4 1955 : 284–301 1 fig.

Kuroda, T. & Habe, T. Volutidae in Japan. Illust. Catalogue Japanese Shells No. 5 1950 pp. 31–38 6 text-figs. pls. 5–7.

Kuroda, T. & Habe, T. (1). Nomenclatural notes. Illust. Catalogue Japanese Shells No. 4 1950 p. 30.

Kuroda, T. & Habe, T. (2). Nomenclatural notes. Illust. Catalogue of Japanese Shells No. 2 1950 p. 16.

Kurosawa, A. see Tomita, G.

Kwiatkowski, S. Conulaire pentagonale du carbonifère inférieur du Massif de la Ste. Croix. Ann. Soc. géol. Pologne 23 (1953) 1955: 157–164 pl. xi. [Russian and French summaries.]

Labruto, G. see Dulzetto, F.

Laidlaw, F. F. Notes on species of the genus Amphidromus from islands lying off the west coast of Sumatra, with descriptions of new races based on collections made by Dr. E. Jacobson in 1913. Zool. Meded. 33 1954: 75-81 2 figs.

Lal, M. B. & Baugh, S. C. Studies in histopathology—effects of the presence of a plagiorchid metacercaria on the tissues of the snail *Vivipara bengalensis* (Lamarck). Proc. Indian Acad. Sci. 42B 3 1955: 114-122 7 figs.

Lapparent, A. F. de. Sur quelques gisements cénomaniens de la falais du Dahar (extrême Sud Tunisien). Bull. Soc. Sci. nat. Tunis 6 1954: 191-1951 fig.

Laseron, C. F. Revision of the New South Wales Turridae. Sydney (Royal Zool. Soc. N.S.W.) 1954 56 pp. 12 pls.

Laseron, C. F. (1). Revision of the Liotiidae of New South Wales. Aust. zool. 12 1954: 1-25 49 figs.

Laseron, C. F. (2). Revision of the New South Wales Eulimoid shells. Aust. 2001. 12 1955: 83-107 77 figs.

Laseron, C. F. (3). The identity of *Phasianella virgo*. Proc. R. zool. Soc. N.S.W. 1953-54 [1955]: 77-78 2 figs.

Laseron, C. F. (4). The genus *Tolema* and its allies. Proc. R. 2001. Soc. N.S.W. 1953-54 [1955]: 70-74 12 figs.

Laseron, C. F. (5). Revision of the New South Wales Triphoridae, Rec. Aust. Mus. 23 4 1954: 139-158 30 figs.

Laseron, J. Sexual dimorphism in Strombus auris-disnae Linne. Proc. R. zool. Soc. N.S.W. 1952-53 1954: 41.

Laseron, J. (1). Notes on Haurakia trajectus Watson. Rissoa (Alvania) trajectus a good species.] Proc. R. zool. Soc. N.S.W. 1953-54 [1955]: 77.

Latif, A. & Qadir, G. A new snail pest *Macrochlamys* sp. (Fam. Zonitidae) at Lyallpur. Proc. 3rd Pakistan sci. Conf. pt. 3 1951: 44-45.

- Laviolette, P. Étude cytologique et expérimentale de la régénération germinale après castration chez *Arion rufus* L. Ann. Sci. nat. (Zool.) (11) 16 1955: 427-535 4 pls.
- Lawson, J. D. The geology of the May Hill inlier. Quart. J. geol. Soc. Lond. 111 1 1955: 85-113 figs. 1-5 1 map.
- Lay Yoe Soey. Giant "Kima" [Tridacna gigas] pearls from tropical waters. Circ. Mus. biblio. zool. Habana 1954: 1043.
- Lefranc, J.-P. Fossiles du Jurassique supérieur et des "Grès de Nubie" de la region de Sana (Yémen). Bull. Soc. géol. Fr. 6 4 7-9 1954 : 680-681 pl. 28 figs. 9, 10.
- Leloup, E. Chaetopleura unilineata n. sp., chiton de l'Equateur. Bull. Mus. Hist. nat. Belg. 30 40 1954 : 1-6 2 figs.
- Leloup, E. (1). A propos du Cryptochiton stelleri (Middendorf, 1846). Bull. Mus. Hist. nat. Belg. 31 14 1955: 1-3 1 fig.
- Leloup, E. (2). Chiton (Radsia) goodalli Broderip, 1832 = Chiton (Radsia) chierchiae Nierstrasz, 1908. Bull. Mus. Hist. nat. Belg. 31 23 1955; 1–10 2 figs.
- Leloup, E. (3). A propos de Chiton heterodon (Pilsbry, 1892) et de Chiton peregrinus Thiele, 1910. Bull. Mus. Hist. nat. Belg. 31 42 1955: 1-7 3 figs.
- Lemcke, H. Neurosecretion and incretory glands in a tectibranch mollusc. Experientia 11 8 1955: 320-322 3 figs.
- Leonhardt, H. Über nerven im Aplysia-Herzen. Pubbl. Staz. zool. Napoli 27 1955: 37–42 pl. 1. [English summary.]
 - Leuchtenberger, C. see Ito, S.
- Lewis, J. R. Observations on a high level population of limpets. J. Anim. Ecol. 23 1954: 85-100 1 pl. 6 figs.
 - Li Chien-min see Chang, Hsi.
- Lippincott, J. W. Hunting tree snails in Cuba. Circ. Mus. biblio. zool. Habana 1954: 1151-1155.
 - Lloyd, R. M. see Kummel, B.
- Loosanoff, V. L. The European oyster, in American waters. Science 121 1955: 119-121.

- Loosanoff, V. L., Engle, J. B. & Nomejko, C. A. Differences in intensity of setting of oysters and starfish. Biol. Bull. Woods Hole 109 1955: 75-81 2 figs.
- Loosanoff, V. L. & Nomejko, C. A. Growth of oysters with damaged shelledges. Biol. Bull. Woods Hole 108 1955: 151-159 1 fig.
- Lopes, H. de S. Sôbre duas espècies do gênero "Pomacea" Perry, com um estudo da genitália em ambos os sexos (Mesogastropoda, Architaenioglossa, Mollusca). Rev. bras. Eiol. 15 2 1955: 203-210 26 figs. [English summary.]
- Lopes, H. de S. (1). Sobre "Entodina lundi" (Mörch, 1871) (Mollusca, Pulmonata, Systrophiidae). Rev. bras. Biol. 15 3 1955: 243-246 7 figs.
- Ložek, V. Hippeutis riparius (Wester'und) im Donau-Tiefland. Arch. Molluskenk. 84 1955: 107.
- Ložek, V. (1). Die Molluskenfauna des Altvatergebirges. Přírod. Sborn. Ostravsk. Kraje 15 1 1954 : 16–65 2 pls. 12 maps. [German summary.] See also p. 169–
- Ložek, V. (2). Měkkýši Československého Kvartéru. (Mollusken des Tschechoslovakischen Quartärs). Rozpr. geel. Ust. čsl. 17 1955: 1–510 pls. 1–12 text-figs. 1–37. [Résumé in Russ. p. 449; in German p. 473.]
- I.ožek, V. & Mácha, S. Geographische Verbreitung der Schliessmundschnecke. Čas. nár. Mus. 123 2 1954: 148–155. [German summary.]
- Lucena, D. T. de. Tropicorbis nordestensis n. sp. du Nord-Est du Brésil. J. Conchyliol. Paris 95 1955: 20-22 text-fig. 1a-c.
- Ludbrook, N. H. The molluscan fauna of the Pliocene strata underlying the Adelaide Plains. Part 1. [New names of 78 species listed but to be described later, given here.] Trans. roy. Soc. S. Aust. 77 1954: 42-64 3 figs.
- Ludbrook, N. H. (1). The molluscan fauna of the Pliocene strata underlying the Adelaide Plains. Part II—Pelecypoda. Trans. roy. Soc. S. Aust. 78 1955: 18–87 pl. 1–6.
- Ludeman, D. Über die Bekämpfung der Wandermuschel *Dreissena poly*morpha in Industrieanlagen. Z. angew. Zool. 1954: 329–352.

- Lundgren, G. The land mollusca of Värmland and remarks on their ecology. Ark. zool. 6 1954: 443-484 20 maps.
- McCamley, F. Shell collecting—further afield. Proc. R. zool. Soc. N.S.W. 1952-53 1954: 41-42.
- McCauley, J. E. & Marriage, L. D. The intertidal mussel, piddock, and abalone resources of Oregon's outer coast. Fish Comm. Res. Briefs 6 1955: 4-13 4 figs.
- McCullough, F. S. & Duke, B. O. L. Schistosomiasis in the Gambia. 1.—Observations on the potential snail vectors of Schistosoma haematobium and S. mansoni. Ann. trop. Med. Parasit. 48 1954: 277–286 1 pl.
- MacGinitie, G. E. Distribution and ecology of the marine invertebrates of Point Barrow, Alaska. [Moll. pp. 172-180.] Smithson. Misc. Coll. 128 9 1955: 1-201 8 pls.
- McGinty, T. L. New marine mollusks from Florida. Proc. Acad. nat. Sci. Philad. 107 1955; 75-85 2 pls. 1 fig.
- McGinty, T. L. (1). Spirula spirula Linné, a remarkable find. Nautilus, Philad. 69 1955: 35.
- McGinty, T. L. (2). Todd Lopez Moise. Nautilus, Philad. 68 1955: 142-143.
- McGowan, J. A. Observations on the sexual behavior and spawning of the squid, *Loligo opalescens*, at La Jolla, California. Calif. Fish Game **40** 1954: 47-54 5 figs.
- McGowan, J. A. & Pratt, I. The reproductive system and early embryology of the nudibranch *Archidoris montereyensis* (Cooper). Bull. Mus. comp. zool. Harv. 111 1954: 259–276 2 pls. 1 fig.
- McHugh, J. L. Growth of oysters at Gloucester Point, Virginia. Virginia J. Sci. N.S. 6 4 1955: 246.
- Macha, S. Die Mollusken des Oder-Tales in der Mährischen Pforte. Přírod. Sborn. Ostravsk. Kraje 15 2–3 1954: 342–349. [German summary.]
- Macha, S. (1). Die Molluskenfauna der Wälder in der Umgebung von Michálkovice (Michalkowitz bei Ostrau). Sborn. přír. Společ. Mor. Ostravé 16 3 1955: 396–399 1 pl. 1 fig.
 - Mácha, S. see Ložek, V.

- McLauchlan, C. F. Three new land shells from New South Wales. Aust. Zool. 12 1954: 39 1 fig.
- McLauchlan, C F. (1). Land shells originally introduced. Aust. Zool. 12 1954: 40 4 figs.
- McMichael, D. F. The identity and validity of *Hyridella australis* (Lamarck) 1819. Nautilus, Philad. **69** 1955: 6-13.
- McMichael, D. F. (1). Review of "Australian Seashores": by Dakin & Bennett & Pope, 1952. Occ. Pap. Mollusks, Cambridge, Mass. 1 18 1954: 448–449.
- MacMillan, G. K. A preliminary survey of the land and freshwater Gastropoda of Cape Breton, Nova Scotia Canada. Proc. N.S. Inst. Sci. 23 4 1955: 389-408.
- McMillan, N. F. The range of *Planor-barius corneus* (L.) in the British Isles. J. Conch., London 24 1955: 63-65.
- Macnae, W. On four species of the genus Aplysia common in South Africa. Ann. Natal Mus. 13 1955: 223-241 7 figs.
 - MacPhail, J. S. see Medcof, J. C.
- Macpherson, J. H. Preliminary revision of the families Patellidae and Acmaeidae in Australia. Proc. roy. Soc. Vict. N.S. 67 2 1955: 229-256 12 pls. figs.
- Mahmoud, I. G. E.-d. Sur un nouveau genre albien d'Ammonite, Neposiella nov. gen. C.R. Soc. géol. Fr. 14 1953: 286-287.
- Mahmoud, I. G. E.-d (1). Les céphalopodes crétacés du Massif du Moghara (Sanaï-Egypte). Bull. Inst. Egypt. 36 1 1955 : 221-226.
- Malaroda, R. Contributo alle conoscenze paleontologiche del Pliocene dei dintorni di Strongoli, nel Crotonese (Catanzaro). R.C. Accad. Lincei (8) 19 1-2 1955: 50-58 1 pl.
- Malatesta, A. & Nicosia, M. L. Contributi alla conoscenza delle Faune Neogeniche e Quaternarie della Sicilia I. I fossili del Pliocene e Pleistocene di Agrigento della collezione Lomi. Boll. Uff. geol. Ital. 77 2-3 1955: 173-180.
- Mancuso, V. Azione del ciánuro di potassio sullo sviluppo di *Physa rivularis* Ph. R.C. Accad. Lincei (8) 19 1-2 1955: 71-73 6 figs.

Mancuso, V. (1). L'azione dell'azide sodico sullo sviluppo dell'uovo di *Physa* rivularis Ph. Riv. Biol. N.S. 47 2 1955 : 203–207 1 pl. 2 figs. [English summary.]

Mandahl-Barth, G. The freshwater mollusks of Uganda and adjacent territories. Ann. Mus. Congo belg. 8vo 32 1954: 1–206 96 figs.

Mandahl-Barth, G. (1). The anatomy and the systematic position of the Tanganyika snails Syrnolopsis and Anceya. Ann. Mus. Congo belge. N.S. 4to Zool. 1 1954: 217–221 6 figs.

Manolessos, N. A further aid for the knowledge of the geology of the island Kythira. Ann. géol. Hellen. 1 6 1955: 51-75 pls. 8-11.

Marche-Marchad, M. Mollusques testaces marins de l'Indochine—I. Nassidae. J. Conchyliol., Paris 95 1955: 28–37.

Marche-Marchad, M. (1). Les Mollusque testacés marins de l'Indochine. II. Cymatiidae et Bursidae. J. Conchyliol., Paris 95 1955: 100–107.

Marcus, E. Opisthobranchia from Brazil. Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: 89–261 30 pls.

Marcus, E. B.-R. & Marcus, E. The nudibranch Rhodope from South America. Comun. 2001. Mus. Montevideo 4 68 1952: 1–8 2 figs.

Marcus, E. & E. (1). Über Sand-Opistobranchia. Kieler Meeresforsch. 11 1955: 230–243 3 figs.

Marcus, E. & E. (2). Über Philinoglossacea und Acochlidiacea. Kieler Meeresforsch. 10 1954: 215–223 19 figs.

Margalef, R. Materiales para hidrobiologia de la isla de Mallorca. Publ. Inst. Biol. apl. Barcelona 15 1953: 5-111 1 pl. 7 figs. [English summary.]

Marriage, L. D. see McCauley, J. E.

Mars, P. Faunules malacologiques des lagunes de Marsala, Ganzirri et Faro (Sicile). Rapp. Comm. int. Mer Médit. 12 1954: 225–226.

Martin, W. E. Seasonal infections of the snail, Cerithidea californica Haldeman, with larval trematodes. Essays in the nat. Sciences in Honor of Capt. A. Hancock, Los Angeles 1955: 203-210 2 pls.

Martinis, B. Contributo alla paleontologia del Luteziano di Rocca Bernarda (Udine). Riv. ital. Paleont. 61 4 1955: 187-200 pl. 17.

Masatani, K. see Kobayashi, T.

Mason, J. & Colman, J. S. Note on a short-term marking experiment on the scallop *Pecten maximus* (L.) in the Isle of Man. Rep. Mar. biol. Sta. Pt. Erin 67 1954 (1955): 34-35.

Masuda, K. Miocene mollusca from Noto peninsula, Japan. Part 1. Trans. palaeont. Soc. Japan, N.S. 20 1955: 119-127 pl. 19.

Matsumoto, T. The bituberculate pachydiscids from Hokkaido and Saghalien (Studies on the Cretaceous ammonites from Hokkaido and Saghalien—8). Mem. Fac. Sci. Kyushu Univ. 5D 3 1955: 153–184 8 figs. 7 pls.

Matsumoto, T. (1). Family Kossmaticeratidae from Hokkaido and Saghalien. (Studies on the Cretaceous Ammonites from Hokkaido and Saghalien—VI.) Jap. J. Geol. Geogr. 26 1–2 1955: 115–164 3 pls. 13 figs.

Matsumoto, T. (2). Evolution of Peroniceratidae. Trans. palaeont. Soc. Jap. N.S. 18 1955: 37-44 text-figs. 1-3 1 table.

Matsumoto, T. & Obata, I. Some upper Cretaceous desmoceratids from Hokkaido and Saghalien. (Studies on the Cretaceous ammonites from Hokkaido and Saghalien—7). Mem. Fac. Sci. Kyushu Univ. 5D 3 1955:119–151 13 text-figs. 2 pls.

Matteson, M. R. Studies on the Natural History of the Unionidae. Amer. Midl. Nat. 53 1 1955: 126-145 7 figs.

Mattox, N. T. Observations on the Brachiopod communities near Santa Catalina Island. [Moll. list pp. 78-79, 81.] Essays in the nat. Sciences in Honor of Capt. A. Hancock, Los Angeles, 1955: 73-86 2 pls.

Mattox, N. T. (1). List of the Mollusca identified from samples of the benthos of San Pedro Basin, California. Allan Hancock Pacif. Exped. 19 1 1955: 154-158.

- Mattox, N. T. (2). Studies on the Opisthobranchiata. 1: A new species of the genus Tritoniopsis from Southern California. Bull. S. Calif. Acad. Sci. 54 1 1955: 8-13 2 pls.
- Maubeuge, P. L. Sur quelques documents ichnologiques du lias de la province de Luzembourg. Bull. Mus. Hist. nat. Belg. 30 5 1954: 1-6 3 pls.
- Maubeuge, P. L. (1). Sur l'âge de "l'Oolithe ferrugineuse" du Callovo-Oxfordien de Chaumont (Haut Marne) à Châtillon-sur-Seine (Côte-d'Or). Bull. Soc. belge Géol. Pal. Hydr. 64 1 1955: 38-44.
- Maubeuge, P. L. (2). Les Ammonites aaléniennes, bajociennes et bathoniennes du Jura suisse septentrional 11e partie. Schweiz. palaeont. Abh. 71 1955: 1-48 pls. 1-11.
- Mazzotti, L. Lymnaea obrussa Say, huespid intermediario de Fasciola hepatica. Rev. Inst. Salubr. Enferm. trop. Méx. 15 3 1955 : 163–165. [English summary.]
- Mead, A. R. The proposed introduction of predatory snails into California. Nautilus, Philad. 69 1955: 37-40.
- Medcof, J. C. Day and night characteristics of spatfall and of behaviour of oyster larvae. J. Fish. Res. Bd. Can. 12 1955: 270–286 2 figs.
- Medcof, J. C. & MacPhail, J. S. Survey of bar clam resources of the Maritime Provinces [of Canada]. Bull. biol. Bd. Can, 102 1955: 1-6 3 figs.
- Medeiros, L. C. M. de see Fraga de Azevedo.
- Meenakshi, V. R. A quantitative study of the distribution of calcium in the soft parts of Pila virens (Lamarck). J. zool. Soc. India 7 1955: 35-40.
- Meenakshi, V. R. (1). Protein catabolism in the developing eggs of Pila virens. Curr. Sci. 24 6 1955: 199-200.
- Meenakshi, V. R. (2). Role of carbonic anhydrase in the female genital duct of the apple snail [Pila virens]. Curr. Sci. 24 12 1955 : 415-416.
- Meenakshi, V. R. (3). Studies in the physiology of digestion in Pila virens (Lamarck). J. Anim. Morph. Physiol. **1** 1954 : 35–47 2 figs.

- Meenakshi, V. R. (4). Physiology of digestion in Melania crenulata. J. Anim. Morph. Physiol. 1 No. 2 1955: 33-41 7 figs.
- Melchinger, H. Unterschiede im Gehäusebau und Verhalten der beiden Heideschnecken Helicella ericetorum und Helicella obvia. Zool. Jb. (Syst.) 83 1955: 185-236 20 figs.
- Menzel, R. W. Some phases of the biology of Ostrea equestris Say and a comparison with Crassostrea virginica (Gmelin). Publ. Inst. Mar. Sci. Univ. Tex. 4 No. 1 1955: 69-153 7 pls.
- Menzel, R. W. & Hopkins, S. H. The growth of oysters parasitized by the fungus Dermocystidium marinum and by the trematode Bucephalus uculus. J. Parasit. 41 1955: 333-342 3 figs.
- Meyer, K. O. Naturgeschichte der Strandschnecke Ovatella myosotis (Drapaunaud). Arch. Molluskenk. 84 1955: 1-43 pls. 1-2.

Miani, N. see Munari, P.

- Mihailović, M. Description stratigraphique et paleontologique de quelques especes d'Ammonites des couches de Klauss de Greben (Serbie orientale). Ann. géol. Pén. balkan. 22 1954 : 123-134 pls. 1-3 text-fig. 1. French résumé.]
 - Miller, A. K. (1) see W. M. Furnish (1).
- Miller, A. K. & Furnish, W. M. Aturias from southern Chile. J. Paleont. 29 3 1955: 467-468 pl. 51 figs. 4-9.
- Miller, A. K. & Furnish, W. M. (1). The Carboniferous guide fossil Tylonautilus, in America. J. Paleont. 29 3 1955: 462-464 pl. 50.
- Miller, A. K. & Furnish, W. M. (2). The Carboniferous guide fossil Tylonautilus in America. J. Paleont. 29 3 1955: 462-464 pl. 50.
- Miller, A. K., Furnish, W. M. & Paguaga, A. S. Aturias from Costa Rica. J. Paleont. 29 3 1955: 465-466 pl. 51 figs. 1-3.
- Miller, A. K. & Garner, H. F. Lower Mississippian cephalopods of Michigan. Part III. Ammonoids and summary. Contr. Mus. Geol. Univ. Michigan 12 8 1955 : 113-173 pls. 1-7 text-figs. 1-16.
 - Miller, A. K. see Furnish, W. M.

Miller, A. K. see Glenister, B. F.

Miyake, M. see Horiguchi, Y.

Miyawaki, M. Notes on the shell structure of the oyster, *Gryphaea* (*Ostrea*) gigas from the Akkeshi Lake. J. Fac. Sci. Hokkaido Univ. (Zool.) 12 1954: 116-119 2 figs.

Miyawaki, Y. see Kotake, N.

Molnárné, D. I. Komárváros környékének levantei és pleisztocén rétegei. Les couches levantines et pléistocènes de environs de Komárváros. Jber. ung. geol. Aust. 1953 l 1954: 287–295 pl. l l map. [French and Russian summaries.]

Monastero, S. Secondo decennio di osservazioni entomologiche sui principali insetti nocivi della Sicilia (1940–1950). Atti Accad. Palermo (4) 12 1 1953: 5-44.

Mongin, D. Description d'un nouveau Lamellibranche: Nucula renouardi n. sp. du Portlandien du Liban. Bull. Soc. géol. Fr. 6 4 7-9 1954: 501-503 pl. 23a fig. 1, 2.

Moore, D. V. see Sandground, J. H.

Moore, H. B. X-rays reveal the inner beauty of shells. Nat. geogr. Mag. 107 3 1955: 427-434 figs.

Mori, K. see Kobayashi, T.

Morretes, F. L. de. Adenda e corrigenda ao ensaio de catalogo dos molusculos do Brasil. Arch. Mus. Paranaen. 10 1953: 37–76.

Morretes, F. L. de. Nova Thais do Brasil, Arch. Mus. Paranaen. 10 1953; 339–340 2 pls.

Morretes, F. L. de (1). Dois novas moluscos do Brasil. Arch. Mus. Paranaen, 10 1953: 331–336 l pl.

Morretes, F. L. de (2). Sobre Megalobulinus paranaguensis Pilsbry and Iering. Arch. Mus. Paranaen. 10 1953: 343-344.

Morris, D. The snail-eating behaviour of Thrushes and Blackbirds. Brit. Birds 47 1954: 33-49 8 pls. 4 figs.

Morrison, J. P. E. Conus eldredi, new name for one of the poison cones. Wash. Acad. Sci. 45 1 1955: 32.

Morrison, J. P. E. (1). Ecological notes on the mollusks and other animals of Raroia. Atoll res. Bull. No. 34 1954: 1–18.

Morrison, J. P. E. (2). Notes on American cyclophoroid land snails, with two new names, eight new species, three new genera, and the family Amphicyclotidae, separated on animal characters. J. Wash. Acad. Sci. 45 5 1955: 149-162 31 figs.

Morrison, J. P. E. (3). A correction [Cyclochittya schermoi Morrison to schumoi]. Nautilus, Philad. 69 1955: 72.

Morton, J. E. The evolution of vermetid gastropods. Pacif. Sci. 9 1 1955: 3-15 4 figs.

Morton, J. E. (1). The structure and function of the stomach and sorting caecum in *Lunella smaragda* (Martyn) (Turbinidae). Proc. Malac. Soc. London 31 1955: 123–137 5 text-figs.

Mosebach, R. see Einsele, G.

Motoki, E. see Kawamoto, N.

Mozley, A. Sites of infection. Unstable areas as sources of parasitic diseases: Schistomiasis and Fascioliasis, London (H. K. Lewis & Co.) 1955 pp. x+96 14 figs.

Mukherji, R. N. & Kanungo, M. J. Invertase in the digestive gland of the common Indian freshwater Pelecypod. Lamellideus sp. Sci. & Cult. 20 5 1954: 253-254.

Müller, A. H. Beiträge zur Stratonomie und Ökologie des germanischen Muschelkalkes. Geologie 4 3 1955: 285–297 pls. 1–3.

Mullerried, F. K. G. Nuevo radiolitido supracretacico en terrenos de Coahuila, Noreste de Mexico. Ciencia, Mex. 12 1-2 1952: 36-38 4 figs.

Munari, P. F. & Miani, N. La struttura sub-microscopica del neuro-axoplasma normale e degenerato osservato al microscopio elettronico. Monit. zool. ital. 63 1955: 170–187 14 figs. [English summary.]

Mundhenke, G. Untersuchungen zur Frage des Farbsehens einiger euthyneurer Schnecken. Zool. Jb. (Allg. Zool.) 66 1955: 33-61 17 figs.

Munro, I. S. R. Eggs and larvae of the sabre-toothed oyster blenny, Dasson steadi (Whitley) (Blenniidae). Aust. J. Mar. Freshw. Res. 6 1 1955: 30–34 2 figs. Muraour, P. Sur quelques sediments dragues au large du littoral compris entre le Cap Djinet et l'embouchure de l'Oued Sebaou. [Short list p. 101.] Bull. Sta. Aquic. Pêche Castiglione (N.S.) 7 1955: 89-108 5 pls. 4 figs.

Murti, N. N. see Subrahmanyan, T. V.

Mutihac, V. see Filipescu, M. G.

Mutvei, H. see V. Jaanusson.

Nagabhushanam, R. Tolerance of the marine wood borer, Martesia striata (Linn.) to waters of low salinity. J. zool. Soc. India 7 1955: 83-86.

Nagao, T. & Huzioka, K. Fossil Acila from Hokkaido and Karahuto (Saghalin). J. Fac. Sci. Hokkaido Univ. (4) 6 1941: 113–141 3 pls.

Nagao, T. see Matumoto, T.

Nair, N. B. The digestive enzymes of *Bankia indica* Nair. Curr. Sci. 24 4 1955: 126-127.

Nair, N. B. (1). Cellulase activity of the crystalline style of the wood-boring pelecypod *Bankia indica* Nair. Curr. Sci. 24 6 1955: 201.

Nakazawa, K. A study of the pelecypod-fauna of the Upper Triassic Nabae group in the northern part of Kyoto Prefecture, Japan, Part 2. Bakevelliids. Mem. Coll. Sci. Kyoto 21B 2 1954: 213-222 4 pls.

Nakazawa, K. & Shimizu, D. Discovery of *Glyptophiceras* from Hyogo Prefecture, Japan. Trans. palaeont. Soc. Japan N.S. 17 1955: 13-18 pl. 3 1 text-fig.

Nash, C. B. see Dingle, A. N.

Nath, V. & Chopra, H. C. On the origin of the golgi dictyosomes from the mitochondria and acrosome formation in the spermatogenesis of the slug Anadenus altivagus (Theobald). Res. Bull. Panjab Univ. No. 74 1955: 91-100 2 pls. figs.

Nelson, L. Adenosinetriphosphatase of Mytilus Spermatozoa. 1. Effects of pH, calcium and magnesium and concentration of enzyme and substrate. Biol. Bull. Woods Hole 109 1955: 295-305 6 figs.

Newell, G. E. The marine fauna of Whitstable. Ann. Mag. nat. Hist. (12) 7 1954: 322-350.

Newell, G. E. see Smith, J. E.

Newton, W. L. The establishment of a strain of Australorbis glabratus which combines albinism and high susceptibility to infection with Schistosoma mansoni. J. Parasit. 41 1955: 526-528 1 fig.

Newton, W. L. & Brand, T. von. Comparative physiological studies on two geographical strains of *Australorbis glabratus*. Exper. Parasit. 4 1955: 244–255.

Nicol, D. An analysis of the arctic marine pelecypod fauna. Nautilus, Philad. 68 4 Ap. 1955: 115-122.

Nicol, D. (1). Morphology of Astartella, a primitive heterodont pelecypod. J. Paleont. 29 1 1954: 155-158 text-figs. 1-4.

Nicol, D. (2). Nomenclatural review of the Paleozoic pelecypod family Conocardiidae. J. Paleont. 29 3 1955: 552-553.

Nicosia, M. L. La fauna miocenica di "La Portella" presso Roccapalumba (Sicilia). Contributi alla conoscenza delle Faune Neogeniche a Quaternarie della Sicilia. Boll. Uff. geol. Ital. 77 2-3 1955: 363-376 pls. 1-2.

Nicosia, M. L. see Malatesta, A.

Niel, J. P. van. Anentome helena (v. d. B.) in een aquarium. Penggemar Alam 34 3-4 1954: 107.

Nomejko, C. A. see Loosanoff, V. L.

North, W. J. Size distribution, erosive activities, and gross metabolic efficiency of the marine intertidal snails, *Littorina planaxis* and *L. scutulata*. Biol. Bull. Woods Hole **106** 1954: 185–197 7 figs.

Nümann, W. & Beth, K. Die Ansiedlungszeiten der wichtigsten Bewüchsorganismen in der nördlichen Adria. Hidrobiologi, Istanbul 3B 1 1955: 3–33 4 figs.

Nüno, H. On a manganese nodul and *Perotrochus* dredged from the banks near the Izu Island, Japan. Rec. oceanogr. Wks. Japan 2 2 1955: 120–124 2 pls. I fig.

Obata, I. see Matsumoto, T.

Oberling, J. J. Shell structure of West American Pelecypoda. J. Wash. Acad. Sci. 45 4 1955: 128–130 2 figs. Obr, S. Contribution to the study of the fauna of springs, lakes and torrents in the mountains Liptovské hole (Tatra-Czechoslovakia). Mem. Soc. zool. tcheosl. 19 1955: 10-26.

Odhner, N. H. Some notes on *Pseudo-chama*. Nautilus, Philad. **69** 1955: 1-6 pl. 1.

Ojima, Y. & Ikenaga, K. Studies on the pearl-sac in molluses. II. Distribution of Alkaline phosphatase in the mantle and pearl-sac of the fresh water pearl mussel (Hyriopsis schlegelii). Zool. Mag., Tokyo 63 1954: 26-29 6 figs. [English summary.]

Okada, H. see Imai, T.

Okutani, T. On a new species of Carinaria C. japonica. Venus, Fukuyama 18 4 1955: 251-258 3 text-figs.

Oldfield, E. (née Whitehead). Observations on the anatomy and mode of life of Lasaea rubra (Montagu) and Turtonia minuta (Fabricius). Proc. Malac. Soc. London 31 1955: 226–249 11 text-figs.

Oliveri, A. see Baroni, C.

Olsen, A. M. Some facts about scallops from diving investigations. Pap. roy. Soc. Tasm. 88 1954: xiii-xiv.

Olsen, A. M. Underwater studies on the Tasmanian commercial scallop, Notovola meridionalis (Tate) (Lamellibranchiata; Pectinidae). Aust. J. mar. freshw. Res. 6 1955; 392-409 5 figs.

Olsson, A. A. see Pilsbry, H. A.

ômori, M. Conchocele compacta Ishizaki and its subspecies minor from the Tertiary of Japan. Sci. Rep. Tokyo Kyoiku Daig. 3C 17-21 1954: 87-97 1 pl. 1 fig.

ômori, M. (1). On some fossil new species of the genus *Propeamussium* from Japan. Sci. Rep. Tokyo Kyoiku Daig. 4 1955: 7-22 2 pls.

Orbigny, A. d'. Los moluscos de Cuba en la historia física, politica y natural de la Isla de Cuba, Parts I-VI [Cephalopoda, Pteropoda, Gastropoda]. Circ. Mus. biblio. zool. Habana 1953: 717-728: 741-762: 781-797: 821-854: 881-904: 927-948.

Orr, P. R. Heat death, I. Time-temperature relationships in marine animals. Physiol. Zoöl. 28 1955: 290-294 1 fig.

Ostergaard, J. M. Some Opisthobranchiate Mollusca from Hawaii. Pacif. Sci. 9 2 1955: 110-136 2 pls. 17 figs.

Oyama, K., Yamamoto, T. & Tokioka, T. Invertebrate fauna of the intertidal zone of the Tokara Islands. VIII. Molluscan shells. Publ. Seto Mar. biol. Lab. 4 1 1954: 1-15.

Ozaki, H. & Shikama, T. On three skytic molluses from Gunma Prefecture, Central Japan. Bull. nat. Sci. Mus. Tokyo N.S. 1 2 1954: 42-45 1 pl. 5 figs.

Ozaki, H. Stratigraphy of the basal comglomerate of the Pliocene Nagarai formation in the Tyôsi City, Kantô region. Bull. nat. Sci. Mus. Tokyo N.S. 121954:46-615 figs.

Ozaki, K. see Hayasaka, I.

Paguaga, A. S. see Miller, A. K. & other'.

Pain, T. New freshwater gastropod mollusks on the African genus *Lanistes*. Breviora 31 1954: 1–4 7 figs.

Pain, T. (1). Notes on some New Caledonian *Placostylus*. J. Conchyliol., Paris 95 1955: 11-19 9 text-figs.

Palmer, K. v. W. Request for a ruling on the question of the type species of *Ancilla* Lamarck, 1799. Bull. zool. Nom. 9 1954: 219.

Palmer, K. v. W. (1). Comment on the type species of *Ancilla* Lamarck, 1799 (Class Gastropoda). Bull. zool. Nom. 9 1954: 269.

Palmer, K. v. W. (2). New name for *Hipponix floridanus* Palmer, 1953. J. Paleont. 29 3 1955: 558.

Pantić, N. see Stevanović, P. M.

Papp, A. Brack- und Süsswasserfaunen Griechenlands. 2. Brack- und Süsswasserarten von Kythera. Ann. géol. Hellen. 1 1 1947: 112–119 pl. 9 text-fig. 1. Summary in Greek.

Papp, A. (1). Brack- und Süsswasserfaunen Griechenlands. 3. Über Lebensspuren an Süsswassergastropoden. Ann. géol. Hellen. 1 1 1947: 126–128 pl. 10. [Greek summary.]

Papp, A. (2). Brack- und Süsswasserfaunen Griechenlands. IV. Süsswassermollusken aus dem Pliozän von Elis (Peloponnes). Ann. géol. Hellen. 1 5 1953: 107–113 pl. 23.

- Papp, A. (3). Brack- u. Süsswasserfaunen Griechenlands. V. Bemerkungen über Melanopsiden der Untergattung Melanosteira Oppenheim 1891. Ann. géol. Hellen. 1 6 1955: 122–132 pl. 20.
- Papp, A. (4). Nubecularien aus den sarmatischen Schichten von Wiesen (Bgld.). Burgenländ. Heimat Bl. 17 2 1955: 49-55 1 pl.
- Papp, A. (5). Bemerkungen über Vorkommen und Variabilität der Bivalvengattung *Oncophora*. Verh. geol. Bundesanst. 2 1955: 120–133 pls. 1–2.
- Papp, A. & Psarianos, P. Uber einige Süsswassergastropoden aus pliozänen Ablagerungen Griechenlands. Ann. géol. hellen. 1 6 1955: 145–150 pl. 23.
- Paraense, W. L. Self and cross-fertilization in *Australorbis glabratus*, Mem. Inst. Ows. Cruz 53 1955: 277-291. [English translation p. 284-291.]
- Paraense, W. L. & Deslandes, N. Reproductive isolation between Australorbis glabratus and A. nigricans. Mem. Inst. Ows. Cruz 53 1955: 321-327. [English transl.]
- Paraense, W. L. & Deslandes, N. (1). Observations on the morphology of *Australorbis glabratus*. Mem. Inst. Osw. Cruz 53 1955: 87-103 4 pls. 9 figs.
- Paraense, W. L. & Deslandes, N. (2). Observations on the morphology of Australorbis nigricans. Mem. Inst. Osw. Cruz 53 1955: 121-134 3 pls. 11 figs.
- Paraense, W. L. & Deslandes, N. (3). Studies on Australorbis centimetralis I. Morphology in comparison with A. glabratus. Rev. bras. Biol. 15 3 1955: 293-307 28 figs.
- Parker, R. H. Changes in the invertebrate fauna, apparently attributable to salinity changes, in the bays of central Texas. J. Paleont. 29 2 1955: 193-211 text-figs. 1-8 2 tables.
- Parmalee, P. W. Some ecological aspects of the Naiad fauna of Lake Springfield, Illinois. Nautilus, Philad. 69 1955: 28-34 tables & map.
- Parodiz, J. J. Un nuevo gastrópodo terres tre del Eoceno de Patagonia. Phys is 20 57 1949: 174-179 1 fig.
- Parodiz, J. J. (1). Austraborus n. nom. pro Microborus Pilsbry, 1926. Physis 20 57 1949: 189–190.

Parodiz, J. J. (2). Notes sobre Strophocheilus fósiles de Argentina. Physis 20 57 1949: 180–184 1 fig.

[1955]

- Parodiz, J. J. (3). Métodos de conquiliometria. Physis 20 58 1951 : 243—248 3 figs.
- Parodiz, J. J. (4). Una nueva especie de *Plectostylus* (Gastr. Pulm.) de la Argentina. Physis 20 58 1951 : 334–335 1 fig.
- Pašić, M. Sur la position stratigraphique de l'espèce Glauconia coquandi d'Orb, et de ses variétés (Repelin) dans les couches du Crétacé supérieur chez nous. Zborn. Radova geol. Inst. Beograd 7 1954: 54–67 (French résumé p. 66).
- Pasternak, S. I. Ancycloceras bipunctatum Schlüter from the Maastrichtian Flags of Volychno-Podolskoe. Gheol, Sborn. 1 1954: 157-159 text-fig. A.
- Paul, W. J. Varices on Cassids. Bull. Auckland Mus. Conch. Cl. No. 8 1952: 10.
- P[aulian], R. Un nouveau mollusque Malgache. Nat. malgache 7 1955: 226.
- Pelluet, D. & V. E. The effects of visible radiation on the ovotestis of the slug *Deroceras reticulatum* (Mull.). Proc. N.S. Inst. Sci. 23 4 1955: 420-421.
- Penndorf, H. Die Ceratiten-Schichten am Meissner in Niederhessen. Abh. senckenb. Naturf. Ges. No. 484 1951: 1-24 pl. 1-6.
- Perry, L. M. & Schwengel, J. S. Marine shells of the western coast of Florida. With revisions and additions to Louise M. Perry's Marine Shells of the southwest coast of Florida. Ithaca, N.Y. (Palaeontological Res. Inst.) 1955 pp. 318 frontis. 55 pls. 5 figs.
- **Petrbok, J.** České Pseudoanodonty. Čas. nár. Mus. **123** 2 1954 : 226-227 3 pls.
- Petter-Receveur, G. Les Lamellibranchs de Padern (Aude) [S. France]. Ann. Paléont. 41 1955: 55-74 pl. 1-3 text-fig. 1.
- Peyer, B. Sull'accoppiamento nel genus *Limax*. Boll. Soc. ticin. Sci. nat. 49 1954: 82-87 1 pl.
- Pfeiffer, K. L. Die Albinarien des Dodekans (Moll. Clausiliidae). Arch. Molluskenk. 84 4/6 1955: 109–153 pls. 8–11 3 maps & text-figs.

Pfitzner, I. Vorkommen von Trichia sericea Drap. in der Mark Brandenburg. Mitt. Berl. Malak. Nr. 6 1954: 51-55

Pfitzner, I. & Jaeckel, S. Beitrag zur Molluskenfauna des Harzes. Mitt. Berl. Malak. Nr. 6 1954: 28–50.

Pieltain, C. B. y. El museo Mexicano de Malacologia (Coleccion Carlos). Ciencia, Méx. 14 7–8 1954: 172– .

Pilsbry, H. A. Another Floridan *Conus.* Nautilus, Philad. **69** 2 Oct. 1955: 47–48 pl. 3 figs. 10, 11.

Pilsbry, H. A. & Brunson, R. B. The Idaho-Montana slug Magnipelta (Arionidae). Notul. nat. Acad. Philad. No. 262 1954: 1-6 11 figs.

Pilsbry, H. A. & Olsson, A. A. Miocene land shell fossils from the Dominican Republic. Notul. nat. Acad. Philad. No. 266 1954: 1-44 figs.

Pitchford, G. W. Lauria anglica (Wood) at Meathop Fell, Westmorland. J. Conch. London 24 1955: 62.

Pitchford, G. W. (1). Helix nemoralis (L.) in a calcium deficient habitat. J. Conch. London 24 1955: 54.

Pitchford, G. W. (2). Dreissena polymorpha (Pallas) in a closed pond. J. Conch. London 24 2 1955: 46.

Pitchford, G. W. (3). Snails preyed upon by jays and magpies. J. Conch. London 24 1955: 40.

Plate, H.-P. Zur Molluskenfauna der Greifswalder Oie und des Ruden. Arch. Fr. Naturg. Mecklenburg 1 1954: 177–192.

Plate, H.-P. see Frömming, E.

Plate, H.-P. & Frömming, E. Über das Verhalten von Landlungenschnecken gegenüber verschiedenen Sorten der Zierpflanze Chrysanthemum indicum. Z. angew. Zool. 1954: 187–195.

Plöchinger, B. Eine neue Subspezies des Barroisiceras haberfellneri v. Hauer aus dem Oberconiac der Gosau Salzburg. S.B. öst. Akad. Wiss. Abt. I. 164 4-5 1955: 203-206 l pl. 2 figs.

Poel, L. Van de. Structure du test et classification des nucules. Bull. Mus. Hist. nat. Belg. 31 3 1955: 1-11.

Pohl, L. L. Understanding the cultured pearl. How Japanese put a shell-fish to work. Nat. Hist. N.Y. **64** 1955: 319–321 4 figs.

Poisson, H. Contribution a l'étude de la faune malacologique marine de l'Archipel des Chagos et de l'ile d'Agalega. Bull. Acad. malgache N.S. 31 1954: 63-70 2 pls.

Pokorný, V. The age of the clay-balls and origin of fauna in the Tortonian sands at Žabčice (Moravia, extra-Alpine Neogene Basin). Mem. Soc. Sci. Bohème 1951 [1953] 8:1-7. [English summary.]

Pokorný, V. see Augusta, J.

Pollini, A. La serie stratigraphica del retico di Monte Castello (Zogno-Val Brembana). Atti Soc. ital. Sci. nat. 94 3-4 1955; 329-368 1 pl. 1 fig.

Pomeroy, L. R. & Haskin, H. H. The uptake and utilization of phosphate ions from sea water by the American oyster, *Crassostrea virginica* (Gmel.). Biol. Bull. Woods Hole 107 1954: 123–129 2 figs.

Pope, E. C. Bather's itch, or schistosome dermatitis. Aust. Mus. Mag. 11 9 1955: 288-291 figs.

Popovici, Z. & Angelescu, V. La economia del Mar y sus relaciones con la alimentación de la humanidad. Publ. Ext. cult. Mus. argent. Cienc. nat. No. 8 1954: 2 vols. [see Vol. 1 pp. 486-501; Vol. 2 pp. 729-731].

Porta, J. de see Solé, N.

Portmann, A. La métamorphose "abritée" de Fusus (Gast. Prosobranches). Rev. suisse Zool. 62 Suppl. 1955: 236–252 11 figs.

Pouderoyen see Altena, C. O. v. R. and others.

Powell, A. W. B. Additions to the recent Mollusca fauna of New Zealand. Bull. Auckland Mus. Conch. Sect. No. 9 1953: 14–20.

Powell, A. W. B. (1). "Old windows for new ones." Bull. Auckland Mus. Conch. Sect. No. 10 1954: 13-15.

Powell, A. W. B. (2). The Molluscan land operculate genus *Liarea*. Rec. Auckland [N.Z.] Inst. 45954:271-2935 pls. 4 figs.

Powell, A. W. B. (3). Mollusca of the southern islands of New Zealand. Cape Exped. Ser. Bull. No. 15 1955: 1-151 5 pls. figs.

Powell, L. H. A study of the Ozarkian faunas of Southeastern Minnesota. Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935: 1–80 17 pls.

Prantl, F. Biostratigrafický význam českych Goniatitú, Cas. nár. Mus. 123 1 1954: 15–19.

Prantl, F. & Růžička, B. Straba nov. gen., a new Devonian pelecypod from Bohemia. Acta Mus. nat. Prag. Series B 10 3 1954: 1-29 pl. 1-4 text-figs. 1-10.

Prantl, F. & Růžička, B. (1). Šufia nov. gen., nový mlž z českého devonu (Šufia nov. gen., a new Devonian pelecypod from Bohemia). Sborn. geol. Ust. Čsl. 21 ser. palaeont. 1954: 303— 316 (1–14) 1 pl. 1 text-fig. [English & Russian résumé.]

Prantl, F. & Růžička, B. (2). *Mnataia* nov. gen., a new Devonian pelecypod from Bohemia. Acta Mus. nat. Prag. Series B 11 5 1955: 1–18 f. 1–9 pl. 1. [English summary.]

Pratt, I. see McGowan, J. A.

Prenant, M. see Gabe, M.

Přibyl, A. & Růžička, B. Neklania nov. nom. for Dalila Barrande, 1881 (Lamellibranchiata). Přírod. Sborn. Ostravsk. Kraje 14 1–2 1953: 57–59. [English summary.]

Price, W. A. Oyster reefs of the Gulf of Mexico. Fish. Bull. U.S. 55 1954: 491.

Priolo, O. Nuova revisione dell conchologie marini di Sicilia. Memoria II-VII. Atti Accad. Gioenia (6) 7 1951: 95-119, 261-285 (6) 8 1953: 81-109 1 pl., 164-190 (6) 9 1954: 53-91, 186-222.

Prud'homme van Reine, W. J. Plants and animals of the seashore. A handy guide containing drawings and descriptions of over 450 species of birds, shell-fish, seaweeds, fish, etc. London (John Murray) 1955: pp. vi + 138, frontis. col., 34 pls.

Pruvot-Fol, A. Le bulbe buccal et la symétrie des mollusques. II. Arch. Zool. exp. gén. 91 1954 : 235-330 33 figs.

Pruvot-Fol, A. (1). Mollusques Opisthobranches. Faune de France 58 1954: 1–460 1 pl. 173 figs.

Psarianos, P. S. Die Pliocänbildungen von Achaia. Ann. géol. hellen. 1 3 1951: 193-214, German summary p. 205. Psarianos, P. (1). Hydrogeologische Untersuchungen im Becken von Arta. Ann. géol. hellen. 1 5 1953: 65-74 text-figs. 1-3 5 tables. [German summary.]

Psarianos, P. (2). Vivipara mitzopoulosi, eine neue Art aus den neogenen Bildungen von Achaia (Nordpeloponnes). Ann. géol. hellen. 1 5 1953: 93-96 pl. 22.

Psarianos, P. (3). Beiträge zur Kenntnis des Neogens von Lakonien (Peloponnes). Ann. géol. hellen. 1 6 1955: 151–184 pl. 24. [German summary.]

Psarianos, P. see Papp, A.

Puffer, E. L. & Emerson, W. K. Catalogue and notes on the gastropod genus *Busycon*. Proc. biol. Soc. Wash. 67 1954: 115–147.

Punt, A. The contracture of the anterior byssal muscle of *Mytilus*. Pubbl. Staz. zool. Napoli 27 1955: 313-317.

Purchon, R. D. A note on the biology of *Tridacna crocea*. Proc. Malac. Soc. London 31 1955: 95-110 6 text-figs.

Purchon, R. D. (1). A note on the biology of the Lamellibranch Rocellaria (Gastrochaena) cuneiformis Spengler. Proc. 2001. Soc. Lond. 124 1954: 17-33 11 figs.

Puri, H. S. Hermanites, new name for Hermania Puri, 1954. J. Paleont. 29 3 1955: 558.

Quadir, G. see Latif, A.

Quattrini, D. Osservazioni sulla struttura del nucleo in alcuni tessuti ed organi di *Papillifera bidens* L. Monit. zool. ital. 63 1955: 239–257 2 pls. 4 figs. [English summary.]

Quick, H. E. Journey to the Gughi Highlands (Southern Ethiopia), 1948– 1949: Gastropoda, slugs from the high mountains. J. linn. Soc. (zool.) 42 1954: 382-386 17 figs.

Radoman, P. Deux représentants nouveaux du gastéropode *Horatia* Bourguignat dans le lac d'Ohrid. Rec. Trav. Sta. Hydrobiol. Ohrid No. 1-6 1953: 63-71 9 figs.

Radoman, P. see Stanković, S.

Raesinck, M. J. see Cate, J. T.

Răileanu, G. Explorations géologiques dans la région Svinița-Fața Mare. Bul. sti. Acad. Republ. rom. 5 2 1953: 307–409 2 pls. 38 figs. [French résumé.]

Răileanu, G. (1). La présence du genre Anthracosia dans le Permien inférieur de la zone Svinitza-Svinecea Mare. Rev. Biol., București 2 1955: 97-100 3 figs.

Ramamoorthi, K. Amino acids in the reserve nutrition materials of the egg of *Pila virens* (Lamarck). Curr. Sci. **24** 10 1955: 339-340 1 fig.

Ramamoorthi, K. (1). Studies in the embryology and development of some Melaniid snails. J. zool. Soc. India 7 1955: 25-34 8 figs.

Rancurel, P. Teredo tritubulata Moll 1941: Description des valves. Bull. Inst. franç. Afr. noire 17 4 1955: 1145-1148 2 figs.

Rancurel, P. (1). Teredo thomsoni Tryon et Teredo lieberkindi Roch Transformations morphologiques des palettes au cours de la croissance. Bull. Inst. franç. Afr. noire 17 4 1955: 1149-1156 5 figs.

Rangnekar, P. V. The nutritive capacity of the blood of some representative crustaceans and molluscs. J. Anim. Morph. Physiol. 2 1955: 101–104.

Rangnekar, P. V. (1). A note on the colourless blood on the fresh water mussels Lamellidens lamellidens. J. Univ. Bombay N.S. 21B 3 1955: 88-89.

Ranzoli, F. Osservazioni citometriche su ovociti in accrescimento. Boll. Zool. 21 1954: 129-132.

Rao, K. P. Tidal rhythmicity of rate of water propulsion in *Mytilus* and its modifiability by transplantation. Biol. Bull. Woods Hole 106 1954: 353-359 5 figs.

Rapson, A. M. Feeding and control of toheroa (Amphidesma ventricorsum Gray). Eulamellibranchiata populations in New Zealand. Aust. J. mar. & freshw. Res. 5 3 1954: 486-512 5 pls. 1 fig.

Raven, C. P. & Beenakkers, A. M. T. On the nature of head malformations obtained by centrifuging the eggs of Limnaea stagnalis. J. Embryol. exp. Morph. 3 1955: 286-303 2 pls. 2 figs.

Raven, C. P., Roon, A. C. de & Stadhouders, A. Morphogenetic effects of a heat shock on the eggs of *Limnaea stagnalis*. J. Embryol. exp. Morph. 3 1955: 142-159 2 pls. 1 fig.

Ray, S. M. & Chandler, A. C. Dermocystidium marinum a parasite of oysters. Exper. Parasit. 4 2 1955: 172—figs.

Reed, J. J. see Wellman, H. H.

Rees, W. J. Notes on the distribution of Cephalopods in the Eastern Mediterranean. J. Conchyliol, Paris 95 1955: 83–85.

Rees, W. J. (1). The larvae and late larval stages of Octopus macropus Risso, Proc. Malac. Soc. London 31 1955: 185–189 pls. 9, 10.

Reesinck, M. J. see Cate, J. T.

Rehder, H. A. Gulf of Mexico, its origin, waters and marine life. Mollusks. Fish. Bull. U.S. 55 1954: 469-474.

Rehder, H. A. (1). The genus *Turcicula* Dall. Proc. Malac. Soc. London **31** 1955: 222–225 pl. 12.

Rehren, E. Die Abhängigkeit der Atmung vom Sauerstoffpartialdruck bei wirbellosen Wassertieren. Zool. Jb. (Allg. Zool.) 65 1955: 237–266 55 graphs.

Reichel, H. Uber die temperaturabhängigkeit der Spanning im Zustand des Tonus und der Kontraktur im glatten Schliessmuskel von *Pinna nobilis*. Pubbl. Staz. zool. Napoli 27 1955: 73–79 figs. 1–3. [English summary.]

Reichelt, H. (1). Örtliche Anhäufungen mariner Fauna in dem am linken Niederrhein allgemein fossilfreien Horizont über Flöz Katharina auf der Leche Friedrich Heinrich. Neues Jb. Geol. Paläont. B. 4–5 1955: 185–192 text-figs. 1–4 1 table.

Reine, W. J. v. P. see Prud'homme van Reine.

Reish, D. J. & Winter, H. A. The ecology of Alamitos Bay, California, with special reference to pollution. Calif. Fish Game 40 1954: 105-121 1 fig.

Reiss, Z. Remarks on the age of some Late Cretaceous and Early Tertiary stratigraphic units of Israel. Bull. res. Counc. Israel 5B 1 1955: 121-126.

- Rendall, R. Coalfish feeding on Patina pellucida. J. Conch. London 24 1955: 39.
- Rendall, R. (1). Mya arenaria in Orkney. J. Conch. London 24 1955: 39.
- Reyment, R. A. Some examples of homeomorphy in Nigerian Cretaceous Ammonites. Geol. Fören. Stockh. Förh. 77 4 1955: 567-594 text-figs. 1-3.
- Ricci, E. Essai sur la mytiliculture en Tunisie. Bull. Sta. oceanogr. Salammbo No. 52 1955: 39-45 map.
- Richards, H. G. The geological history of the Cayman Islands. Notul. Nat. Acad. Philad. No. 284 1955: 1-11 3 figs. map.
- Richards, H. G. (1). Mollusks from the Mississippi Delta. Notul. nat. Acad. Philad. No. 263 1954: 1-84 figs.
- Richards, H. G. & Ruhle, J. L. Mollusks from a sediment core from the Hudson submarine canyon. Proc. Pa. Acad. Sci. 29 1955: 186–190 1 fig.
- Richardson, E. Neothais smithi (Brazier) 1899. Bull. Auckland Mus. Conch. Cl. No. 8 1952: 13.
- Ricketts, H. O. List of new molluscan names proposed by J. R. le B. Tomlin, J. Conch. London 24 1955: 33–39.
- Ricketts, H. O. & Salisbury, A. E. List of papers on Mollusca and Obituaries of Conchologists published by the late J. R. le B. Tomlin. Proc. Malac. Soc. London 31 1955: 87-94.
- Riedel, A. Molluses of the surroundings of Kazimierz on the Wisla. Fragm. faun., Warszawa 7 1954: 147-185.
- Ringuelet, R. A. Ubicacion zoogeografica de las Islas Malvinas. Rev. Mus. La Plata (Zool.) 6 1955: 419-464.
- Rios, J. M. & Almela, A. Estudio geológico de la Sierra de Ricote, en la región de Mula (provincia de Murcia). (Paleontología). Bol. Inst. geol. Esp. 66 1954: 89-99 (73-83) pls. 1-6 2 geol. maps.
- Risbec, J. Considérations sur l'anatomie comparée et la classification des Gastéropodes Prosobranches. J. Conchyliol., Paris 95 1955: 45–82 22 figs.
 - Roback, S. S. see Wurtz, C. B.
- Roberts, R. V. Recent fossils from Wellington harbour. Bull. Auckland Mus. Conch. Cl. No. 8 1952: 11.

- Roch, G. F. Über ein jugendliches Exemplar von Teredo arenaria (Linné) und die systematische Stellung dieser Art. Beaufortia 5 1955: 15-21 7 figs.
- Roch, [G.] F. (1). Die Terediniden Ost und Westindiens der holländischen Museums-Sammlungen zu Amsterdam und Leiden. Zool. Meded. 34 1955: 125-151 7 figs.
- Roch, G. F. (2). Die holz- und steinzerstoerenden Tiere der Afrikanischen Küstengewässer. Riv. Biol. colon. 13 1955: 71–91.
 - Roger, J. see Dartevelle, E.
- Ronchetti, C. R. Revisione della fauna Neocretacica della Libia: Fam. Ostreidae. Ann. Mus. libico Stor. nat. 4 1953: 111-167 3 pls.
- Ronchetti, C. R. (1). I tipi della "Conchiologia fossile subapennina" di G. Brocchi. [Lyria, Halia, Volutilithes, Cancellaria.] Riv. ital. Paleont. 61 1 1955: 251-266 text-figs. 134-141.
- Ronchetti, C. R. (2). I tipi della "Conchiologia fossile subapennina" di G. Brocchi [Cancellaria, Sveltia, Marginella, Conus]. Riv. ital. Paleont. 61 2 1955: 267–282 text-figs. 142–150.
- Ronchetti, C. R. (3). I tipi della "Conchiologia fossile subapennina" di G. Brocchi [Conus, Genota]. Riv. ital. Paleont. 61 3 1955: 283–298 text-figs. 151–159.
- Ronchetti, C. R. (4). I tipi della "Conchiologia fossile subapennina" di G. Brocchi. [Genota, Cythara, Daphnella, Pleurotomella, Clavatula, Turris.] Riv. ital. Paleont. 61 4 1955: 299-314 text-figs. 160-168.
 - Roon, A. C. de see Raven, C. P.
- Roscoe, E. J. Aquatic snails found attached to feathers of White-faced Glossy Ibis. Wilson Bull. 67 1955: 66.
- Roscoe, E. J. (1). Additional South Dakota terrestrial gastropod records. Amer. Midl. Nat. 54 1955: 511-512.
- Roscoe, E. J. (2). Additional records of *Helix aspersa* Muller from Utah. Gt. Basin Nat. 14 3-4 1954: 90.
- Roscoe, E. J. & L. O. The type locality of *Triodopsis cragini* Call (Pulmonata: Polygyridae). Gt. Basin Nat. 14 3-4 1954: 87-89 1 fig.

Roscoe, E. J. & L. O. (1). Gastropods of the Brighton area, Wasatch mountains, Utah. Nautilus, Philad. 68 1955: 127-131 map.

Rosenberg, G. Einige Beobachtungen in Nordteil der Weyerer Struktur (Nördliche Kalkalpen und Klippenzone). S.B. öst. Akad. Wiss. Abt. I 164 3 1955: 145–161.

Rösler, A. Die Fauna aus den "Bornicher Schichten" (Unter-Devon) des Gemeindè-Steinbruches von Holzhausen a. d. Haide (Blatt Katzenelnbogen/Taunus). Notizbl. Hess. Landesamt. Bodenforsch. 81 1953: 138–153 2 pls.

Rossi, F. & Graziadei, P. Il nervo tentacolare del braccio degli Octopoda non è ganglionare e non è unico. Boll. Zool. 21 1954: 391–394.

Rost, H. A report on the family Arcidae (Pelecypoda). Allan Hancock Pacif. Exped. 20 2 1955: 175–248 6 pls. 17 figs.

Rost, H. & Soot-Ryen, T. Notes on a small collection of Pelecypods from Madagascar. Astarte 11 1955: 1-5 2 figs.

Ruhle, J. L. see Richards, H. G.

Rullán, J. B. Contribuciones a la paleontología del néogeno de Mallorca. Pectínidos (segunda parte). Bol. Soc. esp. Hist. nat. Secc. Geol. 49 1–3 1951: 131–152 2 pls.

Rusconi, C. Fosiles Cambricos y Ordovicios al oeste de San Isidro, Mendoza. Rev. Mus. Hist. nat. Mendoza 8 1-4 1955: 3-64 5 pls.

Russell, H. D. A new clam industry in New England. Nautilus, Philad. 69 1955: 53-56.

Rutland, K. Adelaide, gales and shells. Proc. R. zool. Soc. N.S.W. 1953–54 [1955]: 76.

Rutland, K. & Kerslake, J. Descriptions of the animals of three New South Wales cowries. Proc. R. zool. Soc. N.S.W. 1953-54 [1955]: 75.

Ruzhentzev, V. E. & Shimanskii, V. N. Lower Permian Nautiloidea from the southern Ural. Trav. Inst. paléozool. Acad. Sci. U.R.S.S. 50 1954: 1-152 pls. 1-15 text-figs. 1-28.

Ružička, B. Pectinidae Lamarck of the Silurodevonian of Bohemia. Přírod. Sborn. Ostravsk. Kraje 12 2 1951: 170–177.

Ružička, B. (1). Cypricardinia Hall of the Silurian of Bohemia (Lamellibranchiata). Přírod. Sborn. Ostravsk. Kraje 12 3 1951: 304–312. [English summary.]

Růžička, B. see Dopita, M.

Růžička, B. see Prantl, F.

Růžička, B. see Přibyl, A.

Sacarrão, G. da F. Quelques remarques sur une comparaison des ontogenèsis d'Octopus et d'Eledone. Res. Fac. Cienc. Lisboa (2C) 2 2 1952: 215-222.

Sacarrão, G. da F. (1). Vestiges de nagoires dans le développement des octopodes. Rev. Fac. Cienc. Lisboa (2C) 2 2 1952: 201-214.

Sacarrão, G. da F. (2). Sur la formacion des feuillets germinatifs des céphalopodes et les incertitudes de leur interprétation. Rev. Fac. Cienc. Lisboa (2C) 3 2 1953: 311-364 7 pls. 4 figs.

Sacchi, C. F. Sulla sistematica di alcune Elicelline (Stylomm. Helicidae) della fauna italiana. Annu. Ist. Zool. Univ. Napoli 7 6 1955: 1–16 20 figs.

Sacchi, C. F. (1). Ecologia dei popolamenti di Molluschi psammicoli a Cuma. (Dune del Fusaro e di Licola). Arch. zool. ital. Napoli 38 1954: 195–231.

Sacchi, C. F. (2). Il contributo dei molluschi terrestri alle ipotesi del Ponte Siciliano. Arch. Zool. (ital.) Napoli 40 1955: 49–181 10 figs. [English summary.]

Sacchi, C. F. (3). Contributo alla conoscenza faunistica della Campania. Ricerche malacologiche nella regione sorrentina. Annu. Ist. Zool. Univ. Napoli 6 No. 8 1954: 1–11.

Sacchi, C. F. (4). Fattori ecologici fenomeni microevolutivi nei molluschi della montagna mediterranea. Boll. Zool. 22 fasc. 2 1955: 563-652 3 pls.

Sacchi, C. F. (5). Sul significato zoogeografico della malacofauna di Tremiti a Pianosa. Boll. Zool. 22 1955: 93-111 1 pl.

- Sacchi, C. F. (6). Biogeografia e polimorfisme nelle Euparypha (Helicidae) del Marocco occidentale. (Nota prelimi-Atti Soc. ital. Sci. nat. 94 1 nare). 1955: 85-95.
- Sacchi, C. F. (7). Popolamenti litoral¹ di molluschi terrestri sul medio Adriatico Italiano (Piceno e Abruzzo). Rev. Fac. Cienc. Lisboa (2C) 3 1 1953: 25-52 8 tables. [English summary.]
- Sacchi, C. F. (8). Contributo alla conoscenza faunistica della Campania. Ricerche malacologiche nella regione Sorrentina. Annu. Ist. Zool. Univ. Napoli 6 Nr. 5 1954: 1-38 1 pl.
- Sacchi, C. F. (9). Contributo alla conoscenza faunistica della Campania. Ricerche malacologiche nella regione Sorrentina. Annu. Ist. Zool. Univ. Napoli 6 Nr. 3 1954: 1-13 1 pl.
- Sacchi, C. F. (10). Schizzo delle malacocenosi terrestri sul litorale garganico. Annu. Ist. Zool. Univ. Napoli 6 No. 1 1954: 1–35.
- Sakagami, S. Lower Triassic ammonites from Iwai, Oguno-mura, Nishitama-gun, Kwanto Massif, Japan. Sci. Rep. Tokyo Kyoika Daig. 4 1955: 131-140 2 pls. 3 figs.

Sakai, S. see Imai, T.

- Salisbury, A. E. J. R. le B. Tomlin 1864-1954. J. Conch. London 24 1955: 29-33 portrait.
- Salisbury, A. E. (1). J. R. le B. Tomlin, 1864-1954. Proc. Malac. Soc. London 31 1955: 85-87.
- Salisbury, A. E. see Ricketts, H. O. Sandground, J. H. & Moore, D. V. Notes on the rearing of Oncomelania spp. in the laboratory. J. Parasit. 41 1955: 109–113 3 figs.
- Sarkar, S. S. Sur un genre nouveau Ammonite déroulée. Neocomian, Majorca. C.R. Soc. géol. Fr. 4 1954: 97 - 98.
- Sarwar, M. M. Bionomics and geographical distribution of Limnaea acuminata in the Punjab. Proc. Pakist. Sci. Conf. 5th (3) 1953: 176-177.
- Sarwar, M. M. (1). Distribution of Protostrongylus sp. and Varestrongylus pneumonicus, lungworms of sheep and goats, in the Punjab and bionomics of their intermediate hosts. Proc. Pakistan sci. Conf. 5th Med. & vet. Sci. 1953: 14-15.

- Sarwar, M. M. see Abdussalam, M.
- Sato, H. see Asano, M.
- Sato, T. Hammatoceras de Kitakami, Japan. Jap. J. Geol. Geogr. 25 1-2 1954: 81-106 3 pls. 16 figs.
- Sato, T. (1). Les ammonites recueillies dans le groupe de Kuruma, nord du Japon centrale. Trans. paleont. Soc. Japan N.S. 20 1955: 111-118 pl. 18.
- Sauerzopf, F. Beitrag zur Entwicklungsgeschichte des südburgen-landischen Pannons. Burgenländ. Heimat Bl. 14 1 1952: 1-16 2 pls.
- Sauerzopf, F. (1). Die Planorbiidae aus dem Pannon des Alpenostrandes. Burgenländ. Heimat Bl. 15 2 1953: 49-66 10 pls.
- Sauerzopf, F. (2). Beitrag zur kenntnis der basalen Pannonschichten südburgenländisch - steirischen Becken. Burgenländ. Heimat Bl. 16 4 1954: 145–150 6 figs.
- The world of small Savory, T. H. animals, London (University of London Press), 1955 [slugs p. 144] 160 pp. 17 pls. 30 figs.
- Sawyer, F. C. Books of reference in zoology, chiefly bibliographical. J. Soc. Bibl. nat. Hist. 3 1955: 72-91.
- Saxena, B. B. Physiology of excretion in the common Indian apple-snail, Pila globosa (Swainson). J. Anim. Morph. Physiol. 2 1955: 87–95 2 figs.
- Schäfer, W. Über die Bildung der Laichballen der Wellhorn-Schnecke. Natur u. Volk 85 1955: 92-97 6 figs.
- Schalie, H. v. d. Calvin Goodrich 1874-1954. Nautilus, Philad. 68 1955: 135-142.
- Schalie, H. v. d. & Dundee, D. S. The distribution ecology and life history of Pomatiopsis cincinnatiensis (Lea), an Amphibious operculate snail. Trans. Amer. micr. Soc. 74 2 1955: 119-133 9 figs. & maps.
- Classification of Schenck, H. G. Nuculid Pelecypods. Bull. Mus. roy. Hist. nat. Belg. 10 no. 20 1934: 1-78 pls. i-v.
- Süsswasserschnecken Schermer, E. im Biologieunterricht. Mikrokosmos 44 8 1955 : 174-177 5 figs.

Schindewolf, O. H. Zur Taxionomie und Nomenklatur der Clymenien. Ein Epilog. Neues Jb. Geol. Paläont. B. 10 1955: 417–429 text-figs. 1–2.

Schlechter, A. Über Land- und Wasserschnecken in der Umgebung von Kamenz/Sachsen. Abh. Mus. Tierk. Dresden 22 1954: 88–96.

Schlesselman, G. W. The Gulf Coast oyster industry of the United States. Geogr. Rev. 45 4 1955; 531-541.

Schlickum, W. R. Daudebardia brevipes im Bergischen Land und im angrenzenden Sauerland. Arch. Molluskenk. 84 1955: 181–182.

Schlieper, C. Die Regulation des Herzschlages der Miesmuschel Mytilus edulis L. bei geöffneten und bei geschlossenen Schalen. Kieler Meeresforsch. 11 1955: 139-148 4 pls.

Schmid, F. Die bisherigen Untersuchungen über das Unter/Obermaastricht-Grenzprofil von Hemmoor (Niederelbe), seine Schichtfolge und Leitformen. Mitt. geol. (St.) Inst. Hamb. 24 1955: 75–86 1 table.

Schmidt, G. Stratigraphie und Mikrofaune des mittleren Malm in nordwestdeutschen Bergland mit einer Kartierung am Südlichen Ith. Abh. senckenberg. naturf. Ges. No. 491 1955: 1-76 9 pls.

Schmidt, H. A. Die Besiedlung einer Kulturfläche durch Landschnecken. Arch. Fr. Naturg. Mecklenburg. 1 1954: 193–205.

Schmidt, H. A. (1). Bemerkenswerte Landschnecken im Mecklenburg. Arch. Fr. Naturg. Mecklenburg 1 1954: 206–230.

Schmidt, H. A. (2). Zur Abhängigkeit der Entwicklung von Gehäuseschnecken vom Kalkgehalt des Bodens. Dargestellt bei Oxychilus draparnaldi. Arch. Molluskenk. 84 1955: 167–177 figs. & graph.

Schmidt-Thomé, P. see Ganss, O.

Schmierer, E. T. Obituary notice. Mitt. Berl. Malak. Nr. 4 1953: 1-2.

Schmierer, T. (1). Vertigo moulinsiana Dupuy in Brandenburg (Aus den von Prof. Dr. Th. Schmierer hinterlassen Aufzeichnungen: Zusammengestellt von F. Zimmerman, Berlin). Mitt. Berl. Malak. Nr. 5 1954: 19-27. Schnittmann, F. X. Die Steinbrüche im Malm des Vilstals bei Amberg und seiner weiteren Umgebung. Acta albert. ratisbon. 21 1955: 27-62.

Schouten, A. R. see Altena, C. O. v. R. and others.

Schremmer, F. Bohrschwammspuren in Actaeonellen aus der nordalpinen Gosau. S.B. öst. Akad. Wiss. Abt. I 163 4-5 1954: 297-300 1 pl.

Schwabl, M. Rupertomenia fodiens nov. gen. nov. sp. eine neue Lepidomeniide von der Sudwestküste Schwedens. Öst. zool. Z. 6 1955: 90-146 16 figs.

Schwarzbach, M. Beiträge zur Klimageschichte Islands. I. Allgemeiner Überblick der Klimageschichte Islands. Neues Jb. Geol. Paläont. B. 3 1955: 97–130 text-figs. 1–20.

Schwengel, J. S. New Conus from Costa Rica. Nautilus, Philad. 69 1955: 13-15 pl. 2.

Schwengel, J. S. see Perry, L. M.

Schwetz, J. Quelques considérations sur le problème de la classificationnomenclature des Planorbidae (Planorbinae et Bulininae) de l'Afrique éthiopienne. Ann. Mus. Congo belge N.S. 4to Zool. 1 1954: 274–277.

Schwetz, J. (1). L'influence du milieu sur la taille et la forme du meme Planorbe ou du meme Bulinus. Ann. Soc. zool. Belg. 85 1955: 23-34 8 figs.

Schwetz, J. (2). Sur quelques mollusques africains transmetteurs de bilharziose. Zooleo N.S. No. 31 1955: 64-67.

Sclufer, E. The respiration of Spisula eggs. Biol. Bull. Woods Hole 109 1955: 113-122 5 figs.

Scott, M. I. H. Molucos del biotopo de Cerro Colorado (Salta). Acta zool. Lilloana 6 1948: 229–239 2 pls. 1 fig.

Scott, M. I. H. (1). Moluscos del Noroeste Argentino. Acta zool. lilloana 6 1948: 241–274 1 pl. 13 figs.

Scott, M. I. H. (2). El primer Clausilido de la Argentina. Neotropica 1 No. 1 1954: 1-4 7 figs. [English summary.]

Scott, M. I. H. (3). Nuevas *Epiphrag-mophora* (Gastr. Pulm.) del Noroeste Argentino, Physis **20** 58 1951 ; 252–258 3 figs.

- Scott, M. I. H. (4). Sobre Limnaeidae argentinos (Mol. Pulm.). Physis 20 59 1954: 401–408 6 figs.
- Scott, M. I. H. (5). Una nueva especie de *Bostryx* (Bulimulidae) (Mol. Pulm.). Physis 20 59 1954: 409–413 6 figs.
- Scott, M. I. H. (6). El género Hemisinus (Melaniidae) en la costa fluvial argentina (Mol. Prosobr.). Physis 20 59 1954: 438–443 2 figs.
- Scott, M. I. H. (7). Notas sobre morfologia de *Gundlachia* Pfr. (Ancylidae) (Mol. Pulm.). Physis **20** 59 1954: 467–473 12 figs.
- Scott, M. I. H. (8). Un caso de anomalia de forma en *Tropicorbis* peregrinus (d'Orb.) (Planorbidae). Physis 20 59 1954: 498–499 2 figs.
- Sdzuy, K. Die Fauna der Leimitz-Schiefer (Tremadoc). Abh. senckenb. naturf. Ges. No. 492 1955: 1-73 8 pls. 48 figs.
 - Segura Paguaga, A. see A. K. Miller.
- Seibold, E. Beobachtungen zur Tätigkeit von Bohrmuscheln, Neues Jb. Geol. Paläont. B. 6 1955: 248–251 1 text-fig.
- Seilacher, A. Studien zur Palichnologie II. Die fossilen Ruhespuren (*Cubichnia*). Neues Jb. Min. Geol. Paläont. 98 1 1953: 87–124 pls. 7–13 text-figs. 1–5.
- Seneš, J. Spodnosarmatská fauna pri malej nad Hronom. [Untersarmatische fauna bei Malá n./Hronom]. Geol. sbornik. Slovensk. Akad. 3 3–4 1953: 193–257 pls. 24–33 text-figs. 1–42. [Russian & German résumé.]
- Seneš, J. (1). Prispevok k biocenózam mäkkýšov v morských litorálnych sedimentoch. [Beitrage zu den biozönosen der weichtiere in den meeressedimenten.] Geol. sbornik. Slovensk. Akad. 4 3–4 1953: 883–890 pl. xlvii–l.
- Seneš, J. (2). Stratigrafický a biofaciálny výskum niektorých Neogénnyck sedimentov východného Slovenska na základe makrafauny. [Stratigraphische und biofazielle untersuchung einiger neogener sedimente der ostslowakei auf grund der makrofauna.] Geol. Práce Slovensk. Akad. 40 1955: pls. 1–10 text-fig. & tables. [Russian summary p. 137.]

Senés, J. see Hano, V.

- Shigeno, K. Notes on the biology of the clam *Meretrix lamarckii*. Bull. Jap. Soc. sci. Fish. 21 1955: 218-225 7 figs.
- Shikama, T. Clavagellid fossils from Japan. Sci. Rep. Yokohama Univ. Sect. II No. 3 1954: 63–66 1 pl.
- Shikama, T. (1). On the Tertiary formations of Tomikusa in South Nagano Prefecture. Sci. Rep. Yokohama Univ. Sect. II No. 3 1954: 71–108 5 pls.
 - Shikama, T. see Ozaki, H.
- Shimanskii, V. N. Orthoceratoidea and Bactritoidea from the Sakmarskian and Artinskian beds of the southern Ural. Trav. Inst. paléont. Acad. Sci. U.R.S.S. 44 1954: 1-156 pls. 1-12 & pls. 1-5 in text.
- Shimanskii, V. N. see Ruzhentzev, V. E.
 - Shimizu, D. see Nakazawa, K.
- Shuto, T. Amussiopecten from the Miyazaki group, Miyazaki Prefecture, Japan. Trans. palaeont. Soc. Japan N.S. 20 1955: 101-110 pls. 16-17 text-figs. 1-4.
- Sioli, H. Zur Ökologie des Schistosomiasis-Trägers Tropicorbis (Obstructis) paparyensis F. Baker von Fordlandia (Brasilianisches Amazonasgebeit) und ihre praktische Bedeutung. Arch. Hydrobiol. 51 1955: 153-160.
- Skácel, J. A new finding of fauna in Devonian of Low Jesenik. Přírod. Sborn. Ostravsk. Kraje 13 1–2 1952: 15–22 figs. [English summary.]
- Slanković, S. & Radoman, P. Contribution à la connaissance des gastéropodes Ancylides du lac d'Ohrid. Rec. Trav. Sta. Hydrobiol. Ohrid No. 1-6 1954: 115-122 6 figs.
- Slater, E. The twenty-five known species of land and fresh water mollusks from Great Neck, Long Island, New York, Circ. Mus, biblio, zool, Habana 1954: 1099-1102.
- Sloan, R. E. Paleoecology of the Pennsylvanian marine shales of Palo Pinto County, Texas. J. Geol. 63 5 1955: 412-428 text-figs. 1-23.
- Smith, A. G. Review of the Chitons of the West Coast of North America North of Mexico. Mins. Conch. Club S. Calif. 145 1955: 10-18.

- Smith, A. G. (1). Shells from the bird guano of Southwest Farallon Island, California, with description of a new species of *Liotia*. Proc. Calif. Acad. Sci. (4) 27 13 1952: 383–387 3 figs.
- Smith, A. G. & Hanna, G. D. A rare species of Chiton from Pioneer Seamount off Central California. Proc. Calif. Acad. Sci. (4) 27 14 1952: 389-392 3 figs.
- Smith, E. K. A. Stewart Island Mollusca; additions to the 1939 list of Powell. Bull. Auckland Mus. Conch. Cl. No. 8 1952: 5-8.
- Smith, E. K. A. (1). Stewart Island Mollusca: additions and corrections to my 1952 list. Bull. Auckland Mus. Conch. Sect. No. 9 1953: 5-6.
- Smith, J. E. & Newell, G. E. The dynamics of the zonation of the common periwinkle (*Littorina littorea* (L.)) on a stony beach. J. Anim. Ecol. 24 1955: 35-56 7 figs.
- Snegarova, L. Faune des gastro, podes du Lac d'Ohrid. Acta Mus. maced. Sci. nat. 2 3 1954: 55-64 21 pls.
- Soares, R. de R. L. see Gonçalvez-N. B.
- Solé, N. & J. de. Las formaciones tirrenienses del Cabo de Salou (Tarragona). Mem. Com. Inst. geol. Barcelona 13 1955: 5-35 2 pls. [English & French summaries.]
- Solem, A. New and little known Mexican Helicidae. Nautilus, Philad. 69 2 Oct. 1955: 40-44 pl. 3.
- Solem, A. (1). Mexican mollusks collected for Dr. Bryant Walker in 1926. XI. Drymaeus. Occ. Pap. Mus. zool. Univ. Mich. 566 1955: 1-20 5 pls.
- Soós, L. On Hungarian Daudebardiidae. Acta zool., Budapest 1 1955: 351-360 8 figs.
 - Soós, L. see Bartha, F.
- Soot-Ryen, J. A report on the family Mytilidae (Pelecypoda). Allan Hancock Pacif. Exped. 20 1 1955: 1-174 10 pls. 78 figs.
 - Soot-Ryen, J. see Rost, H.
- Sparks, B. W. Notes on four Quaternary deposits in the Cambridge region. J. Conch. London 24 1955: 47-53 figs. A-C.
 - Stadhouders, A. see Raven, C. P.

- Starmühlner, F. Eine Schnecke wird sesshaft. [Viviparus viviparus.] Aquar. Terrar. Z. 7 1954: 90-93 5 figs.
- Starmühlner, F. (1). Beiträge zur Kenntnis der Molluskenfauna des Arlberges. Öst. zool. Z. 4 1954: 587–632 1 fig.
- Starmühler, F. (2). Zur Molluskenfauna des Felslitorals und submariner Höhlen am Capo di Sorrento. Öst. zool. Z. 6 1955: 147-249 9 figs. & text map.
 - Steinberg, J. see Hand, C.
 - Stelck, C. R. see Warren, P. S.
- Stephenson, L. W. Additions to the fauna of the Raritan Formation (Cenomanian) of New Jersey. Prof. Pap. U.S. geol. Surv. 264-B 1954: 25-43 pls. 6-8.
- Stephenson, L. W. (1). Basal Eagle Ford fauna (Cenomanian) in Johnson and Tarrant Counties, Texas. Prof. Pap. U.S. geol. Surv. 274-C 1955: 53-67 pls. 4-7.
- Stephenson, L. W. (2). Owl Creek (Upper Cretaceous) fossils from Crowleys Ridge, southeastern Missouri. Prof. Pap. U.S. geol. Surv. 274-E 1955: 97-140 pls. 14-24 text-figs. 21.
- Stevanović, P. M. & Pantić, N. Sarmatische Flora und Fauna in den Eisenbahneinschnitten bei Boždarevac (30 km. südl. von Beograd). Ann. géol. Pén. balkan 22 1954: 1–26 pls. 1–5. [In Russian and German.]
- Stewart, G. A. Age relations of the Middle Devonian limestones in Ohio. [Characteristic species listed.] Ohio J. Sci. 55 1955: 147-181 3 figs.
- Stirewalt, M. A. Effort of snail maintenance temperatures on development of *Schistosoma mansoni*. Exp. Parasit. 3 1954: 504-516.
- Stolting, W. H. and others. Fish and shellfish preferences of household consumers. Res. Dep. U.S. Fish Serv. 41 1955.
- Stratton, L. W. Clausilia dubia Draparnaud in the Malham area. (Presidential Address.) J. Conch. London 24 1955: 41–46.
- Strausz, L. Adatok a várpalotai miocén faunához. (Zur Fauna des Mittelmiozäns von Varpalota.) Földt. Közl. 85 2 1955 : 198–210 pl. 12 textfigs. 1–2. [German summary p. 210.]

Strausz, L. (1). Szarmata fauna a karádi mélyfúrásból. (Sarmatische Fauna in der Tiefbohrung von Karád (Ungarn, Kom. Somogy).) Földt. Közl. 85 3 1955 : 381–384. [German summary p. 384.]

42

[Strong, A. M.] [Fauna of West Mexico and the Panamic Province.] Terebridae. Mins. Conch. Club S. Calif. 150 2-5.

[Strong, A. M.] (1). [Fauna of West Mexico and Panama.] Mitridae, Volutidae. Mins. Conch. Club S. Calif. 151 1955: 2-5.

[Strong, A. M.] (2). [Marine fauna of West Mexico and Panama.] Naticidae. Mins. Conch. Club S. Calif. 152 1955: 8-11.

[Strong, A. M.] (3). [Marine mollusca of Western Mexico and Panama.] Columbellidae. Mins. Conch. Club S. Calif. 153 1955: 2-10.

Strong, A. M. see Hertlein, L. G.

Sublette, J. E. The physico-chemical and biological features of Lake Texoma (Denison Reservoir), Oklahoma and Texas: A preliminary study. Texas J. Sci. 7 2 1955: 164–182 5 figs.

Subrahmanyam, T. V., Karandikar, K. R. & Murti, N. N. Marine Gastropoda of Bombay. Part II. J. Univ. Bombay N.S. 21B 3 1952: 26-73 187 figs.

Svagrovský, J. Geologické pomery a fauna severnej časti Košickej kotliny. [Die geologischen verhältnisse und die fauna des nordteiles des Košicer Kessels.] Geol. sborn. Slovensk. Akad. 3 3-4 1953: 259–288 pls. 36–42. [Russ. summary p. 288. German summary p. 293.]

Švagrovský, J. (1). Neogénna fauna východného Slovenska. (Neogene fauna der ostslowakei.) Geol. Práce Slovensk. Akad. 37 1954: 2–64 pls. 1–7. [Russian summary p. 45; German summary p. 54.]

Švagrovský, J. (2). Neogenna fauna východného Slovenska, II. [Die neogene fauna der ostslowakei: II Tiel. Clithon (Vittoclithon) pictus (Férussac) im Miozän der Ostslowakei.] Geol. sbornik. Slovensk. Akad. 6 3-4 1955: 198-226 pls. 4-6. [Russian and German summaries.] Swanker, W. A. Shells collected at Loughborough Lake, Ontario, in July. Nautilus, Philad. 69 1955: 18-20.

[1955]

Sweet, W. C. (1). Cephalopods from the Fremont formation of central Colorado. J. Paleont. 29 1 1955: 71-82 pls. 16-18 1 text-fig.

Sweet, W. C. (2). Some early Paleozoic cephalopods from Wyoming and Utah. J. Paleont. 29 1 1955; 83-85 pl. 18 figs. 1, 3, 4.

Swinnerton, H. H. Proposed use of the plenary powers for the purpose of validating the specific name *minimus* Miller (J. S.) 1826, as published in the combination *Belemnites minimus*. Bull. zool. Nom. 9 1954: 284–285.

Swinnerton, H. H. (1). A monograph of the British Lower Cretaceous Belemnites. Palaeontogr. Soc. [Monogr.] 5 1955: xxxiii-xl, 63-86 pls. 16-18 text-fig. 8.

Taki, I. Fauna of chitons around Japanese islands. Stenoplax of Japan. Bull. nat. Sci. Mus. Tokyo N.S. 1 2 1954: 72-78 4 pls.

Taki, I. (1). Fauna of chitons around Japanese Islands. *Lepidozona* of Japan. Bull. nat. Sci. Mus. Tokyo 2 1 1955: 20–25 2 pls.

Taki, I. (2). New species of *Titiscania*, a rare Archaeogastropod Mollusk, from Hachijō-jima. Bull. nat. Sci. Mus. Tokyo 2 2 1955: 50–53 2 pls.

Taki, I. (3). Distribution of mopalioid chitons in the Pacific. Recoceanogr. Wks. Japan 2 1 1955: 202–209 4 figs.

Taki, I. & Habe, T. Gastrochaenidae in Japan. Kuroda. Ill. Cat. Jap. Shells 2 1955: 1-6 4 figs.

Talmadge, R. R. Variation factors in the *Haliotis*. Mins. Conch. Club S. Calif. 148: 4-6.

Tammes, P. M. L. & Dral, A. D. G. Observations on the straining of suspensions by mussels. Arch. néerl. Zool. 11 1955: 87-112 12 figs. & graphs.

Tamura, M. see Kobayashi, T.

Tasch, P. The concept of psephone-crocoenosis applied to "dwarf" fossil faunas. Amer. Midl. Nat. 53 1955: 205-212.

Tasch, P. (1). Paleoecologic observations on the Orthoceratid Coquina beds of the Maquoketa at Graf, Iowa. J. Paleont. 29 3 1955: 510-518 text-figs. 1-8.

Tauber, A. F. Die fossilen Terediniden der Burgenländischen und Niederösterreichischen Tertiarablagerungen. Wiss. Arb. Burgenland No. 3 [1954]: 1–59 8 pls. 10 figs.

Tauc, L. Réponse de la cellule nerveuse du ganglion abdominal d'Aphysia punctata activé par voie synaptique. J. Physiol. Path. gén. 47 1955: 286–287. See Bull. Soc. biol. Arcachon N.S. No. 8 1956.

Tauc, L. (1). Étude de l'activité élémentaire des cellules du ganglion abdominal de l'Aplysie. J. Physiol. Path. gén. 47 1955: 769-792 16 figs. See Bull. Soc. biol. Arcachon N.S. No. 8

Tavani, G. Fossili pliocenici marini rimaneggiati della Caverna delle arene candide. (Finale Ligure.) Mem. Soc. tosc. Sci. nat. 61A 1954: 11-15.

Tavani, G. (1). Fossili del macigno di Calafuria (Livorno). Mem. Soc. tosc. Sci. nat. 61A 1954: 16-25 1 pl. 2 figs.

Tavani, G. (2). Faune del Quaternario di Rossignano Marittimo (Livorno). Mem. Soc. tosc. Sci. nat. 61A 1954: 241-248.

Taylor, A. E. R. see Fox, H. M.

Taylor, D. W. A new pleistocene fauna and new species of fossil snails from the high plains. Occ. Pap. Mus. Zool. Univ. Mich. 577 1954: 1-16.

Teskey, M. C. The golden cowry. Hobbies 35 4 1955: 75-76.

Teskey, M. C. (1). The Mollusks of Warm Springs, Georgia. Nautilus, Philad. 69 1955: 69-71.

Tesmer, I. H. Restudy of Upper Devonian (Chautanguan) stratigraphy and paleontology in southwestern New York State. [Paleontology note p. 19.] Circ. N.Y. St. Mus. No. 42 1955: 1–22 4 figs.

Thomas, Jr., L. J. The localization of heparin-like blood anticoagulant substances in the tissues of *Spisula solidissima*. Biol. Bull. Woods Hole 106 1954: 129–138.

Thomson, W. P. Notoplax cuneata. Bull. Auckland Mus. Conch. Cl. No. 7 1951: 1-2. Thomson, W. P. (1). Austrosuccinea archeyi (Powell) at Mount Maunganui. Bull. Auckland Mus. Conch. Cl. No. 8 1952: 9-10.

Thomson, W. P. & Jones, A. H. Further records of Austrosuccinea archeyi Powell. Bull. Auckland Mus. Conch. Sect. No. 10 1954: 8-10.

Thornley, G. Additions to the N.S.W. check list. Proc. R. zool, Soc. N.S.W. 1952-53 1954: 32 7 figs.

Thornley, G. (1). Some Helmet-shells (Cassididae) from my collection. Proc. R. zool. Soc. N.S.W. 1952-53 1954: 33-55 2 figs.

Tokioka, T. A measurement on Protoconchs found in the egg-capsules of Syrinx proboscidiferus (Lamarck). Venus, Fukuyama 18 1955: 278–282 2 text-figs.

Tokioka, T. (1). On some plankton animals collected by the Syunkotumaru in May-June 1954. II. Shells of Atlantidae (Heteropoda). Publ. Seto mar. biol. Lab. 4 1955: 227-236 2 pls. 5 figs.

Tokioka, T. (2). Shells of Atlantidae (Heteropoda) collected by the Sôyômaru in the southern waters of Japan. Publ. Seto mar. biol. Lab. 4 1955: 237-250 2 pls. 10 figs.

Tokioka, T. see Oyama, K.

Tomila, G., Koizumi, S., Kurosawa, A. & Kubota, S. Effect of tropolone and its allied compounds on the ciliary movement of the oyster and of *Paramecium*. Sci. Rep. Tôhoku Univ. (4) 21 1 1955: 8-12.

Tomkins, I. R. Elliptio spinosus in the Altamaha River. Nautilus, Philad. 68 1955; 132-133.

Tompkins, P. D. see Hill, H.

Torre y Huerta, C. de la. El genero *Polymita*. Mem. Soc. cubana Hist. nat. **20** 1 1950 : 5-20 11 pls. col.

Trigance, S. Deep water molluses from Timaru area. Bull. Auckland Mus. Conch. Sect. No. 9 1953: 4–5.

Trigance, S. (1). Trawled specimens from off Timaru. Bull. Auckland Mus. Conch. Sect. No. 10 1954: 13.

Trueman, A. & Weir, J. A monograph of British Carboniferous nonmarine Lamellibranchia. Palaeontgr. Soc. [Monogr.] 8 1955: 207-242 pls. 27-29 text-figs. 29-30.

Trussell, P. C., Fulton, C. O., Cameron, C. & Greer, B. Marine borer studies: Evaluation of toxicants. Canad. J. Zool. 33 1955: 327-338 1 pl.

Truyols, Y. J. see Crusafont Pairó, M.

Tsujii, T. Studies on the midgutgland in Invertebrata. IV. Experimental pathological observations of the midgut-gland in snail. Rep. Fac. Fish. Univ. Mie 1 1954: 442–448 pl. 1.

Tsuruta, A. see Chiba, T. and others.

Turner, H. J. How clam drills capture Razor clams. Nautilus, Philad. 69 1955: 20-22.

Turner, R. D. Review of "Ensaio de Catalogo dos Moluscos do Brasil" by F. Lange da Morretes 1949. Occ. Pap. Mollusks, Cambridge, Mass. 1 18 1954: 449.

Turner, R. D. (1). Supplement to John Gould Anthony. Occ. Pap. on Mollusks, Cambridge, Mass. 1 18 1954: 442.

Turner, R. D. see Clench, W. J.

Ullman, H. Observations on a new cercaria developing in *Melanopsis praemorsa* in Israel. Parasitology 44 1954: 1-15 20 figs.

Uozumi, S. On some molluscan fossils from the Palaeogene Wakkanabe formation in Hokkaido. Trans. palaeont. Soc. Japan N.S. 19 1955: 73–80 pl. 12 text-figs. 1–2.

Urbański, J. Molluscs from interglacial sediments at Wola, Warsaw. Bull. Serv. géol. Pologne 69 1954: 141-147. [English summary.]

Vägvölgyi, J. The coenological examination of the mollusks of the Töreki Marsh. Ann. hist.-nat. Mus. hung. N.S. 6 1955: 197-204 2 figs.

Valentine, J. W. Upwelling and thermally anomalous Pacific Coast Pleistocene Mollusca faunas. Amer. J. Sci. 253 1955: 462–474 3 figs.

Valovirta, E. J. Clausilia dubia (Drap.), Suomelle uusi sulkuetana Jyväskylästä. [Finland.] Arch. Soc. zool.-bot. fenn. 9 Suppl. 1955: 401–404 2 figs. [German summary.]

Varma, A. K. Studies on the nature, incidence, distribution and control of nasal schistosomiasis and fascioliasis in Bihar. Indian J. vet. Sci. 24 1954: 11-33 1 pl. 1 fig.

Venmans, L. A. W. C. Notes on Oxyloma verrili (Bland). Basteria 19 1955: 73-75 text-fig.

Venmans, L. A. W. C. (1). Notes on Succinea avara (Say). Basteria 19 1955: 21-22 text-fig.

Venmans, L. A. W. C. (2). Rectification and addition. [For article on Streptostele (Raffraya) horei E. A. Smith.] Basteria 19 1955: 76.

Venmans, L. A. W. C. (3) (see appendix by Watson, H.). Notes on *Streptostele* (*Raffraya*) horei E. A. Smith 1890. Basteria 19 1955: 6-14 text-map & figs.

Verdcourt, B. The cowries of the East African coasts. Suppl. 1. J. E. Afr. nat. Hist. Soc. 22 5 1955: 163-164 1 fig.

Veselinović, D. Le Lias dans la région de "Calcaires de Ravanica" (La zone calcaire à l'ouest de Kucaj). Zborn. Radova geol. Inst. Beograd 7 1954: 47-52 pl. 1. [French résumé p. 52.]

Vicente, F. S. Brachylaemus en enfestación experimental y natural. Rev. ibér. Parasit. 15 1955: 301-320 2 pls.

Vielhauer, W. Beobachtungen an einer eingeschleppten schnecke *Helix* (*Cantareus*) aperta. Natur u. Volk **85** 1955: 289–292 1 fig.

Vilela, H. Objectivos da conquilicultura. Naturalia 5 2 1955 : 85–93 12 figs.

Viola, V. see Baroni, C.

Vlastov, B. V. Sex recognition in the species of Unionidae without the presence of external sexual features. Zool. Zh. 35 1956: 21-28 3 figs. [In Russian.]

Vokes, H. E. Cenozoic pearls from the Atlantic coastal Plain. J. Wash. Acad. Sci. 45 8 1955: 260-262 6 figs.

Vokes, H. E. (1). Notes on Tertiary and Recent Solemyacidae. J. Paleont. 29 3 1955: 534-545.

Voss, C. L. The Cephalopoda obtained by the Harvard-Havana Expedition off the coast of Cuba in 1938-39. Bull. mar. Sci. Gulf & Caribbean 5 2 1955: 81-115 5 figs.

Voss, G. (1). Cephalopoda of the Gulf of Mexico. Fish. Bull. U.S. **55** 1954: 475-478.

Wada, S. K. see Dan, J. C.

Waldén, H. W. The land gastropoda of the vicinity of Stockholm. Ark. Zool. 7 1955: 391-449 33 figs.

Wallace, D. E. see Dow, R. L.

Warren, P. S. & Stelck, C. R. New Cenomanian ammonites from Alberta. Rep. Res. Counc. Alberta No. 70 1955: 63-75.

Warren, S. H. The Clacton (Essex) Channel deposits. Quart. J. geol. Soc. Lond. 111 1955: 283-305 text-figs. 1-2.

Warren, T. P. Collecting Cassididae in the Leigh district [N.Z.]. Bull. Auckland Mus. Conch. Cl. No. 7 1951: 4-5.

Watene, K. P. Ellatriva memorata. Bull. Auckland Mus. Conch. Cl. No. 8 1952: 11–12.

Watlington, F. A. Those prolific snails. [Ampularia cuprina.] Aquarium, Philad. 24 1955: 101.

Watson, H. Appendix Notes on the anatomy of Streptostele (Raffraya) horei Smith. Basteria 19 1955: 14-20.

Watson, H. (1). The names of two common species of *Viviparus*. Proc. Malac. Soc. London 31 1955: 163-174 pl. 8.

Watson, H. (2). The genus Biomphalaria and its relations to other Planorbidae. Rev. zool. Bot. afr. 49 1954: 209-220.

Webb, J. E. & Elgood, J. H. Animal classification, Ibadan (Univ. Press) 1955 pp. ix + 161, frontis.

Weel, P. B. v. The problem of the smooth muscle. Pubbl. Staz. zool. Napoli 27 1955: 10-16.

Weingartner, M. P. Some malacologists of the New York Area. Nautilus, Philad. 69 1955: 48-53.

Wellman, H. H., Reed, J. J., Fleming, C. A. & Hornibrook, N. de B. Marine Pliocene at Resolution Island, Dusky Sound, Fiordland (S156). N.Z. J. Sci. Tech. 35B 5 1954: 378-389 21 figs.

Weir, J. see Trueman, A.

Werner, B. Über die Anatomie, die Entwicklung und Biologie des Veligers und der Veliconcha von *Crepidula fornicata* L. Helgoländ. wiss. Meeresunters. 5 1955: 169-217 22 figs.

Whitehead, E. see Oldfield, E.

Wiborg, K. F. Investigations on zooplankton in coastal and offshore waters of western and northwestern Norway. Fiskeridir. Skr. Havundersok. 11 No. 1 1954: 1–246 102 figs.

Wilbur, K. M. & Jodrey, L. H. Studies on shell formation. V. The inhibition of shell formation by carbonic anhydrase inhibitors. Biol. Bull. Woods Hole 108 1955: 359-365.

Wilbur, K. M. see Jodrey, L. H.

Wilkins, G. L. A Catalogue and historical account of the Banks Shell Collection. Bull. Brit. Mus. (Nat. Hist.) Hist. Ser. 1 3 1955: 69-119 figs. & pls. 14-19.

Wilkinson, A. S. and A. Kapiti bird sanctuary. A natural history of the island. Masterton, N.Z. [List of shells pp. 159–161.] (Masterton Printing Co.) 1952 pp. 190, frontis. 24 pls.

Williams, E. G. Shells from fish. Bull. Auckland Mus. Conch. Cl. No. 8 1952: 1-5.

Williams, G. (1). Seventeen weeks in the life of an Arca novaezelandiae Smith. Bull. Auckland Mus. Conch. Sect. No. 9 1953: 3-4.

Williams, G. (2). Mt. Maunganui, Amphineura or chitons. Bull. Auckland Mus. Conch. Sect. No. 10 1954: 10-12.

Winkler, L. R. A new species of Aplysia on the Southern California coast. Bull. S. Calif. Acad. Sci. 54 1955: 5-7 2 pls.

Winter, H. A. see Reish, D.

Wit, W. F. de. The life cycle and some other biological details of the fresh-water snail *Physa fontinalis* (L.). Basteria 19 1955: 35-44 3 graphs.

Wit, W. F. de (1). The life cycle and some other biological details of the fresh-water snail *Physa fontinalis* (L.) contd. Basteria 19 1955: 45–73 graphs & tables.

Witt, R. M. De. The intrinsic rate and natural increase in a pond snail (*Physa gyrina* Say). Amer. Nat. 88 1954: 353-359 2 graphs & table.

Witt, W. B. De (1). Influence of temperature on penetration of snail hosts by *Schistosoma mansoni miracidia*. Exper. Parasit. 4 1955: 271-276 1 fig.

Witten, H. E. Introduced land molluses of New Zealand. Bull. Auckland Mus. Conch. Sect. No. 11 1955: 1-7.

Woltersdorf, F. Eine Molluskenfauna aus der Niederterrasse von Basel. Verh. naturf. Ges. Basel 65 1 1954: 85-87 1 fig.

Wood, E. J. F. Effect of temperature and rate of flow on some marine fouling organisms. Aust. J. Sci. 18 195: 34-37 6 figs.

Wood, E. J. F. see Dew, B.

Wood, R. A. Toxicity studies on one-, thirty-, sixty day old and adult Stagnicola reflexa. Proc. Ind. Acad. Sci. 64 1955: 291-294 1 fig.

Woolacott, L. New shells from New South Wales. Proc. R. zool. Soc. N.S.W. 1952-53 1954: 37-39 1 pl.

Wright, C. W. see Casey, R.

Wurtz, C. B. The American Camaenidae (Mollusca: Pulmonata). Proc. nat. Acad. Sci. Philad. 107 1955: 99-143 19 pls.

Wurtz, C. B. & Roback, S. S. The nvertebrate fauna of some gulf coast rivers. [Mollusca pp. 185–187.] Proc. Acad. nat. Sci. Philad. 107 1955: 167–206.

Yabe, H. A perisphinctoid ammonite, an ally of *Paraboliceras*, found in the Torinosu series of Zusahara, Sôma-gun, in the Abukuma mountainland. Proc. Imp. Acad. Tokyo 19 9 1943: 582-584 1 fig.

Yabe, Y. see Ichikawa, K.

Yamaguchi, N. & Habe, T. Studies on the Japanese slugs (1). Venus, Fukuyama 18 4 1955: 234-240.

Yamamoto, T. see Oyama, K.

Yamazi, I. Plankton investigation in inlet waters along the coast of Japan. VIII, XVI, XVII. Pub. Seto mar. biol. Lab. 4 1955: 269-320.

Yamazi, I. & Horibata, T. Plankton investigation in inlet waters along the coast of Japan. XVIII. Publ. Seto mar. biol. Lab. 4 1955: 320-336 figs.

Yasuda, J. et al. A note on the spawning season in *Venerupis philippinarum*. Bull. Jap. Soc. sci. Fish. 20 1954: 277-279 1 fig.

Yasuraoka, K. The behaviour of Oncomelania nosophora, the intermediate host of Schistosoma japonicum, to gravity in water. Jap. J. med. Sci. & Biol. 8 4-5 1955: 313-321 figs.

Yasuraoka, K. (1). The behaviour of Oncomelania nosophora, the intermediate host of Schistosoma japonicum, to light in water. Jap. J. med. Sci. & Biol. 8 4-5 1955: 323-329 2 figs.

Yonge, C. M. A note on Arca (Senilia) senilis Lamarck. Proc. Malac. Soc. London 31 1955: 202-208 3 text-figs.

Yoshida, T. see Imai, T.

Yoshimura, K. Studies on the Tropomyosin of squid. Mem. Fac. Fish. Hokkaido 3 1955: 159-176 10 figs.

Young, J. Z. see Boycott, B. B.

Youngquist, W. see W. M. Furnish (1).

Zans, V. A. Verrill's Conch [Strombus gigas verrilli] and the Nicaragua Conch [S. pugilis nicaraguensis] in Jamaica. Nat. Hist. Notes No. 71 1955: 217-219 8 figs.

Zbyszewski, G. L'Aquitanien supérieur de Lisbonne et du Ribatejo. Comun. Serv. geol. Portug. 35 1954: 99-154 pls. 1-13, 16.

Zeiss, A. Pholadomya (Procardia) ziergiebeli n. sp., eine neue Lamellibranchiaten-Art aus dem Callovien Europas. Neues Jb. Geol. Paläont. B. 11 1955: 498-504 1 text-fig.

Ziegelmeier, E. Beobachtungen über den Nahrungserwerb bei der Naticide Lunatia nitida Donovan (Gastropoda Prosobranchia). Helgoländ. wiss. Meeresunters. 5 1954: 1–33 15 figs.

Zilch, A. Die Typen und Typoide des Natur-Museums, Senkenberg. 14. Mollusca Viviparidae. Arch. Molluskenk. 84 1955: 45–85.

Zilch, A. (1). Die Typen und Typoide des Natur-Museums, Senkenberg. 15. Mollusca Cyclophoridae, Cyclophorinae-Cyclophoreae. Arch. Molluskenk. 84 1955: 183–208 pls. 13–15.

Zimmerman, F. Vertigo ronnebyensis Wstld. in Berlin. Mitt. Berl. Malak. Nr. 3 1953: 1-9.

Zimmerman, F. (1). Über den von Vertigo ronnebyensis Westerl. in Berlin-Tegel bewohnten Biotop. Mitt. Berl. Malak. Nr. 5 1954: 13-18. Zimmerman, F. (2). Übersicht der Literatur über die Mollusken von Brandenburg. Mitt. Berl. Malak. Nr. 9 1955: 119–137.

Zimmerman, F. see Schmierer, T.

Zinck, K. Of coral and things. Aquarium, Philad. 24 1955: 340-344.

Znosko, J. see Birkenmajer, K.

Zöbelein, H. K. Funde von Landund Süsswasserschnecken in der chattischen und aquitanischen Bunten Molasse des bayerischen Allgäns. Z. dtsch. geol. Geol. 105 3 1955: 385–395

II.—SUBJECT INDEX

N.B.—A number following an author's name refers to the particular paper in Titles so numbered when more than one work by the same author is recorded.

GENERAL LITERATURE

General Works and Treatises .- Kraken, the legendary giant squid, J. Allan; Collection of molluscs in the oceanographic station of Salammbo, Tunisia, M. Arnould; Possible molluscan tracks in the Namurian of Czechoslovakia. J. Augusta, & V. Pokorny; Use of snails in aquaria, W. Bertholdt; Vertical lobe and memory retention in Octopus, B. Boycott & J. R. Young; Fossil molluscs of Auerbach, Germany, G. Brunner; Hitherto unpublished notes by Dr. C. de la Torre on Polymita from Cuba, J. A. Conde; Study of the Viesca member of the Weches formation (Claiborne, Eocene) Texas, N. M. Curtis; Periodicity with regard to distribution of lamellibranch larvae in Sydney Harbour, B. Dew & E. J. F. Wood; Notes on Trigonia spp. W. M. Draycott; Fossil littoral gastropods as an indication of the position of the littoral zone P.-H. Fischer (5); Census of clams from Pismo Beach, California, J. Fitch (1); Trails and tentacular impressions of fossil cephalopods in America, R. H. Flower (3); The oil content of a plant which will act as a deterrent to animals, E. Frömming (1); On the depth of deposits of Pyritous Ammonites, J. Goguel; Pulmonate gastropods from Roumania, A. Grossu: Notes on three Tellinid species described by V. Bertin,

T. Habe (7); On a giant ammonite donated to the National Science Museum, Tokyo, I. Hayasaka (1); Morphological study of Ammonites flexuosus Buch (Oppeliidae), H. Hölder; Nonmarine mollusca from the Oligocene of Honshu, Japan, Y. Kamada; A study of population genetics of Bracybaena similaris, figs, Komai & Emura; Hunting tree-snails in Cuba, J. W. Lippincott; Setting intensity differences in oysters and starfish, Loosanoff, Engle & Nomejko; Shell collecting in Fiji, F. McCamley.

Monographs.—Cretaceous gasteropoda from Baja California, Mexico, E. C. Allison; English Bathonian Ammonites, W. J. Arkell (3); Cephalopods of central New Caledonia, J. Avias; Analytical comparisons evolution and structural relations, H. B. Baker (1); Distribution and ecology of Littorina obtusata, J. J. Barkman; Neocretaceous lamellibranchs from Libya, C. Baroni & Incitti; Pliocene Pyramidellidae from Florida, P. Bartsch; Jurassic fossils of the Yemen, E. Basse (1); Biology of Spirula, A. F. Braun; On the Neocomian fauna of Algeria and the presence in North Africa of Ammonites of the family Oosterellidae, M. Dalloni: Morphology of the endoceroid fossils, R. H. Flower (1); Cavern fauna of Savona, M. Franciscolo; Bivalvia in Brouns Klassen 3 Abt. 3 Teil 2 1955, F. Haas (4); Shells of the Japanese Islands, S. Hirase; The ammonite Taramelliceras in the Jurassic of south west Germany, H. Hölder; New Devonian gastropod family from Bohemia, R. Horný (1); On the Phylogeny of the Planorbidae, B. Hubendick; Oligocene molluses from Jura and the Haut-Savoie, P. Jodot (1); Malayan Streptaxidae: Discartemon and Oophana, W. S. S. v. B. Jutting; Mollusca from Czechoslovakia, V. Ložek (1); Quaternary molluscs of Czechoslovakia, V. Ložek (2); Pliocene Pelecypoda from Adelaide, Australia, N. H. Ludbrook (1); Freshwater molluscs of Uganda, Mandahl-Barth; Carboniferous cephalopods from Michigan, U.S.A., Miller Garner; Monograph of the Cretaceous Inoceanus of Japan, T. Matumoto; Colour vision in some Euthyneurian snails, G. Mundhenke; Morphology of the fossil pelecypod Astartella, D. Nicol (1); Anentome helena in an aquarium, J. P. van Niel; The economy of the sea with regard to human nutrition, Popovici & Angelescu; Molluscs of the sea-shore, W. Prud'homme van Reine; Ceratites from the German Triassic, H. Penndorf; Marine shells of the west coast of Florida, Perry & Schwengel; The Lamellibranchs of Podern (Aude) S. France, G. Petter-Receveur; Mollusca of the southern islands of New Zealand, A. W. B. Powell (3); Marine molluscs from Sicily, O. Priolo; Opisthobranch molluses from European coasts, A. Pruvot-Fol (1); Nutritive capacity of molluscan blood, P. Rangnekar: Haneomorphy in Nigerian Cretaceous ammonites, R. A. Reyment; Revision of Libyan Neocretaceous Ostreidae, C. R. Ronchetti; Tertiary Planorbiidae of the eastern European Alps, F. Sauerzopf (1); Fresh water snails in the teaching of biology, E. Schermer: Molluscs in relation to marine sedimentation, J. Seneš (1); British Lower Cretaceous Belemnites, H. H. Swinnerton (1); British Carboniferous Naiodites, Trueman & Weir; Molluscs from the stomachs of fish caught in New Zealand waters, E. Williams: Reaction of Oncomelania nosophora to gravity in water, K. Yasuraoka, and to light in water, K. Yasuraoka (1); Conch and other shells for use in an aquarium, K. Zinck.

History.—The Suter Collection of land and fresh-water molluses, R. K. Dell; Mollusks of Linnaeus part 3. Bulla and Voluta, H. Dodge; The Suter Collection of New Zealand molluses, C. Fleming (3); History of Japanese Conchology, Kanamesu; Literature relative to molluses from Victoria, Australia R. C. Kershaw (1); Malacologists of the New York Area, M. P. Weingartner; The Banks Shell Collection, G. L. Wilkins.

Biography.—Merrill E. Champion portrait, Anon; Louise Fischer, 1871–1954, Anon; Capt. Dr. G. Allan Hancock biographical notes and portrait, R. B. v. Kleinsmid; Todd Lopez Moise, T. L. McGinty (2); J. R. le B. Tomlin, 1864–1954, A. E. Salisbury; J. R. le B. Tomlin, 1864–1954, A. E. Salisbury (1); Calvin Goodrich, 1874–1954, H. v. d. Sakalia; E. T. Schmierer, 1879–1953, See Schmierer.

Bibliography.—Vol. 3 (Regnum Lapideum) of 12 ed. Syst. Nat. Lin-

naeus 1768 and corresponding volume in Houttuyn (1785), Gmelin (1793), and Turton, 1806 editions of above work, suppression, Opinion 296; Rejection for nomenclature Geoffroy's Traité Sommaire des Coquilles, 1767, Opinion, 362; Works relating to Cretaceous gasteropoda, particularly from America, E. C. Allison; Stratigraphy and palaeontology of central New Caledonia, J. Avias; Dates of the 'Nautilus', H. B. Baker; Relative to Littorina obtusata, J. J. Barkman: Jurassic fossils of the Yémen, E. Basse (1); Works relative to Miocene fauna, particularly of Africa, E. Dartevelle & J. Rogers; Bibliography relative to Linnaean Genera (Bulla and Voluta), H. Dodge; Works relating to Tertiary Japanese molluses, K. Hirayama (3); On Jurassic Cephalopods, H. Hölder; Abstracts from Wildlife, 1935-1951, N. Hotchkiss; On Roebuck's papers on Scottish nonmarine mollusca, W. & M. Russell Hunter; Works relative to Australian pearl-shells Pinctada spp., J. S. Hynd; Works relating to Belemnitella, J. A. Jeletzky (1); Works relative to Australian mollusca, R. C. Kershaw; Bibliography relative to Quaternary molluscs of Czechoslovakia, V. Ložek (2); On "Review of Australian Seashores", by Dakin and Bennett & Pope, 1952, D. F. McMichael (1); Freshwater molluses of Uganda, Mandahl-Barth; Bibliography of Opisthobranchiata and Nudibranchiata, E. Marcus; Bibliography relative to European Jurassic ammonites, P. L. Maubeuge; Bibliography relative to the biology of Ostrea, R. W. Menzel; Works relating to Carboniferous cephalopods, Miller & Garner; Relative to the genus Busycon, Puffer & Emerson; Relating to the mollusca of the Gulf of Mexico. H. A. Rehder; Molluscan papers and relative essays by the late J. R. le B. Tomlin, Ricketts & Salisbury; Works relating to Cretaceous lamellibranchs, particularly from North Africa, C. R. Ronchetti; On G. Brocchi's "Conchiologica fossile subapennina," C. R. Ronchetti (3); Distribution of land molluscs in Southern Italy and North Africa, C. Sacchi (2); A catalogue with brief notes of books of reference in zoology, F. C. Sawyer; Works relative to British Cretaceous Belemnites, H. H. Swinnerton (1): On a Catalogue of Brazilian mollusca by F. Lange de

Morretes, 1949, R. D. Turner; Works relating to the Solemyacidae, H. E. Vokes (1); Literature on the molluscs of Brandenburg, F. Zimmerman (2).

Taxonomy.—Chitons in the Museo de la Ciudad Eva Perón, Argentina, Ageitos de Castellanos; Key to the Mediterranean Archaeogastropoda, M. Arnould; Systematic review of Czechoslovakian aquatic molluses, J. Brabenec; Taxonomy of Anentome helena, L. J. M. Butot; The lamellibranch genera Trigonioides and Hoffetrigonia belong to the Unionidae and not to the Trigoniidae, L. R. Cox (1); New method for describing fossil Lamellibranchiata, Dopita & Růžička; The sperm as a systematic-phylogenetic character, A. Franzén; Problems in oyster taxonomy, J. Hedgpeth; Taxonomic study of Ammonites flexuosus Buch (Oppeliidae), H. Hölder; Notes on taxonomy of Vitrina, B. Hubendick (4); Revision of the Australian pearl-shells, Pinctada spp., J. S. Hynd; Systematics of Santonian and Campanian Belemnitella, J. A. Jeletzky (1); Systematic list of Tasmanian molluses, R. C. Kershaw; Studies on Australian Charopidae, R. C. Kershaw (2); Australian Charopidae Planate Genera, R. C. Kershaw (3); Japanese Trigonians, Kobayashi & Mori; Revision of New South Wales Turridae, C. F. Laseron; Revision of New South Wales Triphoridae, C. F. Laseron (5); Classification and validity of Hyridella australis (Lmk.) 1819, D. F. MacMichael: Revision of Australian Patellidae and Acmaeidae, J. H. Macpherson; Additions and corrections to the Catalogue of Brazilian molluscs, F. L. de Morretes; Australorbis glabratus, Paraense & Deslandes (1); Australorbis nigricans, W. L. Paraense & Deslandes (2); Morphology of Australorbis centimetralis in comparison with A. glabratus, Paraense & Deslandes (3); Shell structure as a means of classification of the Nuculidae, L. V. de Poel; Liarea from New Zealand, A. W. B. Powell (2); Review of the Classification of Busycon Roding, 1798, Puffer Emerson; Classification and Axodiecal notes of Prosobranch Gastropods, J. Risbec; Description of figured specimens of J. Barrande 1881 (Pectinidae), B. Ružička; Description of J. Barrande's figured specimens of 1881 (Cypricar-dinia), B. Ružička (1); Systematics of

Helicella on the basis of genital characters, G. Sacchi; The Taxonomy and Nomenclature of Clymenia, O. H. Schindewolf; Gastropods of Lake Ohrid, Yugoslavia, L. Snegarova; Genera of British Lower Cretaceous Belemnites, H. H. Swinnerton (1); Key to Cuban Cephalapod, G. L. Voss; Relation of Biomphalaria to other Planorbidae, H. Watson (2); Animal classification, J. Webb.

Nomenclature.—Great Britain, etc. Retention of 34 generic names of nonmarine mollusca, Opinion 335, 336; Rejection of Geoffroy's Traité Sommmaire Coq. ... envir. de Paris, 1767, Opinion 362; Resurrection of Ancient Names, J. Q. Burch (2); Additional names ascribed to C. B. Adams, Clench & Turner (1); List of terms used in gastropod descriptive terminology and discussed, L. R. Cox (2); Status of endoceroid classification, R. H. Flower (1); Pecten nomenclature rectification, G. Grau; Additional names ascribed to Couthouy, R. I. Johnson (1): Nomenclatural notes. Kuroda & Habe (1), Kuroda & Habe (2); Entodina lundi (Mörch, 1871) comb. nov., H. de S. Lopes (1); Nomenclature of the Paleozoic Conocardiidae, D. Nicol (2); New name for an Eocene Hipponix from America, K. V. W. Palmer (2); Remarks on the nomencalture of Melanopsis and Melanosteira, A. Papp (3); The Taxonomy and Nomenclature of Clymenia, O. H. Schindewolf; Planorbidae nomenclature, J. Schwetz; Synonymy of Militidae of W. America, T. Soot-Ryen; Additional names ascribed to Anthony, R. D. Turner (1); On the names of two species of Viviparus, figs., H. Watson (1).

Economics.—Oyster beds of Galicia. Spain, B. Andreu; Uses of oyster shells. J. Baughman; Pearl fishery on Luneburg Heath, Germany, C. R. Boettger (2); The musk rat as an enemy of fresh water mussels, T. Brander; Mussels as food for musk rats, T. Brander (1); The Oysters of the Gulf of Mexico, R. G. Butler; Food bivalves in Setonaikai Inland Sea of Japan, Chiba, Tourata & Ingua; Purification of Oysters, H. A. Cole; Biological and biochemical control of planorbid snails, E. Dias; Bacteriological control of the snail hosts of Schistosomiasis, E. Dias

(1); Bacillus pinottii as a control for snail hosts of Schistosoma, E. Dias & M. Dawood; Abundance of Placopecten magellanicus in the Bay of Fundy, Canada, L. Dickie; Venus mercenaria population in Maine, U.S.A., Dow & Wallace; Spreading of plant-diseases by snails, E. Frömming (2); Herbivorous snails in relation to economically important grasses, Frömming & Plate: Recovery of a community of Placenta placenta in the Gulf of Cutch, India, S. Gokhale: Environmental factors governing infection of Mytilus edulis by Mytilicola intestinalis, B. Hepper; Breeding of Ostrea lurida in Japanese waters, Imai, Sakai & Okada; Ecology of Japanese oyster beds, Ito & Imai; Mollusc contamination controls in New York, M. Krassner; Tapes semidecussata as food, J. Krenmayr; Macrochlamys, a new snail pest from the Punjab, Latif & Qadir; Ostrea edulis in American waters, V. Loosanoff: Dreissena polymorpha and economics in Industry, D. Ludemann; Intertidal edible molluses of Oregon, McCauley & Marriage; Introduction of Gonaxis kiburgiensis E. A. Smith as a control of Achatina fulica, A. R. Mead; Spatfall and behaviour of oyster larvae, J. Medcof; G. of S. Lawrence on resources of the bar clam, Medcof & MacPhail; Tolerance of the wood borer Martesia striata to low salinities, R. Nagabhushanam; Tasmanian scallop fisheries, A. M. Olsen; Studies on Notovola meridionalis from Tasmania, A. M. Olsen; Snail behaviour in relation to different varieties of Chrysanthemum plants, Plate & Fromming; Economy of the sea with regard to human nutrition, Popovici & Angelescu; Use of Placuna shells for window panes in the Philippines, A. Powell (1); Oyster reefs in the Gulf of Mexico, W. H. Price; Culture of Mytilis edulis in Tunisia, E. Ricci; Preservation of wooden harbour constructions with regard to the Teredinidae, G. F. Roch (2); A new industry: the dredging of Arteasa islandica for food, H. D. Russell; Gulf Coast oyster industry of the United States, G. Schlesselman; Report on shellfish preferences of household consumers, W. Stolting & others; Value of toxicants for the marine borer Bankia setacea, Trussell, Fulton, Cameron & Green; Objects of shellfish culture, H. Vilela.

STRUCTURE including HISTOLOGY

General Anatomy.—Doryssa kappleri Vernhout, R. T. Abbot (1); Biomphalaria boissyi as relative to Schistosoma mansoni, E. T. Abdel-Malek; Martelia tanganyicensis, J. Bouillon (1); Possible nautiloid mandibles from the Permian of Arizona, L. F. Brady; Anatomical notes on Spanish snails, F. L. Cabo; Cholinergic innervation of Anodonta heart, Cate, J. Ten; Parallelism in upper Cretaceous ammonites, O. Haas (2); Anatomical studies of Japanese snails, figs., T. Habe (6); Anatomy of Lake Titicaca mollusca, B. Hubendick (2); Choanomphalus, B. Hubendick (5); Comparative anatomy of Radix species, M. Jackiewicz: Chiton heterodon and Chiton peregrinus, E. Leloup (3); Anatomy of $Pomacea\ haustrum\ and\ P$. sordida, H. de S. Lopes; Synolopsis and Anceya Tanganyikan forms, Mandahl-Barth (1); Rhodope veranyi from South America E. & E. Marcus; Philinoglossacea and Acochlidiacea, E. & E. Marcus (2); Anatomy of Tri-toniopsis aurantia, figs., N. T. Mattox (2); Lunella smaragda (Martyn) structure and function of the stomach and sorting caecum, J. E. Morton (1); Pseudochama notes on structure and difference from other genera of Chamidae, figs, N. H. Odhner; Anatomy of Lasaea rubra (Mont.) and Turtonia minuta (Fab.), figs., E. Oldfield; General anatomy of Australorbis glabratus, Paraense & Deslandes (1); General anatomy of Australorbis nigricans, Paraense & Deslandes (2); Studies on Australorbis centimetralis, Paraense & Deslandes (3); Anatomy of Fusus, figs., A. Portmann; Anatomy of molluses, particularly the buccal bulb and the symmetry, A. Pruvot-Fol; Opisthobranch anatomy, A. Pruvot-Fol, (1); Rocellaria (Gastrochaena) cuneiformis R. D. Purchon (1); Anatomy and Classification of Prosobranch Gastropods, J. Risbec; On the animals of some N.S. Wales cowries, Rutland & Kerslake; Morphology of Euparypha (Helicidae), C. Sacchi (6); Rupertomenia fodiens gen. et sp. nov., M. Schwabe; General anatomy of Gundlachia (Ancylidae), M. I. H. Scott (7); Anatomy of Drymaeus, A. Solum (1); Anatomy Daudibardiidae of Hungary, L. Soós; Viviparus viviparus respiratory system, F.

Starmühlner; Anatomy of Streptostele horei E. A. Smith, H. Watson; Anatomical studies of American Camaenidae and Systematics, figs., C. B. Wartz; Anatomy of Arca senilis Lmk., C. M. Yonge.

Nacreous growths.—Giant pearls from Tridacna gigas off South Sumatra, Lay Yoe Soey.

Radula.—Arion lusitanicus Mabille, C. O. v. R. Altena (1); Neritina fiassabensis Binder, Hydrobia guyenoti[†] Binder, Bithynia tournieri Binder, Potadoma vogelii, Binder, Potadoma vogelii Binder, E. Binder; Pseudopartula arborascens Butot, fig., Butot (2); Radula of Planorbis metidjensis, J. Fraga de Azevedo; Anatomical peculiarities in the radula of some Tectibranchs, Gabe & Prenant (1); Orectospira, fig., T. Habe (9); Alderia modesta Loven, figs., Hand & Steinberg; Campoceras hiraseii Walker, B. Hubendick; Tropicorbis Platytaphius canonicus (Cousin), heteropleurus Pils. & Van., Anisancylus crequii Bavay, Littoridina spp. nov. F. Haas, Strombopoma titicacae, Haas, S. gracila Haas, Rhamphopoma spp. nov. F. Haas, Brachypyrgalina carinifera Haas, Heligmopoma umbilicata Haas, B. Hubendick (2); Plesiophysa omata Haas. Indoplanorbis exustus (Desh.), Balinus senegalensis Muller, Physopsis africanus Krauss, Amerianna leopoldi Dupuis, Miratesta celebensis Sarasin, Physastra badae (Bollinger), Camptoceras hirasei Walker, Biomphalaria smithi Preston, Afroplanorbis pfeifferi (Krauss), Austra-Jorbis glabratus heteropleurus Pils & Vanatta, Drepanotiema anatinum Orb, Fossulorbis cultratum Orb, Planorbarius corneus L., Helisoma anceps (Menke), Planorbula jenksi (Car-Promenetus exacuous Say, penter), Menetus cooperi Baker, Carinifex pon-sonbyi Smith, Parapholys effusa Lea, Choancomphalus amacursacceous (Bowes), Planorbis planorbis (L.) Anisus spirorbis (L.), Gyraulus albus (Müll.), Armiger "imbricatus Müller," Hippeutis comand Helicorbisplanatus mearnsi (Bartsch), Segmentina nitida (Müll) Polymytes calathus (Benson), Intha capitis Annandale, Acrorbis petricola Odhner, B. Hubendick (3); Fusilatirus pauli MacGinty, T. L. McGinty; Lunella smaragda (Martyn), figs, J. E. Morton (1); Secretion of teeth by the odontoblastes, A. Pruvot-Fol; Oxyloma verrili (Bland), fig., Venmans; Succinea avara, Venmans.

Muscular system.—C. B. Wurtz.

Alimentary system.—Helix pomatia digestive system, A. Guardabassi; Digestive system of molluses, A. Pruvot-Fol; Alimentary system Camaeidae, C. B. Wurtz.

Reproductive system.—Haliotis lamellosa structure and function of reproductive system, A. Bolognari (1); Genitalia of Planorbis metidjensis, J. Fraga de Azevedo; Ammatorial Organ of Ariophanta maderaspatana, V. K. S. Iyengar; Genitalia of Pomacea haustrum and P. sordida, H. de S. Lopes; Archidoris montereyensis, McGowan & Pratt; calcium distribution in the reproductive system of Pila virens, V. Meenakshi; Genital characters as a basis of the systematics of Helicella spp., C. Sacchi; Genetalia, C. B. Wurtz.

Nervous System.—Innervation of Aplysia heart, H. Leonhardt; Structure of normal and degenerate axoplasma of Sepia officinalis, P. Munari; Nervous system of molluscs, A. Pruvot-Fol; Tentacular nerve of the arm of the Octopoda, Rossi & Graziadei; Nervous System Camaenidae, C. B. Wurtz.

Shell.—Varices on cassids, A. Jones (1); The Turrid Protoconch, C. F. Laseron; Growth of oysters with damaged shell-edges, Loosanoff & Nomejko; Shell structure and form revealed by X-rays, figs., H. B. Moore; On the shell structure of Gryphaea gigus from Akheshi Lake, M. Miyawaki; Varices on cassid shells and the effect on succeeding growth, W. Paul; Shell structure and the classification of the Nuculidae, L. v. de Poel; Measurement of protoconchs of Syrinx probossodiformis (Lmk.), Takioka.

PHYSIOLOGY

General.—Breeding temperatures in Southern and Northern Hemisphere Mytilus, F. E. Allen; ATPase activity of squid mantle muscle (Loligo pealii), G. De Villafranca; Calcium metabolism in fresh water snails, E. Fromming; Lymnaea stagnalis, Horstmann; Action potential in the smooth muscle of snails, N. Ikemoto; Urea-deaminating activity in Corbicula, Ishida, S.; Shell formation: Respiratory metabolism of the oyster mantle, Jodrey & Wilbur;

Distribution of nerves in sub-epithelial conjunctive tissue, M. Kantner; General biology of opisthobranchs, A. Pruvot-Fol (1); Tidal rhythmicity of water propulsion rate in Mytilus, K. Rao; On the heart-beat of Mytilus edulis, C. Schlieper; Biological notes on Meretrix lamarckii, K. Shigeno; Midgutgland in snail, T. Tsujii; Problem of smooth muscle, P. van Weel.

Chemical contents.—Sulphur and nitrogen distribution in Sepia, J. W. Airan & Tosi; Choline in Callista brevisiphonata, M. Asano; Trimethylamine content of Octopus, M. Asano & H. Sato; ATPase activity of squid mantle muscle (Loligo pealii), G. W. De Villafranca; Phosphatases and schiffpositive substances in Lamellibranchs and marine gastropods, M. Genesi; Distribution of various sulfatases in the pearl oyster, Horiguchi & Miyake; Pearl Oyster metabolism. Carbonic anhydrase, K. Ikinaga; Distribution of alkaline phosphatase in mantle tissues of bivalves, Y. Kado; Calcium oxalate in the crystalline style of some molluscs, Kato Kubomura; Nucleic acids and nucleo-proteins in Ostrea laperouse Schrenck, H. Kobayashi; Flavones in some Stylommatophora, V. Kubista; Calcium distribution in Pila virens, V. Meenakshi; Protein catabolism in developing egg of Pila virens, V. Meenakshi (1); Carbonic anhydrase in female genital duct of Pila virens, V. Meenakshi (2); Invertase in digestive gland of Lamellidens, Mukherji Kamungo: Digestive enzymes of Bankia indica, N. Nair; Cellulase activity in crystalline style of Bankia indica, N. Nair (1); ATPase of Mytilus spermatozoa, L. Nelson; Alkaline phosphatase in mantle and pearl sac of Hyriopsis schlegelii, Ojima & Ikenaga; Uptake of phosphate ions by Crassostrea virginica, Pomeroy & Haskin; DNA in nuclei of Papillifera bidens, D. Quattrini; Amino acids in reserve nutritive material of Pila virens egg, K. Ramamoorthi; The oxygen-carrier in the blood of Lamellidens lamellidens, P. V. Rang-neker (1); Isolation of blood anticoagulant from Spirula solidissima, L. Thomas; Tropomyosin of squid, K. Yoshimura.

Reproductory System.—Ammatorial Organ of Ariophanata maderaspatana, V. K. S. Iyengar.

Respiratory System and Respiration.—Gastropod respiration, I. Kretschmann; Respiration of Spisula eggs, E. Sclufer.

Nervous system.—The vertical lobe and memory retention in Octopus, B. Boycott & J. Z. Young; Cholinergic innervation of Anodonta heart, Cate J. Ten; Secretion in Illex brain, H. Haefelfinger; Neurosecretion and incretory glands in a tectibranch molluse, H. Lemche; Normal and degenerate axoplasma of Sepia officinalis, P. Munari; Responses of the nervous cell of the abdominal ganglion of Aplysia punctata, L. Tauc; On the cells of the abdominal ganglion of Aplysia, L. Tauc (1).

Alimentary system and Alimentation.—Physiology of digestion, A. Guardabassi; Physiology of digestion in Pila virens Lmk., V. R. Meenakshi (3); Physiology of digestion in Melania crenulata figs., V. R. Meenakshi (4); Digestive enzymes of Bankia indica, N. Nair; Cellulase activity in crystalline style of Bankia indica, N. Nair (1).

Excretory and Secretory organs.— Secretory material of ammatorial organ of Ariophanata maderaspatana, V. K. S. Iyengar; Excretion in Pila globosa, B. Saxena.

Experimental Studies.—Learning in Octopus, B. Boycott; Light perception in Arcacea, R. Braun; Action of acetylcholine and eserine on heart and intestine of Anodonta cygnea, J. Ten Cate & M. S. Rusinch; Use of radioactive plankton to measure water filtration in Pecten irradians. W. Chipman & J. G. Hopkins: Aerosome reaction in pelecypod spermatozoa, J. Dan S. N. Wada; Effect of heating on the mortality of snails, A. N. Dingle & C. B. Nash; Use of heart of Anodonta for estimation of 5-hydroxytryptamine. R. Fange; DNA content of spermatozoa with regard to the activation of the egg of Spisula solidissima, Ito & Leuchtenberger; Effects of glycine and metals on metabolism of oyster spermatozoa. S. W. Jeffrey; Influence of diluted seawater on pearl oyster heart-beat. Kawamoto & Motoki; Influence of diluted sea water on the pearl oyster. Kawamoto & Moloki (1); Gastropod respiration, I. Kretschmann; Experiments on relative streamlining of coiled cephalopod shells, Kummel & Lloyd; Neurosecretion and incretory glands in

a mollusc, H. Lemche; Carbonic anhydrase in the formation of egg-shell of Pila virens, V. R. Meenakshi (2); Invertase in digestive gland of Lamellidens, Mukherji & Kanungo; Influence of acetylcholin, eserine and curare on the anterior byssus muscle of Mytilus, A. Punt: The oxygen carrier in the blood of Lamellidens lamellidens, P. V. Rangneker (1); Head malformations caused by centrifuging Limnaea stagnalis eggs, Raven & Beenakker; Morphogenetic effects of heat shock on Limnaea stagnalis eggs, Raven, Roon & Stadhauders; Pathological observation of the midgut-gland in snail, T. Tsujii; Effect of temperature and rate of flow on marine molluscs. E. F. Wood: Experiments on Toxicity of Snails, R. A. Wood.

DEVELOPMENT EVOLUTION AND GENETICS

General.—Biological influences on the postembryonic development of Eobania vermiculata, E. Frömming (3); Fertilization of Mytilus galloprovincialis H. Heldt.

Life History.—Life history of Hyridella australis, I. D. Hiscock; DNA content of spermatozoa with regard to activation of egg of Spisula solidissina, Ito & Leuchtenberger; Development of Kerguelenella stewartiana, G. A. Knox; Chromosome numbers of two Japanese species of Succinea, T. Koyama; Spawnballs of Buccinum undatum, W. Schäfer; Biological notes and life cycle of Physa fontinalis (L.), de Wit.

Growth.—Shell growth of Margaritifera margaritifera, C. Boettger (2); Growth in Oysters, measurement and weight, J. L. McHugh; Growth of parasitized oysters, Menzel & Hopkins; Growth of Littorina planaxis, W. North; Changes in the boring plates of Teredo in the course of growth, P. Rancurel (1).

Oogenesis and Oology.—Oology and Naticidae, M. Amio; Ovogenesis in Aplysia depilans, A. Bolognari; Occyte development of Haliotis lamellosa, A. Bolognari; Oviferous capsules of Murex trunculus, F. Dulzetto; Egg masses of the Naticidae, M. Giglioli; Protein catabolism in egg of Pila virens, V. Meenakshi; Rôle of carbonic anhydrase in the formation of the egg-shell of

Pila virens, V. R. Meenakshi (2); Amino acids in reserve nutritive material of Pila virens egg, K. Ramamoorthi; Cytometric observations on occytes, F. Ranzoli; Head malformations obtained by centrifuging Limnaea stagnalis eggs, Raven & Beenakker; Effects of heat shock on Limnaea stagnalis eggs, Raven, Room & Stadhouders; Respiration of Spisula eggs, E. Sclufer.

Spermatogenesis.—Acrosome reaction in pelecypod spermatozoa, J. Dan & S. N. Wada; Spermiogenesis and sperm morphology of certain molluses, A. Franzén; Tapes decussatus, H. Heldt; Effects of glycine and metals on metabolism of oyster spermatozoa, S. W. Jeffrey; Chromosome changes in spermatogenesis of Bithynia tentaculata (L.), H.-G. Keyl; Spermatogenesis of Anadenus altivagus, Nath & Chopra; APT-ase of Mytilus spermatozoa, L. Nelson; Spermatogenesis in Deroceras reticulatum, D. & N. E. Pelluet.

Embryonic stages.—Cleavage of Bithynia tentaculata L., O. Hess; Embryonic development of Bithynia tentaculata L., O. Hess (1); Gastrula of Bithynia tentaculata, O. Hess (2); Early embryology of Archidoris montereyensis, McGowan & Pratt; Embryology of Fusus, figs., A. Portmann; Embryology and development of some Melaniidae, K. Ramamoorthi (1); Comparison of Octopus and Eledone, G. da F. Sacarrão; Germ layer formation in cephalopods, G. F. Sacarrão (2); Embryology of Crepidula, B. Werner.

Larval stages.—Egg masses and larvae of some Naticidae, M. Amio; Lepidopleurus asellus, M. Christiansen; List of molluscan larvae observed at Naples, Knight-Jones; Larvae and late Larval stages of Octopus macropus Risso, W. J. Rees (1).

Development of Fins.—Fin-vestiges in the development of the Octopoda, G. da F. Sacarrão (1).

Shell.—Development and coiling of Nautiloids, figs., R. H. Flower; Shell formation: Respiratory metabolism of the oyster mantle, Jodrey & Wilbur; Growth of oysters with damaged shelledges, Loosanoff & Nomejko; Inhibition of shell formation by carbonic anhydrase inhibitors, K. Wilbur.

Hybridization.—Among molluses from North America, P. Bartsch.

Growth.—Rate of growth of Gastropods, Dehnel; Rate of growth in *Paryphanta busbyi*, R. K. Dell (8).

Variation: General.—Distribution and variation of Cepaea, C. Boettger (3); Variation of Littorina in the North Atlantic, J. Knudson (1); Variation of Helicella ericetorum and H. obvia, H. Melchinger; Factors affecting size in Littorina, W. North; Variation of molluses from mediterranean mountains, C. Sacchi (4); Haliotis, R. R. Talmadge.

Variations in form.—Aberrant cowries of Keppel Islands, Queensland, Australia, J. Kerslake; Environmental influence on the size and form of the shells of Planorbis and Bulinus, J. Shwetz (1); Irregular form of Tropicorbis peregrinus, M. I. H. Scott (8).

Variation in Colour.—Polymita and new subspecies, C. de la Torre.

Terautology.—Abnormality in tentacles of Euhadra, Kishimoto.

Sex.—Sex determination in Unionidae deprived of external sexual features, Vlastov.

Evolution.—Of the family Calyptraeidae, P. H. Fischer (2); Evolution and saltation in Nautiloid coiling, R. H. Flower; Evolution of the endoceroid fossils, R. H. Flower (1); Evolution of Santonian and Campanian Belemnitella, J. A. Jeletzky (1); The Trigoniinae and their derivatives, Kobayashi & Mori; Evolution of Peroniceratidae, T. Matsumoto (2); Evolution of vermetid gastropods, J. E. Morton.

ECOLOGY

HABITAT, HABITS, FIELD NOTES AND TECHNIQUE

General.—Ecology of the pearl fishery of Luneberg Heath, Germany, C. Boettger (2); Ecology of molluses from Czechoslovakia, J. Brabenec (1); Ecology of Mozambique molluses, J. M. Braga; Two gastropods from Basle Botanic Gardens, A. C. von Bruggen (1); Anentome helena, the freshwater snail, L. J. M. Butot (1); Ecology of Amphidesma ventricosa from New Zealand, R. M. Cassie; Placopecten magellanicus abundance in the Bay of Fundy, Canada, L. Dickie; Molluses from Sauerland, Germany, H. Dittmar; Venus mercenaria, population in Maine,

U.S.A., R L. Dow; Biological influences on the postembryonic development of Eobania vermiculata, E. Frömming (3); observations on Acteon tornatilis, Fretter & Graham; Environmental factors governing infection of Mytilus edulis by Mytilicola intestinalis, B. Hepper; Ecology of Japanese oyster beds, Ito & Imai; Ecology of molluses from the Balkans, S. Jaeckel (1); Molluses of the central German mountains, S. Jaeckel (4); The cowry shell (Cypraea moneta and C. annulus) in African art, M. Jeffreys; Faunal analysis of molluses from raised beach deposits in Japan, S. Kanno (1); Shore biological communities at Rio de Janeiro, W. Kegel; Limnology of Rio Grande do Sul. Brazil, H. Kleerekoper: High shore level limpets at Aberystwyth, J. Lewis; Ecology of molluses from Czechoslovakia, V. Ložek (1); Ecology of molluscs from Värmland, Sweden, Lundgren; Hydrobiology of Majorca, R. Margalef: Ecological study of Unionidae, M. R. Matteson; Note on Megalobulimus paranaguensis, F. L. de Morretes (2); Erosive activities of Littorina, W. North; Mode of Life of Lassea rubra (Mont.) and Turtonia minuta (Fab.), E. Oldfield; Studies on Notovola meridionalis from Tasmania. A. M. Olsen; Effect on invertebrate fauna of salinity changes in bays of central Texas, R. H. Parker; Molluses of the Harze mountains, Pfitzner & Jaeckel; Snail behaviour in relation to different varieties of Chrysanthemum plants, Plate & Frömming; Ecological notes on Busycon, Puffer & Emerson; Ecology of molluses from Cuma, Naples, C. Sacchi (1); Distribution of land molluses in Southern Italy and North Africa, C. Sacchi (2); Ecological factors in the microevolution of molluses from mediterranean mountains, C. F. Sacchi; Terrestrial and littoral molluses of the Italian Adriatic, C. F. Sacchi (7); Fauna and ecological value of landshells of the Gargano Peninsula, C. F. Sacchi (10); Colonization by land snails, H. Schmidt; Shells of Planorbis and Bulinus unsuitable as classification basis due to environmental differences in form, J. Schwetz (1); Ecology of gastropods of Lake Ohrid, Yugoslavia, L. Snegarova; Marine gastropods of Bombay, T. V. Subrahmanyan, Karandiker & Murti; Molluscan communities of the Töreki Marsh, Hungary, J. Vágvölgyi.

Habitat.—Mytilus californianus, S. Berry (3); Molluscs from deep-water rocks in New Zealand, R. K. Dell (1); Habitats of the Corbicula japonica group, A. Fuji; Habitats of snail hosts of Bilharzia and liver-fluke, A. Mozley.

Habits: General.—Habits of Pandora inaequivalvis (L.) figs, M. F. and J. A. Allen; Habits of, and observations on Balcis devians (Monts) and Balcis alba (Costa), V. Fretter; Arca senilis, habits, C. M. Yonge.

Breeding habits.—Breeding temperatures in Southern and Northern Hemisphere Mytilus spp., F. E. Allen; Lepidopleurus asellus, M. Christiansen; Periodicity in relation to distribution of lamellibranch larvae, B. Dew & E. J F. Wood; Breeding of Ostrea lurida in tanks, Imai, Sakai & Okada; Breeding experiments of Bradybaema similaris, Komai & Emura; Sexual dimorphism in Strombus auris-dianae, J. Laseron; Setting intensity differences in oysters and starfish, Loosanoff, Engle & Nomejko; Loligo opalescens, J. McGowan; Spatfall and behaviour of oyster larvae, J. Medcof; On fish-eggs deposited in dead bivalve mollusc shells, I. S. R. Munro; Self and cross fertilization in Australorbis glabratus, W. L. Paraense; Reproductive isolation between Austraglabratus and A. nigricans, Paraense & Deslandes; On copulation in Limax, B. Peyer; On spawnballs of undatum, W. Schäfer; Buccinum Spawning of Venerupis philippinarum, J. Yasuda.

Feeding Habits and Food.—Coprophagous Palaeozoic gastropods, A. L. Bowsher; Water filtration of Pecten irradians, W. Chipman & J. G. Hopkins; The oil content of a plant which will act as a deterrent to animals, E. Frömming (1); Molluscan diets (Presidential address), A. Graham; Feeding and control of Amphidesma ventricosum in New Zealand, A. M. Rapson; Mollusk as food of fish, R. Rendall; Feeding—straining of suspensions by Mytilus, Tammes & Dral; How Polinices capture Ensis, H. J. Turner, jun.; Lunatia nitida, E. Ziegelmeier.

Boring habits.—Boring attack on Nassarius, Alexander; On Rock boring gastropods, P. H. Fischer (1); Biology of Rocellaria cuneiformis, R. D. Purchon (1); Stone and wood-boring

molluses, G. Roch (2); Boring habits of Lithodomus, E. Seibold; Lunatia nitida, E. Ziegelmeier.

Movement and Locomotion.—Measurement and speed, M. J. Brown; Tricolia pullus (L.), V. Fretter (1); Swimming movements of lamellibranch larvae as guides to their classification, Knight-Jones.

Enemies.—Nassarius thersites bored through by a predator, F. E. S. Alexander; Sea-stars as enemies of molluscs, F. C. Benitez; The musk rat as an enemy of fresh water mussels, T. Brander; Mussels as food for musk rats, T. Brander (1); Snail-eating behaviour of thrushes and blackbirds, D. Morris.

Parasitism and Commensalism.—Biomphalaria boissyi and Schistosoma; Anatomy, E. T. Abdel-Malek; Limnaea acuminata form rufescens as inter-mediate host of Schistosoma spp. in Pakistan, M. Abdussalam; Cercaria of Parorchis acanthus v. australis parasitic in snails, L. M. Angel; Adaptations of Palaeozoic gastropods attached to echinoderms. A. L. Bowsher; Control of planorbid snails, F. Dias; Bacteriological control of the snail hosts of Schistosomiasis, E. Dias (1); Bacillus pinottii as a control for snail hosts of Schistosoma, E. Dias, M. Dawood; Susceptibility of Planorbis to Schistosoma, J. Fraga de Azevedo; Molluscs as hosts of Schistosoma in Mozambique, J. Fraga de Azevedo; Polychaetes in pallial cavity of *Mytilus galloprovincialis*, **S. Genovese**; The use of Cupric ricinoleate to kill molluses, Gonçalvez & Soares; Glochidium of Hyridella australis parasitic on freshwater fish, I. D. Hiscock; Effects of metacercaria on Vivipara bengalensis, Lal & Baugh; Snail hosts of Schistosoma in Gambia, McCullough & Duke; Seasonal infection by larval trematodes of Cerithidea californica Holdmann, W. E. Martin; Lymnaea obrussa as am intermediate host of Fasciola hepatica. L. Mazzotti: Growth of parasitized oysters, Menzel & Hopkins; Habitats. of snail hosts of Bilharzia and liverfluke, A. Mozley; Development of albino-Australorbis susceptible to Schistosoma infection, W. L. Newton; Pyrazus australis as host of Schistosoma in New South Wales, Australia, E. C. Pope: Parasistology and Oyster, Ray Chandler; On rearing Oncomelania in

the laboratory, Sandground & Moore; Helicella and Macrochlamys as intermediate hosts of sheep lungworm, M. Sarwar (1); Molluscs as carriers of bilharziosis in Africa, J. Schwetz (2); Molluscan Schistosoma carriers in Brazil, H. Sioli; Parasitology experiments, M. A. Stirewalt; Cercaria found in Melanopsis praemorsa from Israel, H. Ullman; Molluscan hosts of nasal schistosomiasis in Bihar, India, A. Varma; Limax and Helix as intermediate hosts of the Trematode Brachybaemus, F. S. Vicente; Influence of temperature and infection by Schistosoma, W. B. De Witt (1); Reaction of Oncomelania nosophora, intermediate host of Schistosoma japonicum, to water, K. Yasuraoka; gravity in Reaction of Oncomelania nosophora, intermediate host of Schistosoma japonicum, to light in water, K. Yasuraoka (1).

Effect of Temperature.—Adaptation to temperature by Helix pomatia, P. Blazka; On tissue hydration in Australorbis, T. V. Brand; Mortality of Ostrea lurida at low temperatures, H. Davis; Effect of heating on the mortality of snails, A. N. Dingle & C. B. Nash; Freezing effects in intertidal molluses, J. Kanwisher; Time and temperature relation, P. R. Orr; Temperature-tension relationship in Pinna adductor muscle, H. Reichel.

Effect of Light.—Light perception in Arcacea, R. Braun; Effect of light, M. J. Brown; Spermatogenesis in Deroceras, D. & N. E. Pelluet.

Vision.—Colour vision in some Euthyneurian snails, G. Mundhenke.

Fluorescence.—Of some gasteropoda, P. H. Fischer (3); Fluorescence of gastropod opercula, P.-H. Fischer (6).

Effect of oxygen.—In invertebrates, H. R. Fox; Tolerance of oxygen, Fox & Taylor.

Effect of Toxins.—Effect of acid on the egg of Bithynia codiella, C. Attardo; Action of acetylcholine and eserine on heart and intestine of Anodonta cygnea, J. Ten Cate & M. J. Reesinck; The action of potassium cyanide on the development of Physa rivularis Ph., V. Mancuso; Action of sodium azide on developing Physa rivularis eggs, V. Mancuso (1); Effect on the oyster heart-beat of a toxic substance derived from the sea urchin Toxopneustes

pileolus, K. Okada; Effect of tropolone on ciliary movement of the oyster gill, G. Tomita, Koizami, Kurosava & Kubata; Value of toxicants for the marine borer Bankia setacea, P. Trussell, Fulton, Cameron & Greer.

Effect of Salinity.—Tolerance of Martesia striata to low salinity, R. Nagabhushanam.

Duration of Life.—Age determination figs. and graphs, H. H. Haskin; Death of pearl oysters caused by bacteria, Kotake & Miyawaki.

Association and effect.—Symbiosis involving Mya arenaria and Melgola manhattensis, F. A. Aldrich; Association of Palaeozoic gastropods and echinoderms, A. L. Bowsher; The association of the fossil, Conularia with a sponge Guadalupia and three brachiopods, R. M. Finks; Effects of metacercaria on Vivipara bengalensis, Lal & Baugh; Traces of boring sponges in Actaeonella from the Austrian Jurassic, F. Schremmer.

Geological aspect.—New method of describing fossil Lamellibranchiata, Dopita & Růžička; Concretions from the German Oligocene, K. W. Geib; Molluses as indicators of ecological conditions, R. W. Imlay; Fossil freshwater gastropods from Greece, A. Papp (1).

Regeneration.—Experiments on regeneration Arion rufus L., figs., P. Laviolette.

Technique.—Sulphur and nitrogen distribution in Sepia, J. W. Airan & Tosi; Trimethylamine content of Octopus, M. Asano & H. Sato; Sampling technique for estimation of abundance of Amphidesma ventricosum, R. M. Cassie; Serological technique, J. N. Cathern: Biological and biochemical control of planorbid snails, E. Dias; Bacteriological control of the snail hosts of Schistosomiasis, E. Dias (1); Bacillus pinottii as a control for snail hosts of Schistosoma, E. Dias & M. Dawood; On rearing Oncomelania in the laboratory, Sandground & Moore; For sorting molluses from the stomach contents of fish, E. Williams; Method for collecting and mounting chitons, G. Williams (2).

Conchometry.—Of gastropod as a function of latitude, Dehnel; New method of describing fossil Lamelli-

branchiata, M. Dopita & Růžička; Helicella ericetorum and Helicella obvia, H. Melchinger; Methods for measuring shells, J. Parodiz (3); Mytilidae of the Velera III and IV, T. Soot-Ryen; Measurement of protoconch in egg capsules of Syrinx proboscidiferus (Lmk.), Tokioka.

Effects of soils.—Ecological factors of soil affecting Virginian land snails, J. B. Burch.

Ethnology.—Conchology as an aid to Archaeology and Ethnology, H. Boekelman; Use of shells in early civilizations, P. Fischer; Use of *Placuna* shells for window panes in the Philippines, A. Powell (1).

ZOOGEOGRAPHY

Dispersal and Distribution,—Distribution and variation of Cepaea, C. Boettger (3); Distribution and Ecology of Mollusca of Point Barrow, Alaska, G. E. MacGinitie; Distribution of Helicella ericetorum and H. obvia, H. Melchinger; Distribution of Cephalopoda in the Eastern Mediterranean, W. J. Rees; Distribution of land molluscs in Southern Italy and North Africa, C. Sacchi (2); Capri-Sorrento-Amalfi distribution of land molluscs, figs., C. F. Sacchi (8); Distribution of Limnoea acuminata in the Punjab, M. Sarwar; Distribution, ecology and life history of Pomatiopsis cincinnatiensis (Lea), Schalie & Dundee; Distribution of mopalioid chitons in the Pacific, I. Taki (3).

General Lists & Catalogues .- Organic matter in relation to distribution of pelecypods, R. E. Bader; Catalogues of Mollusca of Portugal, J. O. de Castro; Additional names to the Catalogue of Truncatellidae (Occ. Pprs. No. 13), Clench & Turner; New names of species of 'Dentalium' compiled from the Zoological Record for 1899-1950, W. K. Emerson; The food of Mollusca a copious illustrative list of the diet of many species, A. Graham; List of Japanese Acmaeid Limpets, T. Habe (3); Mollusca collected at the Galapagos Islands, 1931-32 'Velero III' Exped., Hertlein & Strong (1); Report on the molluses of the Clacton Channel deposits, D. F. W. Baden-Powell; Common sea shells of Los Angeles County coast, Hill & Tompkins; List of molluses from Tryphena Bay, Gt.

Barrier Island, New Zealand, A. Hipkins; New Sections (genera) of Rissoidae of New South Wales, T. Iredale; Catalogue of the genus Liguus in Cuba, W.I., M. Jaume; Gastropods of the "Gr's de Taouilah" Cretaceous, Yemen, J.-P. Lefranc; Catalogue and notes on the Busyconinae, Puffer & Emerson; Cephalopods found East of 23° E. in the Mediterranean, W. J. Rees; Fossils from the Hudson submarine Canyon, Richards & Ruhle; New molluscan names proposed by J. R. le B. Tomlin, H. O. Ricketts; List of British Slugs and key to identification, T. H. Savory; List of specific names used for West American mytilids, T. Soot-Ryen; List of species from Quaternary deposits in the Cambridge area, B. W. Sparks; Mollusca Cape Sorrento, F. Starmühler (1); Species of the family Calyptraeidae, B. Werner; Mollusca of Kapiti Island, N.Z., A. S. & A. Wilkinson; Senkenberg Museum types of Cyclophoridae, A. Zilch (1); Species, subspecies and varieties of Viviparidae, A. Zilch.

Europe.—Stratigraphical list of Radiolites of the Pyrenees, G. Astra; Mollusca of Mar Menor, map, F. M. Cabo (1); List of marine molluscs from Whitstable, England, G. Newell; Mollusca in Hertfordshire, notes on issue of list by L. W. Stratton, [A. E. Ellis]; Mollusca from Czechoslovakia, V. Ložek (1); List of fossils from the Rumanian Lias, G. Răileanu; The mollusca of the Clacton Channel deposits, Essex, S. H. Warren.

Asia.—Marine mollusca of North China, Hsi, Chi, & Li; List of mollusca from dredgings between Cap Djinet and the mouth of Oued Sebaou, P. Muraour; Indo-China-Nassidae, M. Marche-Marchad; Species of Cymatiidae and Bursidae of Indo-China, M. Marche-Marchad (1); Land and water snails of neighbourhood of Kamenz/Sachsen, A. Schlechter.

Africa.—Molluses from dredgings of Algeria, R. Dieuzeide.

North America & West Indies.—Index and catalogue of Cuban molluses, C. Aguayo; Cretaceous molluses from Baja California, Mexico, E. C. Allison; Land snails of Hanover, Henrico and Chesterfield Counties, Virginia, J. B. Burch (1); Minnesota land snails, G. Dawley; Land and Freshwater gastropoda of Cape Breton, Canada, G. K.

MacMillan: San Pedro Basin, California, list of Mollusca identified from benthos samples, N. J. Mattox (1); Mollusca of the Gulf of Mexico, H. A. Rehder; Marine Mollusca of West Mexico and Panama, [A. M. Strong]; Mitridae and Volutidae of West Mexico and Panama, [A. M. Strong] (1); Naticidae of Western Mexico and Panama, [A. M. Strong (2); Columbellidae of Western Mexico and Panama, [A. M. Strong] (3); Shells collected on Loughborough Lake, Ontario, W. A. Swanker; The mollusca of Warm Springs, Georgia, M. C. Teskey (1); Cephalopoda of the Gulf of Mexico, G. Voss (1); Mollusca of some Gulf coast rivers, Wurtz & Roback.

South America.—Species of Argobuccinum, A. Carcelles (1); Check list of Tertiary and recent Australasian Nuculanidae, R. K. Dell (6); Lists of molluscan species and stations in Lake Titicaca, F. Haas (6).

Australia.—Additions to the checklist of molluses from New South Wales, Australia, G. Thornley.

Japan.—List of Japanese *Pleurotomaria* species, **Kuroda**; List of Japanese species of the Conidae, **T. Kuroda** (1).

New Zealand.—List of molluses from Stewart Island, New Zealand, E. K. Smith; List of deep water molluses from Timaru area, New Zealand, S. Trigance.

Oceanic Islands.—Palaeontological index of species present in Central New Caledonia, J. Avias; Mollusca of Palau Panaitan, L. J. M. Butot (2); Land molluscs from the Solomon Islands, R. K. Dell (2); Mollusca collected at Galapagos Islands, Hertlein & Strong; Kapiti Island mollusca, List of Shells, pp. 159-161. A. S. & A. Wilkinson.

Museums, Collections, and Expeditions.—The Titian R. Peale collection, R. T. Abbott; Askoy expedition to Panama, L. G. Hertlein & A. M. Strong; Molluses of the Cape Expedition Series, New Zealand, A. W. B. Powell (3); Arcidae in the Allan Hancock Foundation collection, figs., H. Rost; Cephalopoda of the Harvard and Havana expedition off Cuba, G. L. Voss; The Banks Collection, G. L. Wilkins; Carlos Prieto Collection in the Museo Mexicano de Malacología, C. Pieltain; Types in the Senkenberg Museum of Cyclophoridae, A. Zilch (1):

DISTRIBUTION

(A) Geographical

(a) FOSSIL AND LAND AND FRESHWATER

PALEARCTIC REGION

Scandinavia.—Lower Ordovician stratigraphy in Sweden, Jaanusson & Mutvei; Land molluses of Värmland, Sweden, G. Lundgren.

Finland.—Clausilia dubia (Drap.) from Finland, E. J. Valovirta.

British Isles.—English Bathonian Ammonites, W. J. Arkell (3); A Myalina fauna from the Namurian of north Ayrshire, and its affinities with Naiadites, G. M. Bennison; S.E. England Jurassic a new genus Myrene, R. Casey (1); Isle of Wight. Acanthinula in Holocene at Gotland, A. G. Davis; Isle of Wight and Truntatellina cylindrica britannica, Pils., A. G. Davis; Dorset occurrence records of Truncatellina cylindrica britannica, A. G. Davis; Hertfordshire mollusca, [A. E. Ellis]; Carboniferous Goniatites from Devon, England, R. Goldring; Surrey and Planorbis laevis, M. P. Kerney; Hertfordshire and former occurrence of Vertigo parcedentata (Braun), M. P. Kerney (1); List of Norton Common, Herts, tufa species, M. P. Kerney (1); List of fossils (pp. 110-112), from the Silurian May Hill Inlier, Gloucestershire, J. D. Lawson; British Isles and range of Planorbarius corneus, H. F. McMillan: Westmorland and occurrence of Lauria anglica, G. W. Pitchford; A closed pond in Derbyshire and Dreissena polymorpha, G. W. Pitchford (2); Malham, Yorkshire and Clausilia dubia Drap., L. W. Stratton; British Lower Cretaceous Belemnites, H. H. Swinnerton (1); British Carboniferous Naiadites, A. Trueman & Weir: The mollusca of the Clacton Channel deposits, Essex, S. H. Warren.

France.—Arion lusitanicus in France. C. v. R. Altena & others; Pyrenees Radilites and Statigraphical locations, G. Astre; Synchronisation between the Palaeccene of Gand-Ostende and Ilede-France, L. Feugueur; Oligocene molluscs from Jura and the Haute-Savoie, P. Jodot (1); Ferruginous Oolite of the Callovo-Oxfordien of Chaumont to Châtillon-sur-Seine, France, P. L. Maubeuge (1).

Netherlands.—A deposit of tropical cowries, W. S. S. v. B. Jutting (1).

Belgium.—Occurrence of Succinea arenaria, A. Anteunis; New Oligocene molluses from Belgium, Glibert & Heinzelin; Miocene molluses from Belgium, Glibert & Heinzelin (1).

Switzerland.—Macrocephalites from the Jurassic of Switzerland, A. Jeannet; Jurassic ammonites from the Swiss Jura, P. L. Maubeuge; Molluse fauna of Basle, Switzerland, F. Woltersdorf.

Portugal.—Portuguese mollusca, J. O. de Castro; Miocene Lamellibranchs from Portugal, O. da V. Ferreira (1); Miocene molluscs from Portugal, G. Zbyszewski.

Spain.—Oyster beds at Galicia, B. Andrau; Miocene molluses from Spain, P. M. Crusatont; Miocene gasteropods fron Granada, Spain, P. Jodot; Cretaceous molluses from Spain, Rios & Almela; Spanish Quaternary fossils, N. Solé.

Italy.—Myophoria from the Trias of the Brembo Valley, Italy, B. Accordi; Pliocene fauna of Strongoli, Italy, R. Malaroda; Eocene fossils from Udine, Italy, B. Martinis; Triassic molluscs from Italy, A. Pollini; On G. Brocchi's "Conchiologica fossile subapennina", C. R. Ronchetti (3); Distribution of land molluscs in Southern Italy and North Africa, C. Sacchi (2); Sorrento faunal researches, C. F. Sacchi (3); Terrestrial and littoral molluscs of the Italian Adriatic, C. F. Sacchi (7); Sorrento, Helicella claudia a new species, C. F. Sacchi (9); Pliocene fauna of Savona, Italy, G. Tavani; Oligocene molluscs from Livorno, Italy, G. Tavani (1); Quaternary fauna of Livorno, Italy, G. Tavani (2).

Austria.—New Cretaceous ammonite from Austria, B. Plöchinger; New molluscs from the Tertiary of Austria, F. Sauerzopf; Molluscs from the Austrian Tertiary, F. Sauerzopf (2); Traces of boring sponges in Actaeonella from the Austrian Jurassic, F. Schremmer; Molluscan fauna of Arlberge, Austria, F. Starmühlner (1); Tertiary Teredinidae from Austria, A. F. Tauber.

Hungary.—Molluscs of Szeged, Hungary, A. Horváth; Molluscs of Theiss, Hungary, A. Horváth (1); Eocene molluscs from Hungary, M. Kiss-Kocsisné Bányai; Pliocene and Pleisto-

cene of Komárváros, Hungary, D. I. Molnárné; Hungary, on the Daudebardiidae, L. Soós; Middle Miocene fauna of Várpalota, Hungary, L. Strausz.

Poland.—Comparison of Radix species from Poland, M. Jackiewicz; New species of Conularia from the Carboniferous of Poland, S. Kwiatkowski; Interglacial molluses from Warsaw, J. Urbański.

Greece.—The Neogene of Greece.

S. Charalambakis; Pliocene fauna of Greece, B. v. Freyberg; Miocene molluscs from Greece, O. Kühn (1); The Neogene of Kythira, Greece, N. Manolessos; Tertiary fauna of Kythera, Greece, A. Papp; Fossil fresh-water gastropods from Greece, A. Papp (1); Pliocene molluscs from Elis, Greece, A. Papp (2); New Melanopsis from the Greek Pliocene, A. Papp (3); Fresh-water gastropods from the Pliocene of Greece, Papp & Psarianos; The Pliocene of Achaia, Greece, P. S. Psarianos; Hydrogeology of the Tertiary of Arta, Greece, P. Psarianos (1); New species of vivipara from the Neogene of Achaia, Greece, P. Psarianos (2); The Neogene of Peloponnes, Greece, P. Psarianos (3).

Europe.—Cretaceous ammonites near Belgrade, Yugoslavia, M. Z. Andjelković; Triassic cephalopods of Slovakia, D. Andrusov & Kováčik; Pliocene molluses of Hungary, F. Bartha & L. Soos; The Jurassic of the Central Carpathians, K. Birkenmajer & J. Znosko; The Cretaceous of the eastern Carpathians, M. Filipescu; Fossils of the Lias of Schwäbischen Jura, G. Einsele; The Miocene between Bodensee and Salzach, Gauss & Schmidt-Thomi; The Lias of Fay (Ain) Jura, S. Guérin (1); Mollusc fauna of Grundewald, Berlin, Hansen; Non-marine fauna in the Carboniferous stratigraphy of western Europe, V. S. van der Heide; Two gastropods from south eastern Moravia, V. Hudec; Molluses from the Balkan peninsula, S. Jaeckel (1); Glauconia coquandi in the Cretaceous of eastern Europe, M. Pasić; Mollusc fauna of the Harze mountains, Pfitzner & Jaeckel; Tertiary Planorbiidae of the eastern European alps, F. Sauerzopf (1); Pelecypodichnus, new genus from Jurassic of central Europe, A. Seilacher: New species of *Pholadomya* (*Procardia*) from the Callovian of Europe [Germany],

A. Zeiss; German Tertiary molluses, A. Bernhauser; Germany miocene, F. Braudtner; Fossil molluses of Auerbach, Germany, G. Brunner; Molluses from Sauerland, Germany, H. Dittmar; Devonian molluses of Germany, A. Fuhrmann; Concretions from the German Oligocene, K .- W. Geib; Recent and Holocene mollusea from Hanover, Germany, H.-O. Grahle; Molluses of the Lias between Thoirette and Turgon, Jura, S. Guérin (2); Molluses of Obernburger, Germany, L. Hasslein; Polamopyrqus jenkinsi in Berlin and Brandenburg, H. Herold; Molluses of the Miocene of Holstein, Germany, W. Hinsch; The ammonite Taramelliceras in the Jurassic of south west Germany, H. Hölder; Ferrissia North American fresh-water snail found in Germany, S. Jaeckel (2); Molluse fauna of Thüringen S. Jaeckel (3): Molluses of the central German mountains, S. Jaeckel Limapontia depressa pellucida Kevan, Jaeckel (5); Monacha cartusiana in Germany, H. Janus; Cuttlefish impressions from the Solnhofener Plattenkalk [Jurassic], F. Klinghardt; Two abnormal examples of Poltopleuroceras from Germany, H. Kolb; Piloceras in the upper Lias [Germany], O. Kuhn (2); The Triassic in Germany, A. H. Müller; Oncophora in the Miocene of Germany, A. Papp (5); Ceratites from the German Triassic, H. Penndorf; Trichia sericea in Brandenburg, I. Pfitzner; Molluscs of eastern Germany, H. Plate; German Mesozoic cephalopods, G. Rosenberg; Maastricht of Henmoor, Germany, F. Schmid; Land snails of Mecklenburg, Germany, H. Schmidt (1); Vertigo moulinsiana in Brandenburg, T. Schmierer (1); Jurassic molluses from Germany, F. Schnittmann; Vertigo ronnebyensis in Berlin, F. Zimmerman; Vertigo ronnebyensis in Berlin, F. Zimmerman (1): Molluses of the Aquitanien of Bavaria, H. K. Zöbelein.

Bohemia.—The Cretaceous of Hronov, Bohemia, J. Dvořák; New Silurian gastropods from central Bohemia, R. Horný; New Devonian gastropod family from Bohemia, R. Horný (1); Pseudoanodonta from Bohemia, J. Petrbok; Devonian Goniatites from Bohemia, F. Prantl; New Devonian pelecypod from Bohemia, Prantl & Růžička; New Devonian pelecypod

from Bohemia, **Prantl & Růžička** (2); A Silurian lamellibranch from Bohemia, B. Růžička (1).

Czechoslovakia.-Systematic review of Czechoslovakian aquatic molluses, J. Brabenec; Molluscs of Czechoslovakia J. Brabenec (1); Miocene fauna at Rapovee, Hano & Senes; Czechoslo-vakian molluses, V. F. Hlaváć; New form of Viviparus from Czechoslovakia, O. Kapler; Mollusea from Czechoslovakia, V. Ložek (1); Quaternary molluses of Czechoslovakia, V. Ložek (2); Geographical distribution of Laciniaria moravica Brabenec in Europe, Ložek & Mácha; Molluses from Czechoslovakia, S. Mácha; Molluses of Michálkovice, Ostrau, Czechoslovakia, S. Mácha (1); Jurassie ammonites of eastern Serbia, M. Mihailović; Fresh-water fauna of Liptovské hole, Czechoslovakia, S. Obr; Miocene fauna of Czechoslovakia, V. Pokorný; Miocene fauna of southern Czechoslovakia, J. Seneš; Devonian fauna of Czechoslovakia, J. Skácel; Clithon pictus from the Miocene of Czechoslovakia, J. Svagrovský (2).

Roumania.—List of molluses from the Roumanian Cretaceous, R. Givulescu; List of fossils from the Rumanian Lias, G. Răileanu; Pulmonate gastropods from Roumania, A. Grossu; Jurassic and Cretaceous fossils, D. Jacob; Anthracosia from the Roumanian Permian, G. Răileanu (1).

Yugoslavia.—Cretaceous of Serbia, Yugoslavia, M. Z. Andelkovic; Triassic cephalopods from the Austrian-Yugoslavian alps, O. Kühn; New gastropods from Yugoslavia, P. Radoman; New gastropod from Lake Ohrid, Yugoslavia, Stanković & Radoman; Gastropods of Lake Ohrid, Yugoslavia, L. Snagarova; Miocene fauna of Yugoslavia, P. Stevanović & Pantic; The Lias of eastern Yugoslavia, D. Veselinović.

Estonia.—Pupillidae from Estonia, C. Krausp; The Clausillidae of Estonia, C. Krausp (1); Valloniidae of Estonia, C. Krausp (2).

Russia.—Cretaceous deposits at the mouth of the Dnieper-Don rivers, U.S.S.R., G. I. Bushinskii: Evolution and systematics of Cretaceous Belemnitella, J. A. Jeletzky (1); The Ordovician of Kazakstan [U.S.S.R.?], B. M. Keller; Chlamys from the Tortonian of Russia, L. N. Kugrin; Ancycloceras from the

Maastrichtian of Russia, S. I. Pasternak; Permian Nautiloidea from the southern Ural, Ruzhentzev & Shimanskii; from the Sakmarskian and Artinskian of the southern Ural, V. N. Shimanskii

China.—New Gigantomilax from China, C. v. R. Altena (1); Marine molluses of North China, Chang Hsi, C-yen Chi & C-m. Li.

Syria, —Jurassic ammonites from Syria, O. Haas (1); Cretaceous & Tertiary of Israel with regard to Pecten, Z. Reiss; Cercaria from Melanopsis in Israel, H. Ullman.

Asia.—Jurassic fossils of the Yémen, E. Basse (1); Mollusca of S.-W. Asia, K. Büttner; Gastropods of the "Grès de Taouilah" Cretaceous, Yémen, J.-P. Lefranc; Neposiella from the Cretaceous of Gebel Manzour (Moghara-Sinai-Egypt)., I. G. E.-d. Mahmoud.

Mediterranean Islands.—Pliocene and Pleistocene fauna of Sicily, Malatesta & Nicosia; Hydrobiology of Majorca, R. Margalef; Miocene fauna of Sicily, M. L. Nicosia; Marine molluses from Sicily, O. Priolo; Majorca Neogenc Pectinidae, J. B. Rullán; New genus from the Hauterivian of Majorca, S. S. Sarkar.

North Africa.-Molluses in the collection of the Oceanographic Station of Salammbo, Tunisia, M. Arnould; Neocretaceous lamellibranchs from Libya, C. Baroni & L. Incitti; New Cretaceous ammonite from Tunisia, E. Basse; Quaternary molluses from Casablanea, P. Biberson; Revision of the Neocretaceous molluses of Libya, C. Bricci; The Neocomian fauna and Ammonites in North Africa, M. Dalloni: The Miocene fauna of Angola, E. Dartevelle & J. Rogers; New Miocene gastropods from Algeria, L. David & P. Jodot; On the stratigraphical position of lower Cretaceous Ammonites and a new genus. [Algeria], M. D. Delga; Bacillus sp. as a control for snail hosts of Schistosoma in Egypt, E. Dias & M. Dawood; Liassic molluses from Djebel Beut-Saidane et Dj. Zaghouan [Tunisia], G. Dubar; The presence of Agria marticensis and Praeradiolites pulchellus in the Cretaceous of Algeria, J. Emberger (1); Ammonites in relation to Jurassic formations of the Saharan Atlas (Algeria), J. Emberger (2); Tisoa from the Numidien of Tunisia, C. Gottis; Cyrenaica Eocene Aturia, O. Haas; Cretaceous molluses from Tunisia, A. Lapparent; Cretaceous cephalopods from Moghara, Egypt, I. G. E. Mahmoud (1); New species of Nucula from the Portlandian of Liban [Africa]. D. Mongin; Stone and wood-boring molluses from Africa, G. Roch (2); Revision of Neocretaceous Ostreidae from Libya. C. R. Ronchetti; Euparupha (Helicidae) from Morocco, C. Sacchi (6).

ORIENTAL REGION AND EAST INDIES

India & Pakistan.—Limnoca spp. as intermediate host of Schistosoma turkestanicum in Pakistan, M. Abdussalam; Recovery of a community of Placenta placenta in the Gulf of Cutch, India, S. Gokhale; New Loricata from Manora Island, Karachi, Pakistan, P. Kaas; Macrochlamys a snail pest from the Punjab, Latif Qadir; Helicella and Macrochlamys as intermediate hosts of sheep lungworm in the Punjab, M. Sarwar (1).

Malaysia.—Malayan Streptaxidae: Discartemon and Oophana, W. S. S. v. B. Jutting; Freshwater molluses from Sumba, Indonesia, W. S. S. v. Benthem Jutting (2); Amphidromus from Islands off the coast of Sumatra, F. F. Laidlaw.

Japanese Subregion.-New species of Pleuronectites in the Triassic of Japan, M. Amano; Miocene molluses from Fukushima Prefecture, Japan, S. Aoki; A Japanese Myadora from the Pleistocene, T. Habe (2); Pelecypoda and Scaphopoda from Akkeshi Bay, Japan, T. Habe (5); Notes on three Japanese Tellinid species described by V. Bertin, T. Habe (7); A deposit of shell remains in Ofunato Bay, T. Habe (8); Koninckioceras from the Japanese Permian, I. Hayasaka; A new gastropod from the Permian of Japan, I. Hayasaka (2); New Permian Foordiceras from Japan, Hayasaka & Ozaki; Miocene Perotrochus in Japan, Hirayama; Miocene Lucinoma species from Japan, K. Hirayama (1): New Miocene molluses from Tochigi Prefecture, Japan, K. Hirayama (2); Tertiary molluses from Fukushima Prefecture, Japan, K. Hirayama (3); Perotrochus from the Japanese Miocene, K. Hirayama (4); Neo-Triassic pelecypods from Tokyo Prefecture, Japan, K. Ichikawa; A new subspecies of Eumorphotis from the Shionosawa limestone.

Japan, K. Ichikawa; Non-marine mollusca from the Tertiary of Japan, Y. Kamada; New Miocene molluscs from Fukushima Prefecture, Japan, S. Kanno; Faunal analysis of molluses from raised beach deposits of Kamakura, Japan, S. Kanno; Tertiary mollusca from Nagasaki Prefecture, Japan, S. Kanno (1); Miocene Dosinia from Japan, S. Kanno (3); Argonautinae from the Neogene of Japan, Y. Kaseno; Coloured illustrations of Japanese shells, T. Kira; New genus of the Argonautinae from the Miocene of Japan, T. Kobayashi (1); New species of Aturia from the Japanese Miocene, T. Kobayashi (2): Presence of Aturia in the Miocene of Etchu and Iwami Provinces, Japan, Kobayashi & Mastani; Jurassic Trigonians from Japan, Kobayashi & Mori; Jurassic Trigonians in Japan: Vaugoniinae, Kobayashi & Mori (1); Japanese Jurassic Trigonians, Part IV, Kobayashi & Tamura; Miocene mollusca from Noto Peninsula, Japan, K. Masuda; Cretaceous bituberculate Pachydiscids from Hokkaido and Saghalien, Japan, T. Matsumoto; The Kossmaticeratidae from the Cretaceous of Japan, T. Matsumoto (1); Cretaceous Desmoceratids from Hokkaido and Saghalien, Japan, Matsumoto & Obata: New Miocene Acila species from Japan, Nagao & Huzioka; New Triassic Bakevellia spp. from Japan, K. Naka-zawa; New species of Glyptophiceras from the Mesozoic of Hyogo Prefecture, Japan, Nakazawa & Shimizu; Conchocele compacta from the Tertiary of Japan, M. Omori; New Tertiary species of Propeamussium from Japan, M. ômori (1); Pliocene molluscs from Tyôsi City, Japan, H. Ozaki; Tertiary molluses from Gunma Prefecture, Japan, Ozaki & Shikama; Molluses from the Tokara Islands, Japan, Oyama, Yamamoto & Tohioka; Triassic ammonites from the Kwanto Massif, Japan, S. Sakagami; Hammatoceras from the Japanese Jurassic, T. Sato; Jurassic Ammonites from north central Japan, T. Sato (1); New species of Brechites from the Japanese Pliocene, T. Shikama; Tertiary molluses from South Nagano Prefecture, Japan, T. Shikama (1); Amussiopecten from the Tertiary of Miyazaki Prefecture, Japan, T. Shuto; New species of Titiscania from Hachijō-jima island, Japan, I. Taki (2); Palaeogene fossil molluscs

from Hokkaido, Japan, S. Uozumi; New Jurassic ammonite from Iwaki Province, Japan, H. Yabe; New Japanese slugs, N. Yamaguchi; Slugs of Japan, Yamaguchi & Habe.

AUSTRALIAN AND POLYNESIAN REGION

Australia.—Parorchis acanthus australis cercaria parasitic in snails, in S. Australia, L. M. Angel; Mesozoic stratigraphy of the Canning Desert and Fitzroy Valley, Western Australia, R. O. Brunnschweiler; Mollusca of the raised beaches of South Australia, R. L. Crocker & B. C. Cotton: Australian Nuculanidae Tertiary and recent, R. K. Dell (6); Tertiary fossils from Melbourne, Australia, Gill & Baker; New Oligocene Aturia from South Australia, M. F. Glaessner; New Cretaceous and Tertiary nautiloids from Western Australia, Glenister, Miller & Furnish; Snails from Betsy Island, Tasmania, A. M. Hewer; Systematic list of Tasmanian molluscs, R. C. Kershaw; Literature relative to molluscs from Victoria, Australia, R. C. Kershaw (1); Studies on Australian Charopidae, R. C. Kershaw (2); Studies on Australian Charopidae, R. C. Kershaw (3); Pliocene molluscs of Adelaide Plains, Australia, N. H. Ludbrook; Pliocene mollusca from Adelaide Plains, Australia, N. H. Ludbrook (1); New gastropods from N.S.W., Australia, C. McLauchlan; New gastropods from Australia, C. McLauchlan (1); Cassididae from N.S.W. and Queensland, Australia, G. Thornley (1): 2 new molluses from N.S.W., Australia, L. Woolacott.

New Zealand.—Land molluses of New Zealand, A. Brookes; Amphidesma ventricosum from New Zealand, R. M. Cassie; Molluscs of Pyramid Valley swamp, Canterbury, New Zealand, E. S. Deevey; Molluscs from deep-water rocks in New Zealand, R. K. Dell (1); Land molluscs of Fiordland, New Zealand, R. K. Dell (5); Tertiary and recent Nuculanidae from New Zealand, R. K. Dell (6); New Zealand subfossil land molluses and origin of Paryphanta, R. K. Dell (7); Freshwater molluses of New Zealand, P. Dickinson; Molluses from Pyramid Valley Swamp, Canterbury, New Zealand, J. R. Eyles; Pliocene molluscs from Campbell Island, New Zealand, C. A. Fleming (1); Fossil

mollusc collections in the New Zealand geological survey, C. A. Fleming (2); The Suter Collection of New Zealand Mollusca, C. A. Fleming (3); Miocene molluscs from New Zealand, C. A. Fleming (4); Paryphanta watti an extension of range in New Zealand, N. Gardner; Serpho matthewsi in New Zealand, N. W. Gardner (2); The genus Scrobs in New Zealand, A. Hipkins (1); Mollusca of the southern islands of New Zealand, A. Powell (3); Recent fossil molluscs from Wellington harbour, New Zealand, R. Roberts; Molluscs of Stewart Island, New Zealand, E. K. Smith; Molluses of Stewart Island, New Zealand, E. K. Smith (1); Notoplax cuneata in New Zealand, W. Thomson; Austrosuccinea archeyi from Mount Maunganui, New Zealand, W. Thomson (1); Pliocene molluses from Resolution Island, New Zealand, Wellman, Reed & Fleming; New Zealand, mollusca of Kapiti Island, A. S. & A. Wilkinson; Introduced land molluses of New Zealand, H. Witten.

Pacific Islands.—Stratigraphy and palaentology of central New Caledonia, J. Avias; Pulau Panaitau Mollusca, Butot (2); Solomon Islands on new genus of the Synceridae, W. S. Clench (1); Raiatea, Society Islands new Partula, H. E. Crampton; Terrestrial molluses from Nissau Island, Solomon Islands, R. K. Dell (2); New Lamellidea sp. from Solomon Islands, R. K. Dell (4); New Philippine species of Amerianna, B. Hubendick (1); New Caledonian Placostylus notes on some species, figs., T. Pain; List of molluses from the Falkland Islands (Islas Malvinas), R. Ringuelet.

ETHIOPIAN REGION

East Africa.—East African Vitrinae, B. Hubendick (4).

West Africa.—Veronicellidae from Fernando Po Island, West Africa, L. Forcart (1); Homeomorphy in Nigerian Cretaceous ammonites, R. A. Reyment

Central Africa.—Tanganyika Caelatura aegyptiaca calathus Brgt, J. Bouillon; Freshwater molluses of Uganda, G. Mandahl-Barth; Syrnolopsis and Aceya anatomy of Tanganyikan forms, Mandahl-Barth (1); New Lanistes species from Africa, T. Pain; New slugs from the Gughé Highlands, Southern Ethiopia, H. Quick; Planorbidae, nom-

enclature of African forms, **J. Schwetz**; Molluses as carriers of bilharziosis in Africa, **J. Schwetz** (2).

South Africa.—South African shore life, K. H. Barnard; Molluscs from Mozambique, J. M. Braga; New Cretaceous ammonites from Zululand, E. v. Hoepen; New Cretaceous ammonites from Zululand, E. van Hoepen (1); New Cretaceous ammonites from Zululand, E. van Hoepen (2).

Indian Ocean Islands.—First record of Zonitoides arboreus from Madagascar, R. P[aulian].

Atlantic Islands.—Azores: aspects of fauna and flora, G. Chapman; Jamaican Rudist types of Whitfield from the Cretaceous, L. J. Chubb; The Cretaceous Succession in Jamaica, L. J. Chubb (1); A new Murcx from Cuba, W. J. Clench; Hitherto unpublished note by Dr. C. de la Torre on Polymitas from Cuba, J. A. Conde; Miocene Pectinidae from Santa Maria (Azores), O. da V. Ferreira.

NEARCTIC REGION

Iceland.—Fossils of the Klimage-schichte Islands, M. Schwarzbach.

Alaska.—Jurassic molluses from Alaska, R. W. Imlay; Alaska, distribution and ecology of marine invertebrates, G. E. MacGinitie.

Canada.—Viviparus viviparus L. in eastern Canada, E. L. Bousfield; Ordovician molluses from British Columbia, T. Kobayashi (3); Canada Cape Breton, Land and Freshwater Gastropoda, G. K. MacMillan; Ontario, Shells collected in Loughborough Lake, W. A. Swanker; New Cenomanian ammonites from Alberta, Canada, Warren & Stelck.

North America.—The Carboniferous fossil Tylonautilus in America, Miller & Furnish (2); Fossil molluses from N. America, D. Taylor; West North American Pleistocene Upwelling and thermally anomalous faunas, J. W. Valentine.

United States as a whole.—Organic matter in relation to distribution of pelecypods, R. E. Bader; New Chazyan Orthocones from the Champlain Basin, of North America, R. H. Flower (2); Trails and tentacular impressions of cephalopods in the Corryville of Cincinnati, America, R. H. Flower (3).

North Eastern United States.—Shells of Fire Island, near Long Island, U.S.A., C. Ahders; Virginian Land snails and their relation to ecological soil factors, J. B. Burch; Virginia, Land snails of Hanover, Henrico and Chesterfield Counties, J. B. Burch (1); New species of Endolobus from Carboniferous of Kentucky, C. Collinson; Mississippian Goniatites from Illinois and adjacent states, C. W. Collinson (1); Minnesota land snails, G. Dawley; Ohio, confirmation of occurrence of Pallifera fosteri F. C. Baker into north-eastern districts, H. W. Dexter; A new Emarginula from the Cretaceous New Jersey, C. J. Eichman; Palaeontology of the Ordovician of Missouri, R. Heller; Iowa, occurrence of Discus macclintocki in Bixby State Park, Hubricht (6); Carboniferous cephalopods from Michigan, Miller & Garner; Illinois. Ecological aspects of Unionid fauna of Lake Springfield, P. W. Parmalee; New Ordovician molluscs from Minnesota, L. Powell; Land and fresh-water molluscs of Great Neck, Long Island, New York, E. Slater; Cretaceous molluscs of New Jersey, L. W. Stephenson; Cretaceous molluscs from Missouri, L. W. Stephenson (2); Paleoecology of the Maquoketa coquina beds at Graf. Iowa, P. Tasch (1); Miocene pearls from the Atlantic Coastal Plain, Baltimore, H. Vokes.

South Eastern United States .- Pliocene Pyramidellidae from Florida, P. Bartsch; Palaeoecology of the Weches formation (Eocene) of Texas, N. M. Curtis; Florida, Oysters of the Crystal River, C. E. Dawson; Oklahoma additional localities for Land mollusca, D. G. Dundee; New Conularia from the Permian of West Texas, North America, R. M. Finks; Tylonautilus from the Carboniferous of Oklahoma. Miller & Furnish (1); New name for Hermania Puri, from the Miocene of Florida, H. S. Puri; Mollusca of the Mississippi Delta, H. G. Rickards (1); Carboniferous paleoecology in Texas, R. E. Sloan; Mollusca of the Cenomanian of Texas, L. W. Stephenson (1); Mollusca of Lake Texoma (Denison Reservoir) Oklahoma and Texas, J. E. Sublette: Georgia: The mollusca of Warm Springs, M. C. Teskey (1).

Western United States.—Cretaceous gasteropoda from Baja California,

Mexico, E. C. Allison; Albany County, Wyoming, 28 Land and 17 Freshwater molluses found, D. E. Beetle: A new species of Monadenia from Tuolumne Co., California, figs., G. S. Berry; A new species of Discus from the Mission Range, Montana, figs., S. S. Berry (2); Possible nautiloid mandibles from the Permian of Arizona, L. F. Brady; Protocanites (Goniatite) in the Mississippian of north-eastern Nevada, W. M. Furnish (1); Common sea shells of Los Angeles County coast, Hill & Tompkins; Stratigraphical implications of Belemnitella praecursor in Kansas, J. A. Jeletzky (2); Triassic fauna of Idaho, U.S.A., B. Kummel; Intertidal edible molluses of Oregon, J. McCauley; Western Montana and Magnipelta mycophaga Pils., fig., Pilsbry & Brunson; South Dakota new records, E. S. Roscoe (1); Thayer area, Neosho Co., Kansas a type locality of Triodopsis cragini, Call., E. J. & L. D. Roscoe; Utah, Gastropod fauna of the Brighton Area, Wasatch Mountains, E. J. & L. D. Roscoe; Ordovician cephalopods from central Colorado, W. C. Sweet (1); Palaeozoic cephalopods from Wyoming and Utah, W. C. Sweet (2).

CENTRAL AMERICAN REGION

Mexico.—Didymotis from the Cretaceous, R. W. Imlay (1); Lymnaea obrussa as an intermediate host of Fasciola hepatica, in Mexico, L. Mazzotti; New Durania spp. from the Mexican Cretaceous, F. K. Mullerried; New Helicidae, A. Solem.

West Indies.—List of Cuban molluses, C. Aguayo; Trinidad, Didymotis from the Cretaceous, R. W. Imlay (1); List of the Cuban species of Liguus M. Jaume; New species of Liguus from Cuba, M. Jaume (1); Aturia peruviana from the Tertiary of Costa Rica, Miller, Furnish & Paguaga; Cuba, new subspecies, of Polymita, C. de la Torre; Prolific increase of Ampularia introduced to the Caguitas river, Puerto Rico, W.I., F. Watlington.

SOUTH AMERICAN REGION

Colombia.—Didymotis from the Cretaceous, R. W. Imlay (1).

Venezuela.—Pupillidae from northern Venezuela, S. Arias; Didymotis from the Cretaceous, R. W. Imlay (1).

Ecuador.—Subfossil molluses of Pen. de Santa Elena, R. Hoffstetter; Subfossil molluses from Pen. Santa Elena, R. Hoffstetter (1). Subfossil molluses from the Pen. de Santa Elena, R. Hoffstetter (2).

Brazil.—Tertiary fauna of Paraná, Brazil, K. Beurlen; Land mollusca, F. Haas (5); Limnology of Rio Grande do Sul, H. Kleerekoper; Additions and corrections to the Catalogue of Brazilian molluses, F. L. de Morretes; Molluscan Schistosoma carriers in Brazil, H. Sioli.

Peru.—New land mollusca, F. Haas (5); Lake Titicaca gastropoda, F. Haas (6).

Chile.—Aturia from Southern Chile, Miller & Furnish.

Argentina.—Eurydesma from the Permian of eastern Argentina, H. J. Harrington; Tertiary (?) Strophocheilus, J. Parodiz(2); New speciesof Plectostylus, J. Parodiz(4); Cambrian and Ordovician molluscs from Mendoza, C. Rusconi; New molluscs from the Cerro Colorado, M. Scott; New molluscs from northwestern Argentina, M. Scott (1); New species of Nenia Clausiliidae, M. I. Scott (2); New species of Epiphragmophora M. I. H. Scott (3); Limnaeidae, M. I. H. Scott (4); New species of Bostryx, M. I. H. Scott (5); New species of Hemisinus, M. I. Scott (6).

Patagonia.—New Eccene gastropod, J. J. Parodiz.

(β) Marine ARCTIC

Arctic Seas.—Arctic pelceypod fauna analysis, D. Nicol; New and known Trochidae from the Northern and Far-Eastern seas of Russia, Galkin.

NORTH TEMPERATE

Eastern North Atlantic.—Aplacophora and Placophora from the Baltic and the North Sea, S. Jaeckel; New Opisthobranchiata from France, E. & E. Marcus (1); Sweden S.-W. coast, A new Lepidomenea, M. Schwabl; Mollusca of Stockholm and vicinity, H. W. Waldin; Gasteropod larvae in north-western Norway, p. 197, K. Wilborg.

Gulf of Mexico.—Summary of information on Oysters, P. A. Butler; Effect on invertebrate fauna of salinity changes in bays of central Texas, B. H.

Parker; Marine shells of the west coast of Florida, Perry & Schwengel; A new Florida cone, figs., H. A. Pilsbry; Zoological notes and lists of species of the Gulf of Mexico, H. A. Rehder; Oyster industry of the United States Gulf Coast, G. Schlesselman; Gulf of Mexico—Cephalopoda, G. Voss (1); Gulf of Mexico—On the Oyster reefs, W. A. Price.

Mediterranean.—New Oncidiella from Cap Blanc, Tunisia, Gabe & Prenant; Sicily Faunules of Lakes Marsala, Ganzirri and Faro, P. Mars; Molluses from the northern Adriatic, Nümann & Beth; The Lamellibranchs of Padern (Aude) S. France, G. Petter-Receveur; The Eastern Mediterranean and distribution of Cephalopods therein, W. J. Rees; Molluses of Tremiti islands, Adriatic Sea and Pianosa island, off west Italy, C. Sacchi (5).

Eastern Pacific.—Californian occurrences of Ocythoe tuberculata, Raf., S. S. Berry (1); G. of Californian Phyllonotus, J. & C. Cate; California, occurrence of Alderia modesta Loven on the central coast, figs., Hand & Steinberg; New Lima from deep water off Central California, L. G. Hertlein; Mollusca from benthos samples from San Pedro Basin, California, N. J. Mattox (1); A new species of *Tretoniopsis* from Southern California, figs., N. T. Mattox (2); List of molluscs, collected from Alamitos Bay, California, p. 114, 115, Reish & Winter; West Coast of America Arcidae, figs., H. Rost; Review of Chitons of West Coast of N. America, A. G. Smith; Shells from South-east Farallon Island, California, A. G. Smith (1); Rare chiton from deep water off Central California, Smith & Hanna; Mytilidae of the West Coast of America, T. Soot-Ryen; A new species of Aplysia from S. California, figs., Winkler.

Western Pacific.—Japanese species of Conidae, Anon; Notarchus species from Japan, K. Baba; Eastern Pacific Muricacea. A systematic outline, J. Q. Burch; Ringiculidae and Retusidae in Japan, T. Habe; Hydatinidae, Bullidae and Akeridae in Japanese waters, T. Habe (1); Shells of the Japanese Islands and adjacent waters, S. Hirase; Ecology of Japanese oyster beds, Ito & Ismai; New Pleurotomaria from Japan, T. Kuroda; Volutidae of Japan, Kuroda & Habe; Perotrochus dredged near

the Izu Islands, Japan, H. Niino; New species of Carinaria from Japanese waters, T. Okutani; Chitons round Japanese Islands: Stenoplax, I. Taki; Chitons from the waters round Japanese islands, I. Taki (1); Atlantidae from Japanese waters, T. Tokioka (1); New species of Atlanta from Japanese waters, T. Tokioka (2).

TROPICAL AND SUBTROPICAL

East Atlantic.—New molluses from West African coastal waters, W. Adam; New species from the Ivory Coast, W. Africa, E. Binder; West Africa a new Trivia, J. Knudsen; Aplysia from the South African coast, W. Macnae.

Western Atlantic.—Littorina obtusata and its variation in the North Atlantic, J. Knudsen (1); Ostrea edulis in American waters, V. Loosanoff; A new Tropicorbis from N.-E. Brazil, D. T. de Lucena; New Floridan marine mollusca, T. L. McGinty; Occurrences of Spirula animal at Boynton Beach, Florida, T. L. McGinty (1); Marine shells of the west coast of Florida, Perry & Schwengel; Teredinidae from the East and West Indies, [G.] F. Roch (1); New Conus from Costa Rica, figs., J. S. Schwengel; Two new records of Strombus spp. from Jamaica, V. A. Zans.

Eastern Pacific.—New chiton from Ecuador, E. Leloup; West Mexico and Panama fauna, [A. M. Strong]; Mitridae and Volutidae of West Mexico and Panama, [A. M. Strong] (1); Naticidae of Western Mexico and Panama, [A. M. Strong] (2); Columbellidae of Western Mexico and Panama, [A. M. Strong] (3).

Indian Ocean.—Nassidae of Indo-China, M. Marche-Marchad; Cymatiidae and Bursidae from Indo-China, I. Marche-Marchad (1); Marine molluses of Agalega Isle, Chagos archipelago, Indian Ocean, H. Poisson; Marine Gastropoda of Bombay, Subrahmanyan, Karandiker & Murti; Cowries of the East Africa coasts, B. Verdcourt.

SOUTH TEMPERATE AND SUBANTARCTIC

Western Atlantic.—New *Thais* spp from Brazil, **F. L. de Morretes**; New Brazilian molluscs, **F. L. dé Morretes** (1); New marine gastropods from Argentine and Uruguay, A. Carcelles; South American species of Argobuccinum, A. Carcelles (1); Observations on Nucella, Chorus and Concholepas in South America, A. Carcelles (2); New species of Sunetta from Buenos Aires, M. D. Jurado; New Opisthobranchia from Brazil, E. Marcus.

Southern and Western Pacific.—A new Pecten subspecies from Byron Bay, N. S. Wales, C. A. Fleming; Placostylus gracilis from Viti Levu, Fiji, E. & N. Gardner; Revision of Pinctada spp. the Australian pearlshells, J. S. Hynd; New Cephalopods from Australia, T. Iredale (2); New albino Volutes from Australia, N. Jackson; Revision of the Turridae of New South Wales, C. F. Laseron; Revision of Liotiidae of New South Wales, Australia, C. Laseron; New South Wales Eulimoid shells, C. F. Laseron (2); New Triphoridae from Australia, C. F. Laseron (5); Patellidae and Acmaeidae in Australia, J. H. MacPherson; Tasmanian scallop fisheries, A. M. Olsen; Opisthobranchiate gastropods from Hawaii, J. M. Ostergaard.

Southern Ocean.—Studies on Notovola meridionalis from Tasmania, A. M. Olsen; List of additions to the recent molluses of New Zealand, A. Powell; Liarea from New Zealand, A. W. B. Powell (2); Adelaide gales and shells, K. Rutland; Iredalina aurantia from New Zealand coastal waters, S. Trigance (1); Cassididae in the Leigh district of New Zealand, T. Warren; Ellatriva memorata from New Zealand, K. Watene; Amphineura from Mt. Maunganui, New Zealand, G. Williams (2).

(B) Geological

Quaternary.—Sub-recent gastropod from Singapore, bored through by a predator, F. E. S. Alexander; Quaternary molluses from Casablanca, P. Biberson; List of Pleistocene and Holocene fossils from Belgium, J. de H. Braucourt (1); Mollusca of the raised beaches of South Australia, R. L. Crocker; Holocene Acanthinula lamellata in the Isle of Wight, A. G. Davis; New Zealand subfossil land molluses, and origin of Paryphanta, R. K. Dell (7); Recent and Holocene mollusca from Hanover, Germany, H.-O. Grahle;

Subfossil molluses of Pen. de Santa Elena, Ecuador, R. Hoffstetter; Subfossil molluses from Pen de Santa Elena, Ecuador, R. Hoffstetter (1); Faunal analysis of molluses from raised beach deposits of Kamakura, Japan, S. Kanno; Quaternary molluses of Czechoslovakia, V. Ložek (2); Quaternary fossils from Spain, N. Solé; Four Quaternary deposits in the Cambridge area and list of species, R. W. Sparks; Quaternary fauna of Livorno, Italy, G. Tavani (2); The mollusea of the Clacton Channel deposits, Essex, S. H. Warren.

Pleistocene.—A new Pleistocene Myodora from Japan, T. Habe (2); Pleistocene fauna of Sicily, Malatesta & Nicosia; Pleistocene of Komárváros, Hungary, D. I. Molnárné; Pleistocene mollusca from the submarine canyon Hudson River, Richards & Ruhle; New molluscs from North America, D. Taylor; Interglacial molluscs from Warsaw, J. Urbańoki; Pleistocene upwelling and thermally anomalous faunas of the Pacific Coast, J. W. Valentine

Tertiary.—Tertiary molluses of Germany, A. Bernhauser; Tertiary fauna of Paraná, Brazil, K. Beurlen; Distribution and variation of Cepaea, C. Boettger (3); Check list of Australasian Tertiary Nuculanidae, R. K. Dell (6); Tertiary littoral gastropods, P.-H. Fischer (5); Tertiary fossils from Melbourne, Australia, Gill & Baker; New Tertiary Nautiloids from Western Australia, Glenister Miller & Furnish; Tertiary molluscs from Fukushima Prefecture, Japan, K. Hirayama (3); Tertiary mollusca from Nagasaki Prefecture, Japan, S. Kanno (2); Aturia peruviana from the Tertiary of Costa Rica, A. K. Miller Furnish & Paguaga; Conchocele compacta from the Tertiary of Japan, M. Omori; New Tertiary species of Propeamussium from Japan, M. Ômori (1); Tertiary molluscs in a limestone boulder from Gunma Prefecture, Japan, Ozaki & Shikama; Tertiary fauna of Kythera, Greece, A. Popp; Tertiary (?) Strophocheilus from Argentina, J. Parodiz (2); Hydrogeology of the Tertiary of Arta, Greece, P. Psarianos (1); Early Tertiary in Israel, Z. Reiss; Molluscs from the Tertiary of Austria, F. Sauerzopf (2); Tertiary molluses from South Nagano Prefecture, Japan, T. Shikama (1); Amussiopecten from the

Tertiary of Miyazaki Prefecture, Japan, T. Shuto; Tertiary Teredinidae from Austria, A. F. Tauber; Fossil molluses from the Palaeogene of Hokkaido, Japan, S. Uozumi; Tertiary and Recent Solemyacidae, H. E. Vokes (1).

Pliocene,—Pliocene molluscs of Hungary, F. Bartha & L. Soos; Pliocene Pyramidellidae (Gastropoda) from Florida, P. Bartsch; The Neogene of Greece, S. Charalambakis; Pliocene molluses from Campbell Island, New Zealand, C. A. Fleming (1); Pliocene fauna of Greece, B. v. Freyberg: Argonautinae from the Neogene of Japan, Y. Kaseno: Pliocene molluses of Adelaide Plains, Australia, N. H. Ludbrook; Mollusca from the Pliocene of Adelaide Plains, Australia, N. H. Ludbrook (1); The Pliocene fauna of Strongoli, Italy, R. Malaroda; Pliocene fauna of Sicily, Malatesta & Nicosia; The Neogene of Kythira, Greece, N. Manolessos; Pliocene of Komárváros, Hungary, D. I. Molnárné; Pliocene molluscs from the Na-arai Conglomerate, Tyôsi City, Japan, H. Ozaki; Pliocene molluses from Elis, Greece, A. Papp (2); New Melanopsis from the Greek Pliocene, A. Papp (3); Freshwater gastropods from the Pliocene of Greece, Papp & Psarianos: The Pliocene of Achaia, Greece, P. S. Psarianos: New species of Vivipara from the Neogene of Achaia, Greece, P. Psarianos (2); The Neogene of Peloponnes, Greece, P. Psarianos (3); On A. Brocchi's "Conchiologica fossile subapennina," C. R. Ronchetti (3); Neogene Pectinidae of Majorca, J. B. Rullán; Planorbiidae from the Pannon of eastern Europe, F. Sauerzopf (1); Neogene macrofauna of east Czechoslovakia, J. Senes (2); New species of Brechites from the Japanese Pliocene, T. Shikama; Neogene fauna of Czechoslovakia, J. Švagrovský; Pliocene fauna of Savona, Italy, G. Tavani; New molluse from North America, D. Taylor; Pliocene molluses from Resolution Island, New Zealand, Wellman, Reed & Fleming

Miocene.—Miocene molluses from Fukushima Prefecture, Japan, S. Aoki; Miocene mollusea from Germany, F. Brandtner; Miocene molluses from Spain P. M. Crusafont; The Miocene fauna of Angola, E. Dartevelle & J. Rogers; New Miocene gastropods from Algeria, L. David & P. Jodot; Miocene Pectinidae from Santa Maria (Azores), O. de V.

Ferreira: Miocene Lamellibrancs from Portugal, O. da Veiga Ferreira (1): Miocene molluses from New Zealand. C. A. Fleming (4); The Miocene of the Alps between Bodensee and Salzach, O. Gauss & Schmidt-Thoni; Miocene molluses from Belgium, Glibert Heinzelin: Miocene fauna at Rapovce (Czechoslovakia?], Hano & Genes; Molluses of the Miocene of Holstein, Germany, W. Hinsch; Japanese miocene containing Perotrochus, Hirayama; Miocene Lucinoma species from Japan, K. Hirayama (1); New Miocene mol-luses from Tochigi Prefecture, Japan, K. Hirayama (2); Perotrochus from the Japanese Miocene, K. Hirayama (4); Miocene gasteropods from Grenada, Spain, P. Jodot; New non-marine molluses from the Miocene of Fukushima Prefecture, Japan, S. Kanno; Miocene Dosinia from Japan, S. Kanno (3); Argonautinae from the Miocene or Pliocene of Japan, Y. Kaseno: New genus of the Argonautinae from the Miocene of Japan, T. Kobayashi (1); New species of Aturia from the Japanese Miocene, T. Kobayashi (1); Presence of Aturia in the Miocene of Etchu and Iwami Provinces, Japan, Kobayashi & Mastani; Chlamys, from the Tortonian of Russia, L. N. Kugrin; Miocene molluscs from Greece, O. Kühn (1); Miocene mollusca from Noto Peninsula, Japan, K. Masuda; New Acila species from the Miocene of Japan, Nagao & Huzioka; Miocene fauna of Sicily, M. L. Nicosia: Oncophora in the Miocene of Germany, A. Papp (5); Miocene fossils from Dominica, Pilsbry & Olsson; Fauna of the Miocene of Czechoslovakia, V. Pokorný; New name for *Hermania* Puri, from the Miocene of Florida, H. S. Puri; New molluses from the Pannon of Austria, F. Sauerzopf; Planorbiidae from the Pannon of eastern Europe, F. Sauerzopf (1); Miocene fauna of south Czechoslovakia, J. Senés; Neogene macrofauna of east Czechoslovakia, J. Senés (2); Miocene fauna of Yugoslavia, P. Stevanović & Pantic; Middle Miocene fauna of Várpalota, Hungary, L. Strausz; Miocene fauna of an artesian boring at Karád, Hungary, L. Strausz (1); Neogene fauna of Czechoslovakia, J. Švagrovský; Neogene fauna of east Czechoslovakia, J. Švagrovský (1); Clithon pictus from the Miocene of Czechoslovakia, J. Švagrovský (2); Miocene pearls from the

Atlantic Coastal Plain, Baltimore, U.S.A., H. Vokes; Aquitanien molluscs from Portugal, G. Zbyszewski; Molluscs of the Aquitanian of Bavaria, H. K. Zöbelein.

Oligocene.—On concretions from the Oligocene of Germany, K. W. Geib; New Aturia from the Australian Oligocene, M. F. Glaessner; New Oligocene molluscs from Belgium, Glibert & Heinzelin: Tisoa from the Numidian of Tunisia, C. Gottis: Oligocene molluses from Jura and the Haute-Savoie, P. Jodot (1); Non-marine mollusca from the Joban coal-field, Japan, Y. Kamada; New Acila species from the Miocene-Oligocene of Japan, Nagao & Huzioka; On A. Brocchi's "Conchiologica fossile Subapennina", C. R. Ronchetti (3): Neogene macrofauna of east Czechoslovakia, J. Senes (2); Oligocene molluses from Livorno, Italy, A. Tavoni (1); Molluscs of the Aquitanian of Bavaria, H. K. Zöbelein.

Eocene.—Occurrence of Vasseuria and Belosepiella in the English Eocene, D. Carry; Palaeoecology of the Weches formation (Claiborne, Eocene) of Texas, N. M. Curtis; Tisoa from the Numidien of Tunisia, C. Gottis; An Eocene Aturia from Cyrenaica, O. Haas; Eocene molluses [from Hungary], M. Kiss-Koesisné Bányai; Eocene fossils from Udine, Italy, B. Martinis; Aturia peruviana from the Eocene? of Costa Rica, Miller, Furnish & Paguago; Aturia from Southern Chile, Miller & Furnish; New name for Hipponix floridanus from America, K. V. W. Palmer (2); New Eocene gastropod from Patagonia, J. J. Parodiz.

Mesozoic.—Stratigraphy and palaeontology of central New Caledonia, J. Avias; Mesozoic stratigraphy of the Canning Desert and Fitzroy Valley, Western Australia, R. O. Brunnschweiler; Mesozoic Corbiculidae in Europe, R. Casey; New species of Glyptophiceras from the Mesozoic of Hyogo Prefecture, Japan, K. Nakazawa & Shimizu; German Mesozoic cephalopods, G. Rosenberg; Neogene fauna of Czechoslovakia, J. Švagrovský; Neogene fauna of east Czechoslovakia, J. Švagrovský (1).

Cretaceous.—Gasteropoda from the Cretaceous of Baja California, Mexico, E. C. Allison; Cretaceous ammonites near Belgrade, Yugoslavia, M. Z.

Andjelkovič; Cretaceous of Serbia, Yugoslavia, M. Ž. Andelkovič; Neocretaceous lamellibranchs from Libya, C. Baroni & Incitti; New Cretaceous ammonite from Tunisia, E. Basse; Fossils of the "Grés de Nubie" of Yémen. É. Basse (1); Revision of the Neocretaceous mollusca of Libya, C. Bricci; Cretaceous deposits at the mouth of the Dnieper-Don rivers, U.S.S.R., G. I. Bushinskii; Lower Cretaceous Corbiculidae, R. Casey; Cretaceous Rudist types of Whitfield from Jamaica, L. J. Chubb; The Cretaceous Succession in Jamaica, L. J. Chubb (1); The Neocomian fauna and Ammonites in North Africa, M. Dalloni; On the stratigraphical position of Lower Cretaceous Ammonites in Algeria and a new genus, M. D. Delga; a new Cretaceous Emarginula from New Jersey, C. J. Eichman; The presence of Agria marticensis and Praeradiolites pulchellus in the Cretaceous of Algeria. J. Emberger (1); The Cretaceous of the eastern Carpathians, M. G. Filipescu; List of molluscs from the Cretaceous of Roumania, R. Givulescu; Cretaceous nautiloids from Western Australia, Glenister, Miller & Furnish; Parallelism in upper Cretaceous ammonites, O. Haas (2); New ammonites from the Albian of Zululand, E. C. N. v. Hoepen; New Cretaceous ammonites from Zululand, E. van Hoepen (1); New Cretaceous ammonites from Zululand, E. van Hoepen (2); The Cretaceous of Hunedoara, Rumania, D. Iacob; Stratigraphic and geographic range of Didymotis, R. W. Imlay (1); Stratigraphical implications of Belemnitella praecursor in Kansas, J. A. Jeletzky (2); Cretaceous molluses from Tunisia, A. Lapparent; Gastropods of the Grès de Taouilah, Yemen, J.-P. Lefranc; Neposiella from the Cretaceous of Gebel Manzour (Moghara-Sinai-Egypt), I. G. E.-d. Mahmoud; Cretaceous cephalopods from Moghara, Egypt, I. G. E. Mahmoud (1); Monograph of the Cretaceous Inoceramus of Japan, T. Matsumoto; Cretaceous bituberculate Pachydiscids from Hokkaido and Saghalien, Japan, T. Matsumoto; The Kossmaticeratidae from the Cretaceous of Japan, T. Matsumoto (1); Evolution of Peroniceratidae, T. Matsumoto (2); Cretaceous Desmoceratids from Hokkaido and Saghalien Japan, Matsumoto & Obata; New Durania spp. from the Mexican Cretaceous, F. K. Mullerried;

Glauconia coquandi in the Cretaceous of Eastern Europe, M. Pašić: Ancycloceras from the Maastrichtian of Russia, S. I. Pasternak; The Lamellibranchs of Padern (Aude) [S. France], G. Petter-Receveur; New Cretaceous ammonite from Austria, B. Plöchinger; Upper Cretaceous in Israel, Z. Reiss; Homeomorphy in Nigerian Cretaceous ammonites, R. A. Reyment; Cretaceous molluses from Spain, Rios & Almela; Revision of Neocretaceous Ostreidae from Libya, C. Rossi Ronchetti; New genus from the Hauterivian of Majorca, S. S. Sarkar: Maastrichtian of Henmoor, Germany, F. Schmid; Cretaceous molluscs of New Jersey, U.S.A., L. W. Stephenson; Mollusca of the Cenomanian of Texas, L. W. Stephenson (1); Cretaceous molluses from Missouri, U.S.A., L. W. Stephenson (2); British lower Cretaceous Belemnites, H. H. Swinnerton (1); New Cenomanian ammonites from Alberta, Canada, Warren & Stelck.

Jurassic.—Cephalopods of central New Caledonia, J. Avias; Jurassic fossils of the Yémen, É. Basse (1); The Jurassic of the Central Carpathians, K. Birkenmajer & J. Zuosko; Jurassic Corbiculidae, R. Casey; Upper Jurassic S.-E. England a new genus Myrene, R. Casey (1); Ammonites in relation to Jurassic formations of the Saharan Atlas (Algeria), J. Emberger (2); Syria, Jurassic Ammonites, O. Haas (1); The ammonite Taramelliceras in the Jurassic of south west Germany, H. Hölder; Jurassic of Hunedoara, Rumania, D. Iacob; Molluscs from the Jurassic of Alaska, R. W. Imlay; Macrocephalites from the Jurassic of Switzerland, A. Jeannet; Cuttlefish impressions from the Solnhofener Plattenkalk, F. Klinghardt; Jurassic Trigonians from Japan, Kobayashi & Mori; Jurassic Trigonians in Japan: Vaugoniinae, Kobayashi & Mori (1); Japanese Jurassic Trigonians, Kobayashi & Tamura; Jurassic ammonites from the Swiss Jura, P. L. Maubeuge: Ferruginous oolite of the Callovo-Oxfordien of Chaumont to Châtillon-sur-Seine, France, P. L. Maubeuge (1); Jurassic ammonites of eastern Serbia, M. Mihailović; New species of Nucula from the Portlandian of Liban [Africa], D. Mongin; Hammatoceras from the Jurassic of Japan, T. Sato; Jurassic Ammonites from north central Japan, T. Sato (1); Jurassic molluses

from Germany, F. Schnittmann; Traces of boring sponges in Actaeonella from the Austrian Jurassic, F. Schremmer; Pelecypodichnus, new genus from Jurassic of central Europe, A. Seilacher; New Jurassic ammonite from Iwaki Province, Japan, H. Yabe; New species of Pholadomya (Procardia) from the Callovian of Europe (Germany), A. Zeiss.

Lias.—Liassic molluscs from Djebel Beut-Saïdane et Dj. Zaghouan[Tunisia], G. Dubar; Fossils of the Lias of Schwäbischen Jura, Einsele & Mosebach; The Lias of Fay (Ain) Jura, S. Guérin (1); Molluscs of the Lias between Thoirette and Turgon, Jura, S. Guérin (2); Ammonites annulatus from the upper Lias? of Olten, Switzerland, K. Kapp; Two abnormal examples of Paltopleuroceras from Germany, H. Kolb; Psiloceras in the upper Lias [Germany], O. Kuhn (2); The Lias of eastern Yugoslavia, D. Veselinović.

Trias.—Myophoria from the Trias of the Brembo Valley, Italy, B. Accord; New species of Pleuronectites in the Triassic of Japan, M. Amano; Triassic cephalopods of Slovakia, D. Andrusov & Kováčik; Neo-Triassic Pelecypods from Tokyo Prefecture, Japan, K. Ichikawa; A new subspecies of Eumorphotis from the Shionosawa limestone, Japan, K. Ichikawa; Triassic Cephalopods from Austrian-Yugoslavian Alps, O. Kuhn; Triassic fauna of Idaho, U.S.A., B. Kummel; The Triassic in Germany, A. H. Müller; New Triassic Bakevellia spp. from Japan, K. Nakazawa; Ceratites from the German Triassic, H. Penndorf; Triassic molluses from Italy, A. Pollini; Triassic ammonites from the Kwanto Massif, Japan, S. Sakagami.

Palaeozoic.—New Chazyan Orthocones, R. H. Flower (2); Nomenclature of the Pelecypod family Conocardiidae, D. Nicol (2); Palaeozoic cephalopods from Wyoming and Utah, U.S.A., W. C. Sweet (2).

Permian.—Possible nautiloid mandibles from the Permian of Arizona, L. J. Brady; Permian Nuculana from Australia, R. K. Dell (6); New fossil Conularia from the West Texas Permian, North America, R. M. Finks; Eurydesma from the Permian of eastern Argentine, H. J. Harrington; Koninckioceras from the Japanese Permian, I. Hayasaka; A new Porcellia from the Permian of

Japan, I. Hayasaka (1); New Permian Foordiceras from Japan, Hayasaka & Ozaki; A new subspecies of Eumorphotis from the Shinosawa limestone, Japan, K. Ichikawa; Morphology of the fossil pelecypod Astartella, D. Nicol (1); Anthracosia from the Roumanian Permian, G. Råileanu (1); Permian Nautiloidea from the southern Ural, Ruzhentzev & Shimanskii; Cephalopods from the Sakmarskian and Artinskian of the southern Ural, V. N. Shimanskii.

Carboniferous.—Possible molluscan tracks in the Namurian of Czechoslovakia, J. Augusta & V. Pokorny; New species of Endolobus from Clore limestone, Kentucky, C. Collinson; Mississippian Goniatites from Illinois and adjacent states, C. W. Collinson (1); Protocanites (Goniatite) in the Mississippian of north-eastern Nevada, W. M. Furnish (1); Goniatites from the Pilton beds of Devon, England, R. Goldring; Non-marine fauna in the Carboniferous stratigraphy of western Europe, V. S. van der Heide; New species of Conularia from the Carboniferous of Poland, S. Kwiatkowski; Tylonautilus, from the Carboniferous of Oklahoma, Miller & Furnish (1); The Carboniferous fossil Tylonautilus in America, Miller & Furnish (2); Carboniferous cephalopods from Michigan, U.S.A., Miller & Garner; Morphology of the fossil pelecypod, Astartella, D. Nicol (1); Carboniferous paleoecology in Texas, R. E. Sloan; British Carboniferous Naiodites, A. Trueman and J. Weir.

Devonian.—Devonian molluses of Germany, A. Fuhrmann; New Devonian Gastropod family from Bohemia, R. Horný (1); New Devonian Pelecypod from Bohemia, Prantl & Růžička; Devonian Goniatites from Bohemia, F. Prantl; New Devonian Pelecypod from Bohemia, Prantl & Růžička; Prantl & Růžička (2); Devonian molluses from Germany, A. Rösler; Devonian Pectinidae of Bohemia, B. Růžička; Devonian fauna of Jesenik, Czechoslovakia, J. Skácel; Middle Devonian species of Ohio, G. A. Stewart; Upper Devonian, New York, D. H. Tesmer.

Silurian.—Trails and tentacular impressions of cephalopods in the Corryville of Cincinnati, America, R. H. Flower (3); New Silurian gastropods from central Bohemia, R. Horný; List of fossils (pp. 110-112) from the

Silurian May Hill inlier, Gloucestershire, J. D. Lawson; Pectinidae of the Silurian of Bohemia, B. Růžička; A Silurian Lamellibranch from Bohemia, B. Růžička; Paleoecology of the Maquoketa coquina beds at Graf, Iowa, P. Tasch (1).

Ordovician.—Status of endoceroid classification, R. H. Flower (1); Trails and tentacular impressions of cephalopods in the Corryville of Cincinnati, America, R. H. Flower (3); Palaeontology of the Ordovician of Missouri, R. Heller; The Ordovician Platyurus stage in Sweden, Jaanusson & Mutres; The Ordovician of Kazakstan [U.S.S.R.?], B. M. Keller; Ordovian molluses from British Columbia, T. Kobayashi (3); New Ordovician molluses from Minnesota, L. Powell; Ordovician molluses from Mendoza, Argentina, C. Rusconi; Cephalopods from the Ordovician of central Colorado, America, W. C. Sweet (1); Paleoecology of the Maquoketa coquina beds at Graf, Iowa, P. Tasch (1).

Cambrian.—Cambrian molluscs from Mendoza, Argentina, C. Rusconi.

III.—SYSTEMATIC INDEX

N.B.—Full references are given only in the case of a new genus, species or variety. In other cases reference should be made to the list of titles to the paper by the author whose name is given in the short reference in the systematic list, and under which, in the list of titles the full reference will be found. Where full reference is given in the systematic lis given in the systematic part the colume number is given in small clarendon.

Note.—The arrangement here used follows in general that of Thiele's Handbuch. The most important differences are that Epitonacea and Pyramidellacea are placed immediately preceeding Muricacea in order of grouping and the families Triviidae and Eratoidae are placed in Lamellariacea.

Where more than one page is given in a reference, the one in roman type

is where the new name is first mentioned and the one in *italic* is where the gen, sp. or var. is actually described.

Polyplacophora

Rupertomenia fodiens gen. et sp. nov. S.W. coast of Sweden, M. Schwabl, Öst. Zool. Z. 6, 1955, p. 90, f. 1-16.

Aplacophora and Placophora from the North Sea and the Baltic, S. Jaeckel.

Chitons in the Museo de la Ciudad Eva Perón, Argentina, Z. A. de Castellanos.

CHITONACEA

Chaetopleura unilineata sp. nov. Machalillo, Ecuador, E. Leloup Bull. Mus. Hist. nat. Belg. 30 40 1954: p. 1 f. 1, 2.

Chiton (Radsia) goodalli Broderip, 1832 is thought to equal C. (R.) chierchiae Nierstrasz, 1908, E. Leloup (2).

Chiton heterodon and Chiton peregrinus a general note, E. Leloup (3).

Chiton iatricus winckworthi, var. nov. North of Bhuleji, 20 miles west of Karachi, Pakistan, P. Kaas Zool. Meded. 33 1954: p. 2.

Chitons of West Coast of North America, A. G. Smith.

Crypochiton stelleri a note, E. Leloup (1).

Ischnochiton karachiensis p. 3 f. 1-6—
I. haersoltei p. 5 f. 7-9—I. ranjhai p. 8
f. 10-14 spp. nov. Manora Island,
Karachi, Pakistan P. Kaas Zool. Meded.
33 1954.

Lepidopleurus asellus life history, M. Christiansen.

Placiphorella (Placophoropsis) pacifica, deep water off Central California, Smith & Hanna.

Scrobicoplax subgen. nov. for group typified by empleura Iredale Aust. Zool. 6 1 1929 in text p. 82 under Icoplax.

Notoplax cuneata from New Zealand, W. Thomson.

Lepidozona from Japanese Islands, I. Taki (1).

Gastropoda

Gasteropod which excavate littoral rock, P. H. Fischer (1). Gastropod Fossils of the littoral, P. H. Fischer (5); Gasteropod opercula under fluorescent light, P. H. Fischer (6). Descriptive terminology of Gastropoda, L. R. Cox (2); Rate of growth of gastropoda as a function of latitude, P. A. Dehnel; Gastropoda of the Gulf of Mexico, H. A. Rehdei; Gastropods of the Brighton Area, Wasatch Mountains, E. J. & L. O. Roscoe (1).

STREPTONEURA

Aspidobranchiata

Rhipidoglossa

BELLEROPHONTACEA

Euphremitinae subfam. nov. type genus Euphemites Warthin 1930, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Tropidodiscinae subfam. nov. type gen. *Tropidodiscus* Meek and Worthien 1866, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Knightitinae **subfam. nov.** type genus *Knightites* Moore 1941, **J. B. Knight** J. Wash. Acad. Sci **46** 2 1956 p. 42.

HALIOTIDACEA

Agnesiinae subfam. nov. type gen. Agnesia Koninck 1883, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Burnesella error for Barnesella Bridge and Cloud 1947. Resembles Maclurites (Family Macluritidae, Super-fam. Euomphalacea), but is not assigned to a particular family.

†Coelocaulus chlupáči sp. nov. Silurian, Bohemia, R. Horný Sborn. geol-Úst. Čsl. 20 1953: p. 194 pl. 3 f. 1-3.

†Cinctaspira p. 46 gen. nov. Pleurotomariidae Genotype: Cinctaspira conica sp. nov. p. 46 pl. 3 f. 5; text f. 2 Oneota dolomite, Ordovician, Minnesota, U.S.A. L. Powell Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935.

Coelozoninae subfam, nov. Coelozonides tribe nov. type gen. Coelozone Perner 1907, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42. †Emarginula ladowae sp. nov. Cretaceous, Cliff of ravine at Haddonfield, Camden Co., New Jersey, C. J. Eichman Nautilus Philad. 68 4 1955 133 pl. 4 f. 7-8.

Entemnotrochus rumphii (Schepman) figured, Kuroda.

†Eobucania gen. nov. Bucaniidae, genotype: E. pulchra sp. nov. Ordovician, British Columbia, T. Kobayashi J. Fac. Sci. Univ. Tokyo (2) 9 3 1955: p. 404 pl. 1 f. 1 a-f.

†Eurydesma altum sp. nov. Permian, Argentina, H. J. Harrington J. Paleont. 29 1 1955: pl. 25 f. 1-3, 9.

Haliotis, orogenism, A. Bolognari. Haliotis lamellosa reproduction, A. Bolognari (1); H. variation, R. R. Talmadge.

†Lecanospira depressa p. 36 pl. xi f. 6-9; L. soluta p. 37 pl. xii f. 1-5; L. perplana p. 37 pl. xiii f. 1-6; L. carinata p. 38 pl. xiii f. 9-11 pl. xiv f. 1 spp. nov. Ordovician, North America, R. Heller Bienn. Rep. Mo. Bur. Geol. Min. (2) 35 1954; L. tenuis p. 66 pl. 4 f. 18-20, text-fig. 4, No. 2; L. profunda p. 67 pl. 4 f. 21, 22; text-fig. 4, No. 3 spp. nov. Oneota dolomite, Ordovician, Minnesota, U.S.A., L. Powell Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935.

Liospirinae subfam. nov. type genus Liospira Ulrich & Scofield 1897, J. B. Knights J. Wash. Acad. Sci. 46 2 1956 p. 42.

Luciellidae fam. nov. type gen. Luciella Koninck 1883, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

†Lytospira mediosulcata sp. nov. Ordovician, British Columbia, T. Kobayashi J. Fac. Sci. Univ. Tokyo (2) 9 3 1955: p. 408 pl. 1 f. 7a-c.

Monodilepas carnleyensis sp. nov' New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 48 pl. 5 f. 43, text-f. A1.

Nannoscutum gen. nov. type N. forsythi nov. Fissurellidae, Iredale Aust. Zool. 8 1937 p. 244.

Neilsoniinae subfam. nov. type gen. Neilsonia Thomas 1940, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

†Ophileta sardesoni p. 57 pl. 4 f. 4, 5, text-f. 3 No. 1; O. meeki p. 61 pl. 4

f. 6-8, text-f. 3 No. 5 spp. nov. Oneota dolomite, Ordovician, Minnesota, U.S.A., L. Powell Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935.

Ophiletinae **subfam. nov.** type gen. *Ophileta* Vamuxem 1842, **J. B. Knight J.** Wash. Acad. Sci. **46** 2 1956 p. 42.

†?Oxydiscus morcoyanus "n.sp.' Ordovician, Argentina, C. Rusconi Rev: Mus. Hist. nat. Mendoza 8 1-4 1955 p. 46 pl. 5 f. 5.

†Ozarkina parva sp. nov. Oneota dolomite Ordovician, Minnesota, U.S.A., L. Powell Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935: p. 63 pl. 4 f. 15.

†Palaeozygopleura gen. et subgen.
nov. pp. 27-29, genotype: Zygopleura
alinae Perner, 1907; P.(P.) alinae
planicostata p. 32 pl. 3 f. 1; P. (P.)
alinae multicostata p. 33 pl. 3 f. 2
subspp. nov.—P. (P.?) kettneri p. 36
pl. 4 f. 1-4; P. (P.) vesna p. 38 pl. 3
f. 3; P. (P.?) douškai p. 39 pl. 4 f. 6
spp. nov.; P. (Bojozyga) subgen. nov.
p. 41 subgenotype; P. (B.) parallela
p. 41 pl. 3 f. 6; P. (B.) mira p. 42
pl. 3 f. 5 spp. nov.; P. (Palaeozyga)
subgen. nov. p. 44 subgenotype; P.
(Palaeozyga) bohemica p. 49 pl. 6 f. 1, 2;
P. (Palaeozyga) svobodai p. 44 pl. 5
f. 5, 6; P. (Palaeozyga) tenvicostata
p. 46 pl. 4 f. 5; P. (Palaeozyga) tenvicostata
p. 46 pl. 4 f. 5; P. (Palaeozyga) hanuši p. 50 pl. 6 f. 3, 4 spp. nov.
Devonian, Bohemia, R. Horný Sborn.
geol. Úst. Čsl. 21 (paleont.) 1954.

†Palaeozygopleuridae fam. nov. p. 22. Type genus: Palaeozygopleura gen. et subgen. nov. p.*27 genotype: Zygopleura alinae Perner 1907, Devonian, Bohemia, R. Horný Sborn. geol. Úst. čsl. 21 (palaeont.) 1954.

†Palaeozygopleurinae subfam. nov. of Palaeozygopleuridae [q.v.], R. Horný Sborn. geol. Úst. čsl. 21 ser. palaeont. 1954: p. 23.

Perotrochus (Mikadotrochus) beyrichi dredged near the Izu Islands, Japan, H. Niino.

†Perotrochus sp. from the Miocene of Japan, K. Hirayama (4); Pertrochus teramachii sp. nov. south of Bungo Channel, Pacific side of Shikoku Island, Japan, T. Kuroda Venus 18 4 1955: p. 213 pl. 8 f. 1, 2.

†Phragmostoma depressum sp. nov. Devonian, Germany, A. Rösler Notizbl. Hess. Landesamt. Bodenforsch 81 1953: p. 141 pl. 7 f. 1.

Phanerotrematidae fam. nov. type gen. Phanerotrema Fischer 1885, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Planozonides Tribe nov. type gen. Planozona Perne; 1907, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Platyschismatinae subfam. nov. type gen. *Platyschisma* McCoy 1844, J. B. Knight J. Wash, Acad. Sci. 46 2 1956 p. 42.

†Plethospira extensa sp. nov. Ordovician, North America, R. Heller Bienn. Rep. Mo. Bur. Geol. Min. (2) 35 1954: p. 33 pl. 11 f. 2-4.

†Pleurotomaria prope-hirasei — P. yosiwarai "n. spp." name without description. Pliocene, Tyôsi City, Japan, H. Ozaki Bull. nat. Sci. Mus. Tokyo N.S. 1 2 1954: p. 51.

†Pseudozygopleurinae Knight, 1930 placed as subfamily in Palaeozygopleuridae fam. nov. Gastropoda, R. Horný Sborn. geol. Úst Čsl. 21 ser. palaeart. 1954: p. 26.

†Rhachopea coronata p. 48 pl. 5 f. 9-11; R. conica p. 50 pl. 5 f. 5 spp. nov. Oneota dolomite, Ordovician, Minnesota, U.S.A., L. Powell, Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935.

Raphistoma Hall. Validation, Opinion 225.

Rhaphischismatidae Fam. nov. type gen. Rhaphischisma Knight 1936, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Ruedemanniinae subfam. nov. type gen. Ruedemannia Foerste 1944, J. B. Knight, J. Wash. Acad. Sci. 46 2 1956: p. 42.

†Sinuopea virginia sp. nov. Oneota dolomite, Ordovician, Minnesota, U.S.A., L. Powell Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935: p. 53 pl. 6 f. 6.

Prosobranchiata

Prosobranchiata of tropical West Africa, J. Knudsen.

Prosobranch gastropods and comparative anatomy and Classification thereof, J. Risbec.

Docoglossa PATELLACEA

Acmaea Eschscholtz 1833 addition to Official List, Opinion 344.

Actinoleuca campbelli bountyensis subsp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 65 pl. 1 f. 9.

Cellana strigilis flemingi p. 72 pl. 5 f. 45-47; C. strigilis bollonsi p. 73 pl. 5 f. 48-50; C. strigilis oliveri p. 73 pl. 5 f. 51-53 subspp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955.

Hypseloconidae fam. nov. type genus *Hypseloconus* Berkey 1898, J. Wash. Acad. Sci. **46** 2 1956 p. 142.

Patina pellucida as food of Coalfish, R. Rendall.

TROCHACEA

†Agnathodonta brooksi sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 411 pl. 40 f. 6.

†Angaria (Angaria) gwynae sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 412, pl. 40 f. 7, 8.

Archinacellidae fam. nov. type gen. Archinacella Ulrich and Scofield 1897, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

†Astrea tyôsiensis—A. virgata 'n. spp.' name without description. Pliocene, Na-arai Conglomerate, Tyôsi City, Japan, H. Ozaki Bull. nat. Sci. Mus. Tokyo N.S. 1 2 1954: p. 51.

Brookula obscura p. 6 f. 10; B. jacksonensis p. 6 f. 11; B. augeria p. 6 f. 12; B. turbinata p. 7 f. 13; B. orospatia p. 7 f. 14; B. finesia p. 7 f. 15; B. turnida p. 8 f. 16 spp. nov. off Twofold Bay and Manly Beach, N.S.W., Australia, C. Laseron Aust. zool. 12 1954.

Callodix gen. nov. Liotiidae Genotype. C. solida sp. nov. p. 16 f. 25, 25a, b. 14 fathoms off Long Reef, Sydney, Australia, C. Laseron Aust. zool. 12 1954.

Callomphala alta sp. nov. Port Stephens, N.S.W., Australia, C. Laseron Aust. 2001. 12 1954: p. 15 f. 26, 26a.

Cantharidoscops subgen. nov. Trochidae, U.S.S.R., Galkin, Y. I. Tabl. anal. Faune URSS, 51 1955 102.

Cavostella gen. nov. Liotiidae. Genotype: C. radians sp. nov. p. 14 f. 30, 30a, Port Stephens, N.S.W., Australia, C. Laseron Aust. 2001. 12 1954.

Cavotera gen. nov. Liotiidae. Genotype: C. simplex sp. nov. p. 14 f. 31, 31a, Port Stephens, N.S.W., Australia, C. Laseron Aust. zool. 12 1954.

†Cirsochilus gilberti sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 411 pl. 40 f. 20.

Cirsonella reflecta. p 11 f. 28, 28a; C. perplexa p. 11 f. 21, 21a spp. nov. N.S.W. Australia, C. Laseron Aust. zool. 12 1954.

Conicella gen. nov. Liotiidae. Genotype: Cyclostrema porcellana Tate & May; Conicella lacuna p. 12 f. 39, 39a; C. lata p. 12 f. 37, 37a spp. nov. N.S.W. Australia, C. Laseron Aust. zool. 12 1954.

†Encycloscala emersoni sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 411 pl. 40 f.13, 19.

Ethminolia mayi 'sp. nov.' nom. nov. for the shell figured by May. Illustrated Index of Tasmanian shells 1923, pl. 18 f. 21 and there named Minolia angulata Adams, R. C. Kershaw Pap. roy. Soc. Tasm. 89 1955: p. 291.

Helisalia gen. nov. p. 16 Liotiidae Genotype: H. Liliputia p. 16 f. 45, 45a; H. pallida p. 17 f. 46, 46a; H. sucina p. 17 f. 47, 47a spp. nov. N.S.W., Australia, C. Laseron Aust. zool. 12 1954.

Hisseyagibbula subgen. nov. of Gibbula Trochacea, genotype: Littorina hisseyana Tenison Woods 1876, Tasmanian, R. C. Kershaw Pop. roy. Soc. Tasm. 89 1955: p. 291.

Liotia farallonensis sp. nov. Southeast Farallon Island, California, A. G. Smith Proc. Calif. Acad. Sci. (4) 27 13 1952: p. 385 pl. 20 f. 2-4.

Liocarinia gen. nov. Liotiidae. Genotype: Liotia disjuncta Hedley, collected by 'Thetis' expedition, 41–50 fathoms off Cape Three Points, N.S.W., Australia. Genus also includes Omalaxis radiata Hedley, Masthead Islands, Queensland, C. Laseron Aust. zool. 12 1954.

Liotella princeps p. 8 f. 17, 17a; L. littoralis p. 8 f. 18; L. parvirota p. 9 f. 19, 19a; L. patronga p. 9 f. 32, 32a, b.; *L. gravicosta* p. 10, f. 33, 33a spp. nov. N.S.W., Australia, C. Laseron Aust. zool. 12 1954.

Lissotesta arenosa sp. nov. North Harbour, Port Jackson, N.S.W., Australia, C. Laseron Aust. zool. 12 1954: p. 11 f. 22, 22a.

Lunella smaragda (Martyn) structure of stomach and sorting cæcum, figs., J. E. Morton (1).

Margarella hinemoa sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 57 pl. 1 f. 5.

Margarites helicinus Fab. observations on the species, and anatomy figs., V. Fretter (1).

Margerites (Margerites) helicina tenuistriata subspec. nov. N. Pacific, S. Sea of Japan, S. Okhotsk sea, Galkin, Y. I. Tabl. anal. Faun. URSS 57 1955 78.

Margerites (Margerites) vehlii var. angulata, var. nov, Bering, Okhotsk, Sea of Japan, Galkin, Y. I. Tabl. anal. Faun URSS 57 1955 102.

†Margarites eos sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 111 pl. 5 f. 23-32.

Meekospiridae fam. nov. type gen. Meekospira Ulrich 1897, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Microcarina gen. nov. Liotiidae. genotype: M. surgerea sp. nov. p. 17 f. 42, 42a, b. North Harbour, N.S.W., Australia, C. Laseron Aust. zool. 12 1954.

Orectospira Dall type species Basilissa babelica Dall now transferred to Turritellidae, **T. Habe** Zool. Mag. Tokyo **64** 1955 p. 259.

Orectospira Dall on its Systematic position, T. Habe (4).

Orectospira include the following species: Echinella tactiformis Watson 1888; Turritella wardana, Yokoyama 1890; Cerithium excelsum Yokoyama 1928; Cerithiopsi sikoensis, Yokoyama 1928; and Besilissa bebetica, Dall 1907, T. Habe (9).

Palaeotrochacea superfam. nov. Palaeotrochidae fam. nov. type gen. Palaeotrochus Hall 1879, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Parisanda gen. nov. Liotiidae. genotype: P. iredalei sp. nov. p. 19 f. 49, 49a. Manly Ocean beach, N.S.W., Australia, C. Laseron Aust. zool. 12 1954.

Phasionella gregaria sp. nov. abandonment on record Phyllospora Long Reef N. of Sydney, C. Laseron Pr. R. Zool. Soc. N.S.W. 1953-54 1955 p. 77, 78 figs.

Plagiothyridae fam. nov. type gen. Plagiothyra, Whidborne 1892, J. B. Knight J. Wash. acad. Sci. 46 2 1956 p. 42.

Planitrochidae fam. nov. type gen. Planitrochus Perner 1907, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

† Perotrochus in Miocene of Japan, Hirayama.

Rotostoma gen. nov. p. 15 Liotiidae. Genotype: Ethalia brazieri Angas, N.S.W. Australia, C. Laseron Aust. zool. 12 1954: p. 15 f. 24, 24a.

Solariella patriae sp. nov. Atlantic ocean, Mar del Plata, Argentina, A. Carcelles Comun. zool. Mus. Montevideo 4 70 1953: p. 1 pl. 4 f. 22.

Solariella bernaysi sp. nov. Oligocene, Belgium, M. Glibert Bull. Inst. Sci. nat. Belg. 31 86 1955: p. 3.

Stomatolina gen. nov. Stomatiidae type Stomatella rufescens Gray Iredale Aust. Zool. 8 1937 p. 245.

†Teinostoma yabei sp. nov. Miocene, Japan, K. Masuda Trans. Palaeont. Soc. Japan N.S. 20 1955 p. 123 pl. 19 f. 16-18.

Tricolia pullus (L.) observations on the species, V. Fretter (1).

Tubinidae fam. nov. type gen. *Tubina* Owen 1859, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

Zalipais laseroni sp. nov. Tasmania, R. C. Kershaw Pap. roy. Soc. Tasm. 89 1955: p. 291.

†Tectus (Tectus) tomásensis p. 410, pl. 40 f. 5; Tectus? thalmanni p. 410, pl. 40 f. 3 spp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955.

†Turbo amabilis 'n.sp.' name without description. Pliocene, Tyôsi City, Japan, H. Ozaki Bull. nat. Sci. Mus. Tokyo N.S. 1 2 1954: p. 51. Wanganella gen. nov. Liotiidae, genotype: W. fissura sp. nov. p. 13 f. 40, 40a, Port Stephens, N.S.W., Australia, C. Laseron Aust. zool. 12 1954.

LOXONEMATACEA

†Loxonema plasi sp. nov. Silurian, Bohemia, R. Horný Sborn. geol. Úst. čsl. 20 1953 : p. 193 pl. 2 f. 4, 5.

†Morania gen. nov. Loxonematidae, p. 190, genotype: M. V-sinuata sp. nov. p. 191 pl. 1 f. 1-3. Silurian, Bohemia, R. Horný Sborn. geol. Úst. Čsl. 20 1953.

NERITACEA

†Clithon (Vittoclithon) pictus ventrosus subsp. nov. Miocene, Czechoslovakia, J. Senes Geol. Práce Slovensk. Akad. 40 1955: p. 154 pl. vi f. 5, 6.

Neritina (Neripteron) See Alina tiassalensis Bandama R. sp. nov., E. Binder Rev. suisse Zool. 62 Suppl. 1955 p. 74 f. 2 a, b.

†Neritoma (Neridomus) woodfordi p. 413, pl. 40 f. 16; N. (Neridomus) delpeyae p. 413, pl. 40 f. 9, 10 spp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955.

†Theodoxus pappi sp. nov. Pannon (Tertiary) Austria, F. Sauerzopf Burgenland. Heimat Bl. 14 1 1952: p. 12 pl. 2 f. 12.

Titiscania shinkishihataii sp. nov. Hachijo-jima island, Japan, I. Taki Bull. nat. Sci. Mus. Tokyo 2 2 1955: p. 50 pl. 6-and 7.

CYCLOPHORACEA

Adelopoma brasiliense sp. nov. = Adelopoma tucma L. de Morretes 1949, (non Doering 1884), Brazil, F. L. de Morretes Arch. Mus. Paranaen. 10 1953: p. 331 pl. 24 f. 1-8.

Aperostoma (Aperostoma) schunkei p. 361 f. 70 Chanchamayo Valley; A. (Incidostoma) dilatatum p. 363 f. 71 Contamana, Rio Ucayali; A. (I.) superstructum p. 364 f. 72 Province of Lamas spp. nov. Peru, F. Hass Fieldiana zool. 34 1955.

Amphicyclotidae close to marine family Lacunidae, J. P. E. Morrison J. Wash. Acad. Sci. 45 5 1955.

Amphicyclotiis megaplanus sp. nov. along the Chiapas-Guatemala boundary, J. P. E. Morrison J. Wash. Acad. Sci. 45 5 1955 p. 160 f. 29-31.

Ampullaria pealiana Lea holotype figs., R. T. Abbott.

Atrocyclus subgen. nov. for Cyclotus horridus, Hedley, Iredale Aust. Zool. 1011941 See text p. 58.

Bellamya unicolor ugandae p. 32 f. 8a-d; B. unicolor dagusiae p. 33 f. 9a-e; B. jucunda altior p. 37 f. 11 f-g subspp. nov. Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo. 32 1954.

Buckleyia haughti sp. nov. stream N. of Rio Nuqui, Choco, Colombia, J. P. E. Morrison J. Wash. Acad. Sci. 45 5 1955 p. 150 f. 26-28.

Coxielladda nov. for Paludina gilesi Angus, Iredale & Whitley, S. Aust. Nat. 18 19 p. 66 in text under statistics of larva fannula.

Cyclochittya gen. nov. Cyclophoridae genotype Cyclotus dentistigmatus Chitty p. 154; C. schermoi sp. nov. Miocene Bowden, Jamaica, J. P. E. Morrison J. Wash. Acad. Sci. 45 5 1955 p. 154.

Diplommatina aerari p. 424 f. 1b— D. solomonensis p. 425 f. 1a sp. nov., Solomon Islands, R. K. Dell Pacif. Sci. 9 4 1955.

Ditropis koperbergi sp. nov. W. Borneo, A. Zilch Arch. Mollusk 84 4/6 1955 193 pl. 15 f. 78.

Hildapina gen. nov. Pupinidae for sabpolita, Iredale Australian Naturalist, 10 1940 p. 232.

Incerticyclus gen. nov. Cyclophoridae genotype: Neocyclotus (Ptychocochlis) bakeri Simpson 1895, J. P. E. Morrison J. Wash. Acad. 45 5 1955 p. 156.

Incidostoma jacksoni p. 158 f. 13-15; I. chocolatum p. 158 f. 10-12; I. diminutum p. 159 f. 7-9 spp. nov. Ecuador, J. P. E. Morrison J. Wash. Acad. Sci. 45 5 1955.

Lagocheilus carinulatum sp. nov. New Guinea Zilch Arch. Mollusk. 84 4/6 1955 183 pl. 14 f. 47.

Lanistes (Lanistes) kobelti p. 1 f. 3, 4 probably West Africa; L. (Meladomus) connollyi p. 2 f. 1, 2 Southern and Northern Rhodesia spp. nov; L. (M.) ovum mweruensis subsp. nov, p. 3 f. 5-7 Belgian Congo, Lake Mweru, T. Pain Breviora **31** 1954.

Liarea turriculata waipoua further records from New Zealand, N. Gardner (1).

Liarea hochstetteri alta p. 280, pl. 45 f. 13, 14; L. egea tessellata p. 286 pl. 44 f. 5; pl. 47 f. 24-26; L. aupouria tara p. 287 pl. 48 f. 34; L. turricuatal waipoua p. 291 pl. 44 f. 3, pl. 48 f. 36; L. turriculata partula p. 291 pl. 48 f. 35 subspp. nov. Liarea ornata p. 282 pl. 48 f. 37; L. aupouria aupouria p. 286 pl. 44 f. 2; pl. 48 f. 33, 33a spp. nov. New Zealand, A. W. B. Powell Rec. Aucklnad [N.Z.] Inst. 4 5 1954.

Maizaniella Bequaert & Clench types M. leonensis (Morelet) t.o.d., J. P. E. Morrison.

Palaina gardneri sp. nov. Solomon Islands, R. K. Dell Pacif. Sci. 9 4 1955: p. 424 f. le, d.

Periclocytis gen. nov. type P. ardeni nov. microevtid, p. 28 pl. ii f. 13.

Pila globosa on the liver fat, George & Desai.

Pila ovata eleanorae subsp. nov. Bujagali, Uganda Mandahl-Barth Ann. Mus. Congo belg. 8vo. 32 1954: p. 41 f. 13a-c.

Pila virens calcium distribution with special reference to the reproductive system, V. Meenakshi.

Pila virens Lmk. Physiology of digestion, V. R. Meenakshi (3).

Pila globosa—physiology of excretion, B. Saxena.

Pomacea haustrum, P. sordida gentalia and anatomy, H. de S. Lopes.

canaliculata chaquensis Pomacea subsp. nov. Argentina, M. Scott Acta zool. Lilloana 6 1948: p. 242.

Poteria clarendonensis nom. nov. for Ptychocochlis taylori Bartsch 1942 non Poteria (Eyclobakeria) welchi taylori Bartsch 1942; p. 155 P. daltei nom. nov. for Ptychocochlis welchi non P. Cyclobaberia welchii Bartsch p. 155.

†Poteria bowdenensis sp. nov. p. 155 Miocene Bowden, Jamaica, J. P. E. Morrison J. Wash. Acad. Sci. 45 5 1955.

Rugicyclotus gen. nov. Cyclophoridae genotype R. perplexus sp. nov. Appleton, St. Elizabeth, Jamaica, J. P. E. Morrison J. Wash. Acad. Sci. 45 5 1955, pp. 153-4.

Taia pseudoshanensis nom. for shanensis Kob non Theobald, A. Zilch Archiv. Mollusk. 84 173 1905 79.

† Vivipara mitzopoulosi sp. nov. Neogene, Greece, P. Psarianos Ann. géol. Hellen. 1 5 1953: p. 93 pl. 22 f. 7, 8. Viviparus viviparus L. in eastern

Canada, E. L. Bousfield.

Viviparus viviparus modestus form. nov. Czechoslovakia, O. Kapler Přírod. Sborn. Ostravsk. Kraje 16 l 1955: p. 89 pl. 2.

Viviparus viviparus anatomy, F. Starmühlner.

Viviparus viviparus, fasciatus lacustris, contectus use and misuse of names and on the identity of two species, figs., H. Watson (1).

†Vivipara mitzopoulosi 'n.sp.' [n. nud.] Pliocene, Greece, P. S. Psarianos 1951.

†Viviparus pulchriformis sp. nov. Pliocene, Greece, A. Papp & P. Psarianos Ann. géol. Hellen. 1 6 1955 : p. 145 pl. xzxiii f. 1-3.

VALVATACEA

† Valvata (Concinna) circinata jurana var. nov. Oliogocene, Jura, P. Jodot Bull Soc. géol. Fr. 6 4 7-9 1954 p. 538 pl. xxxiii b, f. 5, 6.

LITTORINACEA

†Goniophora virgata sp. nov. Devonian, Germany, A. Rösler Notizbl. Hess. Landesamt. Bodenforsch 81 1953: p. 145 pl. 7 f. 16.

Littorina obtusata (L.) and its variation, distribution and zoological notes, J. J. Barhmann; L. obtusata and variation in the North Atlantic, L. Knudson (1); L. planaxis, L. scutulata size distribution and feeding activities, W. J. North; L. Littorea (L.) the dynamics of zonation on a stony beach, J. E. Smith & Newell.

†Littorina costata sp. nov. Miocene, Czechoslovakia, J. Senes Geol. Práce Slovensk. Akad. 40 1955: p. 155 pl. vi f. 14.

Macquariella delli sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 75 pl. 1 f. 10.

RISSOACEA

Badipigrus gen. nov. Rissoidae types Rissoa badia Petterd, Iredale Pr. R. Zool. Soc. N.S.W. 1953-54 1955 p. 81.

Bithinella tenuiplicata (G. Vincent, mss.) sp. nov. Oligocene, Belgium, Glibert & Heinzelin Bull. Mus. Hist. nat. Belg. 30 1 1954: p. 8 Pl. 2 f. 4.

Bithynia and effect of acid sodium on egg, C. Attardo. Bithynia Leach, 1815—support for validation under the Plenary Powers, J. L. Bailey Bull. 2001. Nom. 11 1955: 279; B. (Gabbia) toninirri Davo near Gagnoa, E. Binder Rev. suisse Zool. 62 Suppl. 1955 p. 77 f. 9-10; Bithynia Leach, 1818—proposed valiation, A. E. Ellis; B. tentaculata Embryology, Hess (1); B. tentaculata chromosome changes in spermatogenesis, H. Keyl.

Brachypyrgulina gen. nov. Hydrobiidae genotype Brachypyrgulina carinifera sp. nov. Lago Grande—Sucuné, Lake Titicaca, F. Haas Tr. Lin. Soc. Lond. (3) 1 1955 p. 301 f. 27.

Caporista gen. nov. Rissoidae type R[issoina] iredalei Laseron, Iredale Pr. R. Zool. Soc. N.S.W. 1953-54 1955 p. 81.

Cochliopopsis Mori is a synonym of Stenothyra, Kuroda & Habe Kleest. Catal. Jap Shells No. 2 1956 p. 16.

Cyclochittya schesmoi Morrison J. Wash. Acad. Sci. 45 (5):154 Mar. 1955 should be C. schumoi, J. P. E. Morrison Nautilus Phila. 69 72.

Dardanula roseola lacteola subsp. nov. p. 88 pl. 3 f. 24; D. bollonsi p. 88 pl. 3 f. 25; D. obtusispira p. 88 pl. 3 f. 23; D. convexispira p. 89 pl. 3 f. 22 spp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955.

Dialossa gen. nov. Rissoidae type Diala translucida Hedley, Iredale Pr. R. Zool, Soc. N.S. Wales 1953-54 1955 p. 81.

Emblanda "fam. nom." type Rissoa emblematica Hedley, Iredale Pr. Roy. Soc. N.S.W. 1950-54 (1955) p. 81.

Feldestea 'gen. nov.' type R[issoina] salebrosa Frauenfeld, Iredale Pr. Roy. Soc. N.S.W. 1950-54 (1955) p. 81.

Fluvioressoina gen. nov. types B[ythynella] ramsaii, Iredale Aust. Zool. 10 3 1944 p. 332.

Gabbia (Parabithynia) kichwambae p. 53 f. 21a-d; G. (P.) parva p. 55 f. 21e spp. nov.; G. humerosa edwardi p. 55 f. 23 c-d subsp. nov. Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo. 32 1954.

†Goniochilus winkleri sp. nov. Pannon (Tertiary) Austria F. Sauerzopf Burgenland. Heimat Bl. 14 1 1952: p. 13 pl. 2 f. 1.

Haurakia trajectus (Watson) a valid species, J. Laseron (1).—Haurakia crassicosta p. 80 pl. 3 f. 27; H. minuscula p. 81 pl. 3 f. 26 spp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955.

Herewardia 'gen. nov.' type Rissoina kestoveni Hedley, Iredale Pr. Roy. Soc. N.S.W. 1950-54 (1955) p. 81.

Horatia polinskii p. 64 f. 1-3—H. brusinae p. 67 f. 5-7 spp. nov. Yugoslavia, P. Radoman Rec. Trav. Sta. Hyrdobiol. Ohrid No. 1-6 1953.

Hydrobia guyenoti sp. nov. Bay of Taupah, L. Ebrié, E. Binder Rev. suisse Zool. 62 Suppl. 1955 p. 76 f. 6-7.

†Hydrobia (Solia) 'nov. subg.' Hydrobiidae type: H. (S.) suturatocostata sp. nov. Miocene, Czechoslovakia, J. Senes Geol. Práce Slovensk. Akad. 40 1955: p. 155 pl. vi f. 16–19.—Hydrobia punctum sarmatica var. nov. p. 197 pl. xxiv f. 19–24; H. pseudosetia sp. nov. p. 197 pl. xxiv f. 25–28, Miocene, Czechoslovakia, J. Senés Geol. sbornik Slovensk. Akad. 3 3–4 1953.

†Keilostoma quaidei sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 414 pl. 40 f. 14, 15.

Limnothauma gen. nov. Hydrobiidae genotype Limnothauma crawfordi sp. nov. Isola Titicaca, Lake Titicaca, F. Haas In. Lin. Soc. Lond. (3) 1 1955 p. 302 f. 28.

Martelia tanganyicensis general description, J. Bouillon (1).

†Micromelania tauberi sp. nov. Pannon (Tertiary) Austria, F. Sauerzopf Burgenland-Heimot Bl. 14 l 1952: p. 12 pl. 2 f. 2-3.

†Mohrensternia subnitida p. 156 pl. vii f. 1-4; M. tortonica p. 157 pl. vii f. 11-20; M. krachi p. 158 pl. vii f. 25 sp. nov.; M. subnitida žižčenkovi p. 157 pl. vii f. 5-10; M. tortonica graciliformis p. 158 pl. vii f. 21-22; M. tortonica pauricostata p. 158 pl. vii f. 23-24 subspp. nov. Miocene, Czechoslovakia, J. Seneš Geol. Práce Slovensk. Akad. 40 1955.

Nodulestea 'gen. nov.' type Estea. castella Laseron, Iredale Pr. Roy. Soc. N.S.W. 1950-54 (1955) p. 81.

Notosetia antipoda p. 86 pl. 3 f. 19; N. dilatata p. 86 pl. 3 f. 21; N. sorenseni p. 87 pl. 3 f. 20 spp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955.

Nystia pseudoplicata sp. nov. Oligocene, Belgium Glibert & Heinzelin Bull. Mus. Hist, nat. Belg. 30 1 1954: p. 11 pl. 2 f. 7.

Obescrobs 'gen. nov.' type Rissoa jacksoni Brazier, Laseron, Iredale Pr. Roy. Soc. N.S.W. 1950-54 (1955) p. 81.

Oncomelania nosophora reaction to light and gravity in water, K. Yasuraoka (1).

Orbitestella decorata p. 18 f. 43, 43a, b.; O. aura p. 18 f. 44, 44a, b spp. nov. N.S.W. Australia, C. Laseron Aust. zool. 12 1954.

Paludinella solomonensis sp. nov Solomon Islands, R. K. Dell Pacif-Sci. 6 3 1955: p. 325 f. ld.

Phintorene gen. nov. type Rissoina allanac Laseron, Iredale Pr. Roy. Soc. N.S.W. 1950-54 (1955) p. 81.

Ponatiopsis cincinnationsis (Lea) distribution, ecology and life history; maps, figs. and graph, Schalie &; Dundee.

Potamopyrgus jenkinsi in Berlin and Brandenburg, H. Herold.

Rhamphopoma gen. nov. p. 298, Hydrobiidae genotype Rhamphopoma magnum sp. nov. Choccocoya Lake Titicaea p. 298 f. 24; R. parkum sp. nov. molinoparapa p. 309 f. 25: Lake Titicaea, G. Haas Tr. Linn. Soc. London (3) 1 1955.

†Rissoa (Mohrensternia) spiralocostata p. 201 pl. xxv f. 3-4, Pl. xxvii f. 33-37; R. (M.) acuta p. 203 Pl. xxv f. 7, 8, pl. xxviii f. 13-16; R. (M.) longa p. 206 pl. xxv f. 15, 16; R. (M.) cincta p. 208 pl. xxv f. 21, 22; R. (M.) minor p. 211 pl. xxv f. 26-28; R. (M.) multicostata p. 212 pl. xxvi f. 3, 4; R. (M.) multicostata minor p. 213 pl. xxvi f. 5-8; R. (M.) polonica p. 214 pl. xxvi f. 10, 11; pl. xxviii f. 34, 35; R. (M.) nuda p. 216 pl. xxvi f. 9; pl. xxx f. 12-15; R. (M.) pseudopolonica p. 217; pl. xxv f. 19-20; R. (M.) kicindica p. 219 pl. xxvi f. 22, 23; pl. xxxi f. 7-9; R. (M.) granensis p. 220 pl. xxvi f. 18, 19; R. (M.) pseudogranensis p. 221 pl. xxvi f. 20, 21; R. (M.) irregularis p. 221 pl. xxvii f. 1, 2, pl. xxxi f. 34-41; R. (M.) andrusovi p. 223 pl. xxvii f. 3, 4, pl. xxxii f. 7-12; R. (M.) friedbergi p. 223 pl. xxvii f. 24-27; R. (M.) fragilis p. 224 pl. xxvii f. 5, 6; R. (M.) čechoviči p. 225 pl. xxvii f. 7, 8; R. (M.) pupa p. 225 pl. xxvi f. 28-30 spp. nov. Miocene, Czechoslovakia, J. Seneš Geol. sbornik Slovensk. Akad. 3 3-4 1953.

Scrobs—New Zealand species, A. Hipkins.

Setaepoma gen. nov. Synceridae genotype Japonia (?) hadigeri I. & B. Rench Bougainville Island, Solomans, W. S. Clench Nautilus, Philad. 68 1955 134.

†(Solia) subgen. nov. Hydrobiidae type: Hydrobia (Solia) suturatocostata nov. sp., J. Senes.

Strombopoma gen. nov. p. 297 f. 22, Hydrobiidae genotype Littoridina (Helcobia) ortoni Pilsbry, with S. gracile sp. nov. Lagerville Lake Titicaca p. 298 f. 23, F. Haas Tr. Sci. Soc. London (3) 1 1955.

Subonoba sorenseni p. 76 pl. 2 f. 11; S. alpha p. 77 pl. 2 f. 13; S. beta p. 77 pl. 2 f.14; S. gamma p. 77 pl. 2 f. 17; S. delta p. 78 pl. 2 f. 12; S. fallai p. 78 pl. 2 f. 15; S. oliveri p. 79 pl. 2 f. 16 spp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955.

Taiwanassiminea nom. nov. for Assiminopsis Habe 1942 non Locard 1897, Kuroda & Habe Illust. Catal. Jap. Shells No. 2 1950 p. 16.

Truncatella Risso 1826 validation, Opinion 344.

Truncatellidae additional names, Clench & Turner.

Ultiscrobs gen. nov. Rissoidae type Scrobs pluteus Laseron Iredale Pr. Roy. Zool. Soc. N.S. Wales 1953-54 1955 p. 81.

CERITHIACEA

Anceya anatomy and systematics of the Tanganyikan forms, Mandahl-Barth (1).

Architectonica uruguaya sp. nov. Atlantic ocean, Argentina, A. Carcelles Comun. 2001. Mus. Montevideo 4 70 1953: p. 3 pl. 1 f. 1-6.

†Bittium tasnádii sp. nov. Eccene, Hungary, M. Kiss-Kocsisné Bányai Földt Közl. 85 3 1955: p. 369 pl. xv f. 3a, b.

Caecum (Elephantulum) berthae sp. nov. Brazil F. L. de Morretes Arch. Mus. Paranaen. 10 1953: p. 334 pl. 24 f. 9.

†Cassiope (Cassiope) kleinpelli p. 416 pl. 41 f. 6; C. (Gymnentome) gwynae p. 416 pl. 41 f. 7 spp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955.

Cerithidea californica Haldeman seasonal infection by larval trematodes, W. E. Martin.

†Cerithiopsis? stantoni sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 421 pl. 42 f. 15.

†Cerithium? emersoni sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 420 pl. 42 f. 5.—C. (Thericium) pannonicum sp. nov. Eocene, Hungary, M. Kiss-Kocsisné Bányai Foldt Kozl. 85 3 1955: p. 372 pl. xvi f. 5; C. (Pithocerithium) rubiginosum gracilis p. 60 pl. iii f. 22-24; C. (Chondrocerithium?) incertum minuta p. 61 pl. iii f. 33-42; pl. iv f. 1-4 vars. nov. Neogene, Czechoslovakia, J. Švagrovský Geol. Práce Slovensk. Akad. 37 1954.

†Cimolithium miyakoense tomásensis var. nov. p. 419 pl. 42 f. 2; C. eleanorae sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955.

Cleopatra cridlandi p. 61 f. 27a-b; C. nyanzae p. 63 f. 29a-b spp. nov.; C. cridlandi cylindrica p. 62 f. 27c-d subsp. nov. Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954. †Conocerithium (Benoistia) hungaricum p. 373 pl. xvi f. 2a, e; C. (B.) trochiforme p. 374 pl. xvi f. 1a, b spp. nov. Eocene, Hungary, M. Kiss Kocsisné Bányai Földt. Közl. 85 3 1955.

†Cryptoplocus yemenensis sp. nov. Jurassic, Yemen, E. Basse Bull. Soc. géol. Fr. 6 4 7-9 1954: p. 672 pl. 27 f. 7; text-pl. 1 f. 15.

Doryssa hohenackeri form kappleri Veonkout, anatomy pl. 4, R. T. Abbott (1).

Eularina nom. nov. for Larina Iredale, Aust. Zool. 10 1943 199 in text.

Hemisinus quaraniticus sp. nov. Argentina, M. I. Scott Physis 20 59 1954: p. 439 f. 1, 2.

Isotriphora echina sp. nov. New South Wales, Australia, C. F. Laseron Rec. Aust. Mus. 23 4 1954: p. 155 f. 23, 23a

Magnosinister gen. nov. Triphoridae, p. 157, genotype: M. hedleyi sp. nov. p. 158 f. 28 New South Wales, Australia, C. F. Laseron Rec. Aust. Mus. 23 4 1954.

Melania crenulata Physiology of digestion, V. R. Meenakshi (4); Melania embryology and development of three species, K. Romamoorthi (1).

Melanoides tuberculata butiabae subsp. nov. Lake Albert, Uganda Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954: p. 58 f. 24 f.

†Melanopsis (Canthidomus) freybergi sp. nov. Miocene, Greece, O. Kühn Ann. géol. Hellen. 1 3 1951: p. 186 pl. 17 f. 4, 5; M. (Melanosteira) mitzopoulosi p. 126 pl. xx f. 15-19; M. (M.) skurensis p. 130 pl. xx f. 27-30 sp. nov.; M. (M.) aetolica brevitesta p. 126 pl. xx f. 10-14-M. (M.) conemenosiana graciliformis p. 129 pl. xx f. 26 subsp. nov. Pliocene, Greece, A. Papp Ann. géol. Hellen. 161955.-M. (Canthidomus) costatiformis p. 108 pl. 23 f. 1-8; M. (C.) scalaritesta p. 109 pl. 23 f. 9-12; M. (C.) macrosculpturata macrosculpturata p. 109 pl. 23 f. 13-15 spp. nov.; M. (C.) macrosculpturata longitesta p. 110 pl. 23 f. 16-18 subsp. nov. Pliocene, Greece, A. Papp Ann. géol. Hellen. 1 5 1953; M. (Canthidomus) costatiformis skuphoi subsp. nov. Pliocene, Greece, A. Papp & P. Psarianos Ann. géol. Hellen. 1 6 1955: p. 148

pl. 23 f. 6-9; M. pseudaustriaca 'n. sp.' [nom. nov.] for M. austriaca Prusina, 1902 (non M. austriaca Handmann) p. 13 pl. 2 f. 5; M. pygmaea inflata subsp. nov. p. 13 pl. 2 f.4 Pannon (Tertiary) Austria, F. Sauerzopf Burgenland Heimat Bl. 14 1 1952.

†Mesalia (Mesalia) mauryae sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 414 pl. 41 f. 3.

Neclarina, "I wrote Naclarina in place and perhaps this may come into use" for Robinsonia Neville 1870, Iredale Aust. Zool. 10 1943 199 in text under Larina strangii.

†Nerinea (Plesioptygmatis) bösei p° 424 pl. 43 f. 11 text-f. 3A; N. (Plesioptygmatis) tomásensis p. 425 pl. 43 f. 10, 12 text-f. 3B, 3E spp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955.

Notosinister univitta p. 145 f. 2; N. conferta p. 145 f. 3; N. tricolor p. 146 f. 4; N. sarcira p. 146 f. 5; N. robusta p. 147 f. 7; N. pocula p. 148 f. 8; N. topazica p. 149 f. 11, 11a; N. alborda p. 149 f. 12, 12a, 12b; N. glaciala p. 150 f. 13, 13a; N. jacksonensis p. 150 f. 14, 14a, 25, 25a; N. fulvalinearis p. 153 f. 20, 20a; N. stramentia p. 154 f. 22; N. grandiosa p. 155 f. 30 spp. nov. New South Wales, Australia, C. F. Laseron Rec. Aust. Mus. 23 4 1954.

†Pirenella kytherensis 'n.f.' Pliocene, Greece, A. Papp Ann. géol. Hellen. l 1 1947 : p. 114 pl. ix f. 1-4.

Potadoma vogelis sp. nov. Agnéby à Agboville p. 79 f. 14, 15; P. rehnis sp. nov. Bay Aboisse, p. 81 f. 18, E. Binder Rev. suisse Zool. 62 Suppl. 1955.

†Pseudomalaxis pateriformis sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955: p. 124 Pl. 21 f. 19-21.

†Pseudonerinea stantoni p. 423 pl. 43 f. 7—P. stirtoni p. 423 pl. 43 f. 1, 2 spp. nov. Cretaceous, Mexico E. C. Allison J. Paleont. 29 3 1955.

† Pyrazus (Echinobathra) durhami p. 419 pl. 42 f. 12; Pyrazus? harrissi p. 420 pl. 42 f. 1 spp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955.

†Seila (Seila) costatonodosa p. 62 pl. iv f. 14, 17-21, pl. v f. 1, 6; S. (S.) conica p. 62 pl. v f. 2-5, 8, 9 spp. nov. Neogene, Czechoslovakia, J. Švagrovský Geol. Práce. Slovensk. Akad. 37 1954.

Solosinister gen. nov. Triphoridae genotype: Solosinister pagoda sp. nov. New South Wales, Australia, C. F. Laseron Rec. Aust. Mus. 23 4 1954: p. 157 f. 27, 27a.

Sundaya campbellica sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 92 pl. 2 f. 18.

Syrnolopsis anatomy and systematics of the Tanganyikan forms, Mandahl-Barth (1).

Telescopium mauritsi Butot (Basteria 18 1–13 pl. 1) Butot Treubia 23 1955 p. 110 pl. 4 f. 2.

†Trobus gen. nov. Turritellidae: type species: T. buboanus sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955: p. 127 pl. 22 f. 23-25.

†Turritella? durhami sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3, 1955: p. 415 pl. 40 f. 17; ? Turritella (Archimediella) brebioni nom. nov. for T. (A.) choffati J. Böhm 1929 (non T. choffati Thomas and Peron 1889) E. Dartevelle & J. Roger Comun. Serv. geol. Portug. 35 1954: p. 250.

†*Uchauxia kirki* p. 417 pl. 42 f. 3; *U. beali* p. 418 pl. 42 f. 8 spp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955.

†Voysa? cuniculana sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264-B 1954: p. 38 pl. 8 f. 19.

AMALTHEACEA

†Hipponix levinus nom. nov. for H. floridanus Palmer, 1953 (December) non H. floridanus Olsson and Harbison 1953 (November), Eocene, Florida, K. V. W. Palmer J. Paleont. 29 3 1955: p. 558.

CALYPTRAEACEA

†Calyptraea aokii sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 117 pl. 4 f. 1, 2. Calyptraeidae, Notes on the family, P. H. Fischer (2).

Crepidula gravispinosa n. sp. for Crepidula aculeate (not Gmelin) Kuroda & Habe Illust. Catal. Jap. Shells No. 4 1950 p. 30.—Crepidula fornicata Anatomy and biology, B. Werner.

†Lirpsa? lepida sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264-B 1954: p. 36 pl. 8 f. 6-13.

Progalerinae subfam. nov. type gen. Progralerus Holzapfel 1895, J. B. Knight J. Wash. Acad. Sci. 46 2 1956 p. 42.

STROMBEACEA

†Anchura (Anchura) andersoni p. 427, pl. 44 f. 3; A. (Perissoptera) hannai p. 427, pl. 44 f. 2 spp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955.

†Helicaulax formosa sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955: p. 128 pl. 21 f. 7-9.

†Lispodesthes lirata sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-C 1955: p. 57 pl. 4 f. 6-11.

Strombus gigas verrilli and S. pugilis nicaraguensis, new records from Jamaica, V. A. Zans.

ATLANTIDACEA

Atlanta pacifica sp. nov. Japan, T. Tokioka Publ. Seto. mar. biol. Lab. 4 2-3 1955: p. 249 pl. vvii f. d-g.

Carinaria japonica sp. nov. 36° 21′ N 141° 00′ E about 30 miles off Oarni-saki Ibarage Prefecture Japan Okutani Venus 18 4 1955 251 f. 1–3.

NATICACEA

Lunatia nitida boring and feeding habits, E. Ziegelmeier.

Natica caneloensis sp. nov. Mazatlan, Mexico to Manta, Ecuador, Hertlein & Strong Bull. Amer. Mus. nat. Hist. 107 2 1955 p. 287 pl. 2 f. 13 18.

Naticidae, on the eggs and larvae, M. Amio.—Naticidae: structure and formation of egg masses, M. Giglioli.—Naticidae of Western Mexico and Panama [A. M. Strong] (2).

Polinices duplicatus and method of capturing Ensis directus, H. J. Turner, inr.

LAMELLARIACEA

Ellatriva memorata from New Zealand, K. Watene.

Trivia dartevellei sp. nov. Landana Cabinda, West Africa shore, J. Knudsen Rev. Zool. Bot. Afr. 51 1955 p. 122 f. 8, 9.

CYPRAEACEA

Cypraea deposit of tropical species in the Netherlands, W. S. S. & B. Jutting (1).—C. aurantium Martyn notes and price \$275 fig., M. C. Teskey.

Erronea caurica longior Iredale on Long Reef (notes on the animal), Rutland & Kerslake.

Evenaria ursellus marcia Iredale (earlier name E. kieneri Hidalgo (on the animal) Long Reef, Rutland & Kerslake.

Gratiadueta Xanthodon Swby (notes on the animal) fairly common of central Queensland mainland, Rutland & Kerslake.

TONNACEA

Argobuccinum—the South American species, A. Carcelles (1).

†Argobuccinum (Haurokoa) subgen. nov. Cymatiidae type species: A. (H.) woodi sp. nov. p. 1056 pl. 41 f. 16 Miocene, New Zealand, C. A. Fleming Trans. rov. Soc. N.Z. 82 5 1955.

Bursidae of Indo-China, M. Marche-Marshad (1).

Casmaria perryi first record from Australia, G. Thornley (1) p. 33 f. 6, 8.

Cymateum echo n. sp. for Cymatium parthenopeum (not Salis) Kuroda & Habe Illust. Catal. Jap. Shells No. 4 1950 p. 30.

Cymatiidae of Indo-China, M. Marche-Marchad (1).

Eudolium thompsoni sp. nov. dredged off Tampa, Florida, P. L. McGinty Proc. nat. Acad. Sci. Philad. 127 1955 p. 80 pl. 1 f. 5, 6.

†Haurokoa subgen. nov. of Argobuccinum Cymatiidae. Miocene, New Zealand, C. A. Fleming Trans. roy. Soc. N.Z. 82 5 1955; p. 1055.

Phalium iheringi sp. nov. Atlantic Ocean, Mar del Plata, Argentina, A. Carcelles Comun. zool. Mus. Montevideo 4 70 1953: p. 5 pl. 2 f. 7-10.

Tonna Brünnich 1771 added to official list, Opinion 237.

Xenogalea angasi knighti subsp. nov. Gloucester passage, Bowen, Queensland, Australia, G. Thornley Proc. R. zool. Soc. N.S.W. 1952–53 1954: p. 33 pl. i f. 6.

STENOGLOSSA

EPITONIACEA

Elasmonematidae fam. nov. type gen. Elasmonema Fischer 1885, J. B. Knight J. Wash. Acad. Sci. 45 2 1956 p. 42.

Epitonium trevelyanum in the Netherlands, C. O. & R. Altena.

†Epitonium (Boreoscala) tenerum sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 115 pl. 4 f. 17-19.

Janthina general note, A. H. Jones.

Aglossa

PYRAMIDELLACEA

Balcis devians (Monts.) and B. alba (Costa). Habits and movement and other observations, figs., V. Fretter.

†Bartschella parkeri sp. nov. Pliocene, Florida, P. Bartsch Smithson misc. Coll. 125 2 1955: p. 36 pl. 6 f. 4a, b.

†Chrysallida aldrichi p. 61 pl. 15 f. 13; C. weberi p. 62 pl. 15 f. 8; C. dalli p. 63 pl. 15 f. 6; C. cookei p. 64 pl. 15 f. 10; C. gardnerae p. 64 pl. 15 f. 7; C. harrisi p. 65 pl. 15 f. 12; C. palmerae p. 66 pl. 15 f. 4; C. mansfieldi p. 67 pl. 15 f. 14; C. macneill p. 68 pl. 15 f. 15; C. simpsoni p. 69 pl. 15 f. 11; C. woodringi p. 70 pl. 15 f. 9; C. vaughani p. 70 pl. 15 f. 2; C. simpsoni p. 71 pl. 16 f. 11; C. leai p. 72 pl. 16 f. 8; C. locklini p. 73 pl. 16 f. 10; C. sayi p. 74 pl. 16 f. 13; C. smithi p. 75 pl. 16 f. 12; C. pilsbryi p. 76 pl. 16 f. 7; C. tuomeyi p. 77 pl. 16 f. 14; C. holmesi p. 78 pl. 16 f. 9 spp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955.

Chryseulima gen. nov. Melancllidae genotype Stilifer frazicri Angas p. 87 pl. 1 f. 14, 15; C. solitaria sp. nov. Long Reef, N.S.W. p. 88 pl. 1 f. 16, 17, Laseron Aust. Zool. 12 1955.

Curveulima gen. nov. Melanellidae p. 88 type Curvevlima cornuta sp. nov. p. 89 pl. 1 f. 18, 19 Ocean Beach, Manly; C. obtusa sp. nov. p. 89 pl. 1 f. 20, 21 Ocean Beach, Manly; C. lata 14 fms. off Long Reef p. 90 pl. 1 f. 22; C. subobtusa Long Reef, p. 90 pl. 1 f. 23; C. litoris Manly Beach p. 90 pl. 1 f. 24, 25; C. abrupta Ocean Beach, Manly p. 90 pl. 1 f. 26, 27; C. manifesta 30–35 fms. off Crookhaven p. 91 pl. 1 f. 28, 29 spp. nov. N.S. Wales, Laseron Aust. Zool. 12 1955.

Cuspeulima gen. nov. Melanellidae genotype Leisotraca acutissima Sowerby p. 91 pl. 2 f. 32, 33; C. sobrina sp. nov. Port Stephens, p. 92 pl. 2 f. 35, 36; C. iredalei sp. nov. Pt. Halliday p. 93 pl. 2 f. 37, 38; C. portensis Port Hackling p. 93 pl. 2 f. 39-41; C. incidenta sp. nov. N. Harbour, Port Jackson p. 94 pl. 2 f. 42, 43 N. S. Wales, Laseron Aust. Zool. 12 1955.

†Devonzyga gen. nov. Palaeozygopleuridae fam. nov. (Gastropoda) p. 35, genotype: D. perneri p. 53 pl. v f. 2, 3; D. diversicostata p. 54 pl. 5 f. 1 sp. nov. Devonian, Bohemia, R. Horný Sborn. geol. Úst. Čsl. 21 ser. palaeont. 1954.

†Eulimastoma harbisonae p. 82 pl. 16 f. 1; E. olssoni p. 83 pl. 16 f. 5 spp. nov. Pliocene, Florida, P. Bartsch Smith. misc. Coll. 125 2 1955.

Euliamaustra gen. nov. Melanellidae genotype Eulima proxima Swby, p. 85 pl. 1 f. 1, 2; E. anomala sp. nov. Woolgoolga N. S. Wales p. 85 pl. 1 f. 1, 2, Laseron Aust. Zool. 12 1955.

†Eulimella (Eulimella) tampaensis p. 13 pl. 2 f. 1—E. (Cossmannica) pinellasi p. 14 pl. 2 f. 2 spp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955.

Eulimitra gen. nov. Melanellidae genotype Eulimitra vittata sp. nov. N. Harbour, Port Jackson; p. 94 pl. 2 f. 44, 45; E. waltersi sp. nov. Woollahra Point, Port Jackson p. 94 pl. 2 f. 46, 47, N. S. Wales, Laseron Aust. Zool. 12 1955. Eulimoda gen. nov. Melanellidae genotype Eulima munito Hedley (N. S. Wales), Laseron Aust. Zool. 12 1955 p. 98 pl. 3 f. 66, 67.

Eulitoma gen. nov. Melanellidae genotype Eulitoma nitens sp. nov. 6-9 fms. Sow and Pig Reef p. 96 pl. and f. 54, 55; E. castanea sp. nov. 14 fms. Port Jackson p. 96 pl. 2 f. 56, 57 (N. S. Wales) Laseron Aust. Zool. 12 1955.

† Fargoa gen. nov. Pyramidellidae, genotype: F. calesi p. 80 pl. 16 f. 2; F. archeri p. 81 pl. 16 f. 4 spp. nov. Pliocene, Florida, P. Bartsch Smith. misc. Coll. 125 2 1955.

Fusceulima gen. nov. Melanellidae genotype Fusceulima jacksonensis sp. nov. N. Harbour, Port Jackson p. 95 pl. 2 f. 48, 49; F. flava sp. nov. 40–50 fms. off Twofold Bay p. 95 pl. 2 f. 50, 51; F. sucina sp. nov. 8–10 fms. off Point Halliday p. 96 pl. 2 f. 52, 53 (N. S. Wales) Laseron Aust. Zool. 12 1955.

Hebeulima gen. nov. Melanellidae genotype Leiostraca inusta Hedley p. 97 pl. 3 f. 58, 59; H. tumire sp. nov. Port Stephens p. 97, pl. 8 f. 60, 61; H. crassiceps Port Halliday p. 97 pl. 3 f. 63, 64; spp. nov. N. S. Wales, Laseron Aust. Zool. 12 1955.

†Iolaea wagneri sp. nov. Pliocene, Florida, P. Bartsch Smith. misc. Coll. 125 2 1955: p. 59 pl. 15 f. 3.

†Locklinia gen. nov. Pyramidellidae, genotype: L. fargoi p. 11 pl. 2 f. 8; L. pliocena p. 12 pl. 2 f. 3; L. ornata p. 12 pl. 2 f. 7 spp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955.

†Longchaeus (Pharcidella) calesi p. 9 pl. 1 f. 4; L. (Longchaeus) marionae p. 10 pl. 1 f. 2, 3 spp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955.

Macertexta gen. nov. Melanellidae genotype Macertexta ovitesta sp. nov. N. Harbour, Port Jackson, N. S. Wales, Laseron Aust. Zool. 12 1955 p. 101 pl. 3 f. 75.

Miralda (Miraldella) subgen. nov. type: M. (M.) gordonae sp. nov. Pliocene, Florida, P. Bartsch Smith misc. Coll. 125 2 1955: p. 79 pl. 16 f. 3.

†Miraldella subgen. nov. of Miralda [q.v.] Pyramidellidae Pliocene, Florida, P. Bartsch.

†Mormula vaughani p. 27 pl. 6 f. 3a, b; M. gardnerae p. 28 pl. 4 f. 10a, b; M. mansfieldi p. 29 pl. 4 f. 9a, b; M. cookei p. 30 pl. 5 f. 2a, b; M. woodringi p. 30 pl. 4 f. 8a, b; M. marshalli p. 31 pl. 5 f. 5a, b; M. harrisi p. 32 pl. 5 f. 1a, b; M. palmerae p. 33 pl. 5 f. 4a, b; M. pilsbryi p. 34 pl. 6 f. 2a, b; M. robertsonae p. 34 pl. 5 f. 3a, b; M. teskeyae p. 35 pl. 6 f. 1a, b spp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955.

†Odostomia (Evalea) emeryi p. pl. 17 f. 1; O. (E.) pomeroyi p. 84 pl. 17 f. 2; O. (E.) willcoxi p. 86 pl. 17 f. 3; O. (Odostomia) heilprini p. 87 pl. 17 f. 5; O. (O.) johnsoni p. 87 pl. 17 f. 6; O. (O.) conradi p. 88 pl. 17 f. 9; O. (O.) gabbi p. 88 pl. 17 f. 10; O. (O.) stephensoni p. 89 pl. 17 f. 7; O. (O.) schwengelae p. 90 pl. 17 f. 8; O. (O.) gunteri p. 90 pl. 17 f. 11; O. (O.) matsoni p. 91 pl. 18 f. 6; O. (O.) burnsi p. 91 pl. 18 f. 5; O. (O.) nicoli p. 92 pl. 18 f. 3; O. (O.) bassleri p. 92 pl. 18 f. 2; O. (O.) cooperi p. 93 pl. 18 f. 4; O. (O.) stearnsi p. 93 pl. 18 f. 8; O. (O.) coxi p. 94 pl. 18 f. 7; O. (O.) pinellasensis p. 95 pl. 18 f. 1 spp. nov. O. (Evalea) caloosaensis (Dall) = O. (Syrnola) caloosaensis Dall (part) 1892. Pliocene, Florida P. Bartsch Smith. misc. Coll. **125** 2 1955.

†Orinella (Sulcorinella) locklini p. 15 pl. 2 f. 5; O. (Orinella) pliocena p. 15 pl. 2 f. 9 spp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955.

†Peristichia martschi sp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955: p. 16 pl. 2 f. 6.

Pictobalcis gen. nov. Melanellidae genotype Eulima articulata Swby. Laseron Aust. Zool. 12 1955 p. 98 pl. 3 f. 65.

†Pseudomelania? basicostata sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-C 1955: p. 57 pl. 4 f. 4, 5.

† Pyrgiscus yamai p. 38 pl. 6 f. 6a, b; P. zethusi p. 38 pl. 7 f. 7a, b; P. vishnui p. 39 pl. 7 f. 2a, b; P. venusae p. 40 pl. 6 f. 5a, b; P. tityusi p. 40 pl. 7 f. 1a, b; P. thestiusi p. 41 pl. 7 f. 8a, b; P. tellusae p. 41 pl. 7 f. 5a, b; P. clionae p. 42 pl. 7 f. 4a, b; P. tantalusi p. 43 pl. 7 f. 6a, b; P. sylvanusi p. 43

pl. 7 f. 3a, b; P. somnusi p. 44 pl. 8 f. 4a, b; P. sisyphusi p. 44 pl. 8 f. 5a, b; P. silenusi p. 45 pl. 8 f. 3a, b; P. pyrrhusi p. 45 pl. 8 f. 1a b; P. apolloi p. 46 pl. 8 f. 2a, b; P. dianae p. 47 pl. 9 f. 2a, b; P. latonae p. 47 pl. 9 f. 2a, b; P. latonae p. 47 pl. 9 f. 8, b; P. phaetoni p. 48 pl. 8 f. 6a, b; P. harmoniae p. 49 pl. 9 f. 8a, b; P. cadmusi p. 49 pl. 9 f. 6a, b; P. inoae p. 50 pl. 9 f. 5a, b; P. telamoni p. 51 pl. 9 f. 3a, b; P. hebeae p. 52 pl. 9 f. 4a, b; P. aesoni p. 52 pl. 9 f. 7a, b, c. spp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955.

†Salassia (Salassia) fargoi p. 58 pl. 15 f. 1; S. (Salassiella) balchi p. 58 pl. 15 f. 5 spp. nov. Pliocene, Florida, P. Bartsch Smithson, misc. Coll. 125 2 1955.

Sinusicola gen. nov. Hydrobiidae for Turbonilla (Careliopsis) filiola Yokoyama, Kuroda & Habe Illust. Catal. Jap. Shells No. 2 1956 p. 16.

Stilimella gen. nov. Melanellidae genotype Stilifer lodderae Petterd J. of Conch. IV, 1884 p. 140, Laseron Aust. Zool. 12 1955 p. 100 pl. 3 f. 72-74.

Stricteulima gen. nov. Melanellidae genotype Stricteulima cameroni sp. nov. N. S. Wales, Laseron Aust. Zool. 12 1955 p. 91 pl. 2 f. 30, 31.

Striopyrgus 'new pseudogenus' [hybrid genus] Pyramidellidae type: S. hybridus sp. nov. Florida, P. Bartsch Smithson, misc. Coll. 125 2 1955: p. 57 pl. 10-14 f. 1-61.

Syntharella gen. nov. Melanellidae genotype Eulima topaziaca Hedley (N. S. Wales), Laseron Aust. Zool. 12 1955 p. 100 pl. 3 f. 71.

†Triptychus pliocena sp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955: p. 8 pl. 1 f. 1.

†Turbonilla (Chemnitzia) acisi p. 19 pl. 3 f. 2; T. (C.) admeta p. 19 pl. 3 f. 6a, b; T. (C.) geryoni p. 20 pl. 3 f. 5; T. (C.) hippolyta p. 20 pl. 3 f. 4a, b; T. (C.) adonisi p. 21 pl. 3 f. 8; T. (C.) alemena p. 21 pl. 3 f. 1; T. (C.) iolausi p. 22 pl. 3 f. 3; T. (C.) hydra p. 22 pl. 3 f. 9; T. (C.) augeasi p. 23 pl. 3 f. 7; T. (C.) eurytioni p. 23 pl. 4 f. 1; T. (C.) hesperusi p. 24 pl. 4 f. 2; T. (C.) antaeusi p. 24 pl. 4 f. 5a, b; T. (C.) antaeusi p. 25 pl. 4 f. 6; T. (C.) terra p. 25 pl. 4 f. 3; T. (C.) cacusi p. 26 pl. 4 f. 4;

T. (C.) cerberusi p. 26 pl. 4 f. 7 spp. nov. Pliocene, Florida, P. Bartsch Smithson. misc. Coll. 125 2 1955.

†Ugartea locklini sp. nov. Pliocene, Florida, P. Bartsch Smithson, misc. Coll. 125 2 1955: p. 17 pl. 2 f. 4.

STENOGLOSSA

(a) RACHIGLOSSA

MURICACEA

Chorus from South America, A. Carcelles (2).

Concholepas in South America, A. Carcelles (2).

Liniaxis gen. nov. Parpuridae genotype Liniaxis elongata sp. nov. 30-35 fms. off Crookhaven, Laseron Pr. R. Zool. Soc. N.S.W. 1953-54 1955 p. 72 f. 3-6.

†Morea transenna sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955: p. 129 pl. 22 f. 6-9.

Murex clenchi sp. nov. Atlantic ocean, Argentina, A. Carcelles Comun. zool. Mus. Montevideo 4 70 1953: p. 7 pl. 5 f. 23-28; M. (Murex) finlayi sp. nov. 100 fms. Mantanzas Bay, Cuba, W. S. Clench Breviora No. 44 1955 pp. 1-2 figs. 1-3; M. trunculus, oology, F. Dulzetto.

Muricacea of the Eastern Pacific, A systematic outline, J. Q. Burch.

Neothais smithi another record from New Zealand, E. Richardson.

Nucella—from South America, A. Carcelles (2).

Oppomorus subgen. nov. for Morula nodulifera Menke on reef Lord Howe Id., Iredale Aust. Zool. 8 4 1937, in text p. 258 under Morula nodulifera.

Phyllonotus of the Gulf of California, J. & C. Cate.

Purpura lapilus L. and luminescence, P. H. Fischer (3).

Rhombothais gen. nov. p. 37 Thaididae, super-family Muricacea. Genotype: R. arbutum sp. nov. p. 38 pl. 3 f. 1, 2, Angourie, $3\frac{1}{2}$ miles south of Clarence River Heads, N.S.W., Australia, L. Woolacott Proc. R. zool. Soc. N.S.W. 1952-53 1954.

Thais (Taisella) mariae sp. nov. Brazil, F. L. de Morretes Arch. Mus. Paranaen. 10 1953: p. 339 pl. 25, 26. Tolema Iredale 1929 genotype T. australis Laseron not Purpura sectata Hedley as used by Iredale, Laseron (4).

BUCCINACEA

Acochlidiacea—general note and anatomy, E. & E. Marcus (2).

Aeneator attenuatus nothopanax subsp. nov. Pliocene, New Zealand, C. A. Fleming N.Z. J. Sci. Tech. 35B 5 1954: p. 382 f. 9-12.

Anentome helena notes, Butot (1).

Aquilofusus ravni sp. nov. Oligocene, Belgium, M. Glibert Bull. Inst. Sci. nat. Belg. 31 86 1955: p. 4.

Busyconinae, A Catalogue genera, subgenera and species with notes, Puffer & Emerson.

Columbellidae of Western Mexico and Panama, [A. M. Strong] (3).

†Colus (Aulacofusus) fujimotoi sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 122 pl. 5 f. 1-7.

†Euthriofusus wenzi sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 29 3 1955: p. 428 pl. 44 f. 10-12.

Fusilatirus gen. nov. Fasciolariidae genotype F. pauli sp. nov. dredged off Sombrero Key Light, Florida, T. L. McGinty Proc. nat. Acad. Sci. Philad. 107 1955 p. 79 pl. 1 f. 4, 4a.

Fusus caparti sp. nov. South Atlantic, West African coastal waters, W. Adam & J. Knudsen Bull. Inst. Sci. nat. Belg. 31 61 1955: p. 12 pl. 2 f. 8-10; text-f. 3; Fusus Embryology and Anatomy, figs., A. Portmann.

† Kelletia brevis 'n.sp.' name without description. Pliocene, Na-arai Conglomerate, Tyôsi City, Japan, H. Ozaki Bull. nat. Sci. Mus. Tokyo N.S. 1 2 1954: p. 51.

Lathyrus frenguellii sp. nov. Atlantic Ocean, S.S.E. of Quequén, Argentina, A. Carcelles Comun. zool. Mus. Montevideo 4 70 1953: p. 8 pl. 3 f. 13-16, 19.

†Molopopholus rara sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 119 pl. 5 f. 8-10.

Nassarius attacked and bored, Alexander.

Nassidae of Indo-China, M. Marche-Marchad.

Paxula sorenseni sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 109 pl. 1 f. 6.

Penion Fischer = Verconella Iredale 1914; Penion brazieri sp. nov. p. 1057 pl. 41 f. 22 Miocene, New Zealand, C. A. Fleming Trans. roy. Soc. N.Z. 82 5 1955.

†*Priscofusus ishijimai* sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 121 pl. 5 f. 14–19.

Zemitrella constans sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955 : p. 107 pl. 1 f. 7.

VOLUTACEA

Alcithoe (Leporemax) fusus resolutionensis subsp. nov. Pliocene, New Zealand, C. A. Fleming N.Z. J. Sci. Tech. 35B 5 1954: p. 383 f. 19, 20.

Ancilla Lamarck, on the type species, Hemming Bull. zool. Nom. 9 1954: p. 219; Ancilla Lamarck, request for a ruling on the question of the type species, K. Palmer; Ancilla comment on the type species, K. Palmer Bull. zool. Nom. 9 1954: p. 269.

Benthovoluta gen. nov. Volutidae genotype Phenacoptygma? kiiensis Kuroda = Voluta hilgendorfi v.Martens, Kuroda & Habe Illust. Catal. Jap. Shells No. 5 1950 p. 37 pl. 5, f. 2 pl. 7 f. 1 text-fig. 1.

†Cancellaria rara sp. nov. Miocene, Fukushima Prefecture, Japan, S. Aoki Sci. Rep. Tokyo Kyoiku Daig. 3C 1954: p. 38 pl. 2 f. 17; Cancellaria, C. R. Ronchetti.

†Eoharpa gen. nov. Harpidae, p. 131, type species: Eoharpa sinuosa sp. nov. p. 132 pl. 23 f. 3-6. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955.

Ericusa sowerbyi porcellana subsp. nov. vicinity of Habo Island, Victoria, Australia, N. Jackson Proc. R. zool. Soc. N.S.W. 1952-53 1954: p. 37 pl. 2 f. 2.

Fulgoraria prevostiana magna var. nov. Hitachi, Honshu p. 34 pl. 6 f. 3, 4; F. cancellata sp. nov. off Tosa, Shikoku 100 fms. p. 34 pl. 5 f. 11, Kuroda & Habe Illust. Catal. Jap. Shells. No. 5 1950; F. kaneko [Hirase M.S.) new species = Fulgoraria kaneko Hirase Illust. Thousand shells 4 f. 319, where name and figure is given [which is a description] neotype designated as original type destroyed, Kuroda & Habe.

†Halia, C. R. Ronchetti.

Iredalina aurantia from New Zealand coastal waters, S. Trigance (1).

†Liopeplum rugosum sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955: p. 130 pl. 22 f. 1-5.

†Lyria C. R. Ronchetti (1).

Marginella corderoi sp. nov. Atlantic ocean, Argenaina, A. Carcelles Comun. zool. Mus. Montevideo 4 70 1953: p. 10 pl. 3 f. 17, 18.

†Marginella, C. E. Ronchetti (2).

Melo amphora knighti subsp. nov. Gloucester passage, Bowen, Queensland, Australia, N. Jackson Proc. R. zool. Soc., N.S.W. 1952-53 1954; p. 35 pl. 2 f. 1.

Mitra moisei p. 77 pl. 1 f. 4, 4a; M. olssoni p. 78 pl. 1 f. 3, spp. nov. Florida, McGinty Proc. Acad. Sci., Philad. 107 1955.

Mitridae of West Mexico and Panama, [A. M. Strong] (1).

†Paladmete pristina sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S., geol Surv. 264-B 1954: p. 39 pl. 8 f. 24-26.

Voluta a historical review of the Linnaean genus, H. Dodge.

Volutidae of Western Mexico and Panama, [A. M. Strong] (1).

† Volutilithes, C. R. Ronchetti.

STENOGLOSSA

(β) Toxoglossa

CONACEA

Apispiralia gen. nov. Turridae genotype Clathurella albocincta Angas pp. 39 and 40 pl. 10 f. 203, 204: A. catena 14 fms. off Long Reef p. 40 pl. 10 f. 205, 206; A. maxima 10-15 fms. off Twofold Bay p. 40 pl. 10 f. 207: spp. nov. N.S.W., C. F. Laseron, Review of the New South Wales Turridae, Sydney, 1954.

Apitua gen. nov. Turridae genotype A. delicatula sp. nov. dredged North Harbour Port Jackson, C. F. Laseron, Review of the New South Wales Turridae, Sydney 1954, p. 42 pl. 10 f. 215, 216.

Asperdaphne rugosa 30-35 fm. off Cookhaven p. 46 pl. 11 f. 235-237; A. sepulta Sow and Pigs Reef, Port Jackson p. 46 pl. 11 f. 238, 239; spp. nov. Turridae N. S. Wales, C. F. Laseron, Review of the New South Wales Turridae, Sydney 1954.

Austrocarina gen. nov. Borsoniinae, Turridae, genotype Leucosyrinx recta Hedley, "temporarily placed in the Borsoniinae," C. F. Laseron, Review of the New South Wales Turridae, Sydney 1954 p. 21 pl. 4 f. 88.

Austrodaphnella gen. nov. Daphnellinae genotype A. clathrata sp. nov. dredged Pittwater N.S.W. p. 45 pl. 11 f. 228, 229; C. F. Laseron Review of the Turridae, Sydney, 1954.

Austrogemmula gen. nov. Turridae genotype Hemipleurotoma tasmanica May, C. F. Laseron Review of the New South Wales Turridae, Sydney, 1954, pl. 1 f. 8, 9.

Austropusilla gen. nov., Mangeliinae, Turridae, genotype Mangelia hilum Hedley p. 25, pl. 5, f. 110–112; A. profundis sp. nov. Mangeliinae 70 fms. off Green Cape, N.S. Wales, p. 25 pl. 5 f. 113. C. F. Laseron Review of the New South Wales Turridae, Sydney, 1954.

Austroturris gen. nov. Turridae genotypes Filodrillia steira Hedley, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 6 pl. 1 f. 134.

Clathurella aguayoi sp. nov. San Matías Gulf, Argentina, A. Carcelles Comun, zool. Mus. Montevideo 4 70 1953: p. 12 pl. 4 f. 20.

†Clavatula interrupta vitálisi nov. f. Miocene, Hungary, L. Strausz Földt. Közl. 85 2 1955: p. 202 pl. XII f. 25, 26; Clavatula, C. R. Ronchetti (3).

Conasprella (?) gratacapi (Pilsbry) figured, Anon.

Conidae—list of Japanese species, T. Kuroda (1).

Conticosta gen. nov. Turrinae genotype Inquisitor petilinus Hedley, C. F.

Laseron Review of the New South Wales Turridae, Sydney 1954 p. 13 pl. 3 f. 58.

Conus litteratus L. notes, Anon.; C. ermineus Born, V. J. Conus lithoglyphus Reeve, [J. Q. Burch] (1); C. drangai p. 13 pl. 2 f.1-7; C. andrangae p. 14 pl. 2 f. 8-11; spp. nov.: C. gradatus thaanumi p. 15 pl. 2 f. 12-13; C. recurnis helenae p. 15 pl. 2 f. 14-15; subspp. nov. from Costa Rica, J. S. Schwengle Nautilus Philad. 69, 1955; C. (Gastridium) eldredi nom. nov. for Conus geographus rosea Swby. = intermedius Rve. non Lamarck 1810. Conch. Ill. f. 33 1833, non *C. roseus* Fischer 1807 and Lamarck 1810, J. P. E. Morrison, J. Wash. Acad. Sci. 45 1 1955, 32; C. intermedius Reeve renamed C. mappa Crosse Rev. Mag. Zool. (2) 10, 200-205, 1858 Crosse's name however is preoccupied by C. mappa Golander Portland Catal. 116 No. 2554, 1786 [vide C. eldrcdi nom. nov., J. P. E. Morrison J. Wash. Acad. Sci. 45 l 1955; C. magintyi sp. nov. dredged 165° off Pensacola N.-W. Florida, H. A. Pilsbry, Nautilus, Philad. 69 2 1955 47 pl. 2 f. 10-11.

†Conus, C. R. Ronchetti (2). †Cythara, C. R. Ronchetti (3). †Daphnella, C. R. Ronchetti (3).

Epidirona nodulosa p. 9 pl. 1 f. 21–23, 50 fms. off Crosody Head; E. carinata 40–50 fms. off Manning River p. 9 pl. 2 f. 24–26; E. tuberculata p. 10 pl. 2 f. 27–28; E. costifera 40–50 fms. off Manning River p. 9 pl. 2, f. 29; E. candida 70–80 fms. off Cape Everard p. 16 pl. 2 f. 30; E. molleri 50 fms. off Crosody Head spp. nov. New South Wales, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Etrema constricta dredged in North Harbour, Port Jackson p. 27 pl. 6 f. 125–126; E. levicosta 30–35 fms. off Port Stephens p. 27 pl. 6 f. 127–128; E. pyramis Black at Shellharbour p. 28 pl. 6 f. 129–130; E. kitcheni 8 fms. Eden p. 28 pl. 6 f. 132–134; spp. nov. E. kitcheni attenuata subsp. nov. 30–35 fms. off Sydney p. 28 pl. 6 f. 135, 136, C. F. Laseron Review of the New South Wales Turridae, Sydney, 1954.

Euguraleus thornleyanus 30–35 fms. off Port Stephens p. 32 pl. 7 f. 156–157;

sp. nov.; E. tasmanicus peronianus subsp. nov. 50 fms. 7 m. N of Montague Island p. 33 p.1 8 f. 160, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Exomilus cylindricus sp. nov. 80 fms. 22 m. east of Narrabeen, N.S.W. Turridae, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 48 pl. 12 f. 253-254.

Filodrillia tricarinata intermissa subsp. nov. Mangeliinae 30-70 fms. off Green Cape, N.S.W. p. 23 pl. 5 f. 94-96; F. delicatula 30-35 fms. off Crookhaven p. 23 p. 5 f. 100; F. ordinata 50-70 fms. off Green Cape p. 24 pl. 5 f. 103; F. thornleyana 30-35 fms. off Port Stephens p. 24 pl. 5 f. 104-105; F. vitrea 70-84 fms. off Cape Everard p. 25 pl. 5 f. 106-107; F. teres 20-25 fms. off Crookhaven p. 25 pl. 5 f. 108-109 spp. nov. Turridae N.S. Wales, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Fusidaphne gen. nov. Daphnellinae genotype F. bullata sp. nov. 10-15 fms. Twofold Bay, Turridae N.S. Wales, C.. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 47 pl. 12 f. 246-247.

†Genota, C. R. Ronchetti (3).

Guraleus (type Mangelia picta Hedley tasmantis 70 fms. off Green Cape p. 30 pl. 7 f. 148–144, f. tasmantis tritomus 40 fms. off the Manning R. p. 30 pl. 2 f. 145–146; sp. & subsp. nov., G. asper. 15 fms. off the Clarence River p. 31 pl. 7 f. 150–151; G. nanus Mamly Beach in shell sand, p. 31 pl. 7 f. 152; G. fossa 30 fms. off Twofold Bay p. 31 pl. 7 f. 153; spp. nov. Turridae N.S.W., C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Hindsiclava gen. nov. Turridae genotype Pleurotoma militaris Hinds in Reeve 1843, Hertlein & Strong Bull. Amer. Mus. nat. Hist. 107 2 1955 p. 227.

Isodaphne gen. nov. Daphnellinae genotype, I. garrardi sp. nov. 10–18 fms. Twofold Bay, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 44 pl. 11 f. 226–227.

Leptoconus milne-edwardsi (Jousseaume) figured, Anon.

Lioglyphostoma armstrongi sp. nov. Guayabo, Chiquito, Panama, Hertlein & Strong Bull. Amer. Mus. nat. Hist. 107 2 1955 p. 230 pl. 3 f. 12.

Liotina saxa sp. nov. 14 fathoms off Long Reef, N.S.W., Australia, C. Laseron Aust. zool. 12 1954: p. 2 f. 2, 2a.

Maoritomella thola pl. 4 f. 73 40-50 fms. off Manning River; M. foliacea pl. 4 f. 74, 40 fms. Twofold Bay: spp. nov. Clarinae N.S. Wales, C. F. Laseron Review of the New South Wales Turridae. Sydney 1954 p. 18.

Marita elongata dredged North Harbour (Port Jackson) p. 38 pl. 9 f. 197-198; M. tumida 30-38 fms. off Port Stephens, p. 39 pl. 9, f. 199, 200: spp. nov. Turridae N.S.W., C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Micantapex profundis sp. nov. 300 fms. off Sydney, C. F. Laseron Review of the New South Wales, Turridae, Sydney 1954 p. 8 pl. 1 f. 13.

Microdiscula vitrea p. 4 f. 8, 8a; M. pellucida p. 5 f. 9, 9a; M. fragilis p. 5 f. 48, 48a, b. spp. nov. N.S.W. Australia, C. Laseron Aust. zool. 12 1954.

Microdrillia stephensensis sp. nov. Clarinae, 30-55 fms. off Port Stephens, New South Wales, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 19 pl. 4 f. 76-78.

Microgenia gen. nov. Daphnellinae genotype Clathurella edwini Brazier, N.S.W. Turridae, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954, p. 50 pl. 12 f. 261, 262.

Mitraguraleus gen. nov. Mangeliinae genotype Bela mitralis Adams & Angas, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 32 pl. 7 f. 154-155.

Narraweena gen. nov. Clavinae genotype Asthenotoma subtilinea Hedley p. 19 pl. 4 f. 79-80; N. carrota sp. nov. 30-35 fms. off Port Stephens, N.S. Wales 19 pl. 4 f. 79-80, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Nepotilla serrata shell sand, Port Stephens, p. 49 pl. 12 f. 256–257; N. carinata 30–35 fms. off Crookhaven p. 49 pl. 12 f. 258-259; spp. nov. N.S.W. Turridae, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Notocytharella gen. nov. Turridae genotype Cytharella niobe Dall, Hertlein & Strong Bull. Amer. mus. nat. Hist. 107 2 1955 p. 232.

Paracuneus gen. nov. Turridae genotype Mangelia immaculata Tenison-Woods: P. immaculata perionianus subsp. nov. Quarantine Bay pl. 3 p. 50-51; P. spadix tumulus subsp. nov. pl. 3 f. 54-55; 50-70 fms. Twofold Bay; P. kemblensis sp. nov. 63-75 fms. off Port Kembla, New South Wales Turridae, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 14.

Paradaphne gen. nov. Daphnellinae genotype Daphnella botanica Hedley p. 44 pl. 11 f. 222-223; P. bucklandi 10-15 fms. Twofold Bay p. 44 pl. 11 f. 224-225 Turridae N.S. Wales, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Paramontana gen. nov. Turridae genotype Clathurella modesta Angas p. 41 pl. 10 f. 210, 211; P. fusca Eden, p. 41 pl. 10 f. 213-214; spp. nov. N.S.W., C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

Peraguraleus emina brevostium subsp. nov. 30-35 fms. off Crookhaven p. 34 pl. 8 f. 163; P. howelli 70-80 fms. off Lake Everard p. 34 pl. 8 f. 164-165; P. subitus 55-70 fms. off Green Cape p. 34 pl. 8 f. 166-168; P. tepidus 30-35 fms. off Port Stephens p. 310 pl. 8 f. 169; P. serpentis dredged in North Harbour, Port Jackson p. 35 pl. 8 f. 172; P. alternatus 40-50 fms. off Manning River p. 35 pl. 8 f. 173-175; P. lucidus 14 fms. off Long Reef p. 35 pl. 8, f. 177-178; spp. nov. Turridae N.S.W., C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

†Pleurotomella, C. R. Ronchetti (3).

Pulsarella gen. nov. Clavinae genotype Pleurotoma cognata Smith, C. F. Laseron Review of the New South Wales Turridae, Sydney, 1954 p. 20 pl. 4 f. 84.

Pseudoraphitoma pyramidula sp. nov. dredged in North Harbour, Port Jackson, N.S.W., C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 43 pl. 10 f. 220–221.

Splendrillia molleri sp. nov. Clavinae 50 fms. off Crowdy Heads, N.S. Wales, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p. 17 pl. 4 f. 7.

Stilla anomala sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15: p. 113 pl. 1, f. 8.

Tasmadaphne gen. nov. Daphnellinae genotype T. spicula sp. nov. 30-35 fms. off Crookhaven, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 pp. 46-47 pl. 11 f. 242-243.

Terebra doello-juradoi sp. nov. Atlantic ocean, Argentina, A. Carcelles Comun. zool. Mus. Montevideo 4 70 1953: p. 14 pl. 4 f. 21.

Terebridae of West Mexico and Panama, J. Q. Burch.

Turcicula Dall on the genus and list of species described as such. figs., H. Rehder (1).

Turrella gen. nov. Mangeliinae genotype Clathurella tenuilirata Angas p. 36 pl. 9 f. 179–182; T. crassa 30–35 fms. off Crookhaven p. 36 p.l p f. 185, 186; T. gracilis 70 fms. off Green Cape p. 37 pl. 9 f. 189–190; T. subcostata Crookhaven Heads, p. 38 f. 191, 192; T. asperrima 6 m. east of Jibbon Point N.S.W. p. 38 pl. 9 f. 193, 194; spp. nov. Turridae of New South Wales, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954.

†Turris (Oxytropa) pseudovolgeri sp. nov. Oligocene, Belgium, M. Glibert Bull Inst. Sci nat. Belg. 31 86 1955: p. 6.

†Turris, C. R. Ronchetti (3).

Vexitomina pilazona 10-15 fms. Twofold Bay p. 12 pl. 2 f. 38, 39; V. garrardi dredged in Twofold Bay p. 13 pl. 2 f. 42, 43; V. torquata 30-50 fms. off Sydney p. 13 pl. 3 f. 48-49; spp. nov. New South Wales, C. F. Laseron Review of the New South Wales Turridae, Sydney 1954. Xenoroturris corona sp. nov. 50-70 fms. off Green Cape, N.S.W., C. F. Laseron Review of the New South Wales Turridae, Sydney 1954 p, 6 pl. 1 f. 1, 2.

Opisthobranchiata

BULLACEA

Acteon finlayi sp. nov. off Palm Beach, Florida, T. L. McGinty Proc. Acad. nat. Sci. Philad. 107 1955 p. 81 pl. 2 f. 10; Acteon tornatilis (L.), Fretter & Graham.

†Actaeon pygmaeus sp. nov. Miocene, Czechoslovakia, J. Senes Geol. Prác. Slovensk. Akad. 40 1955: p. 159 pl. IX f. 17-24.

Akeridae of Japan, T. Habe (1).

Austrodiaphana flemingi sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955; p. 115 pl. 3 f. 29.

Bulla orientalis p. 21 pl. 3 f. 10; B. difficilis p. 22 pl. 3 f. 13; spp. nov. Japan, T. Habe Illust. Catal. Jap. Shells No. 3 1950; Bulla Historical review of the Linnaean genus, H. Dodge.

Bullidae of Japan, T. Habe (1).

Bullina nobilis p. 19, pl. 3 f. 11; B. virgo p. 20 pl. 3. 4; spp. nov. Japan, T. Habe Illust. Catal. Jap. Shells No. 3 1950.

†Cylichna stolida sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig 4 1955: p. 125 pl. 5 f. 11-13.

†Eoacteon gen. nov. Acteonidae, p. 132 type species: Solidus linteus Conrad; E. linteus (Conrad) p. 133 p. 23 f. 8-10. Cretaceous, North America, L. W. Stephenson Prof. Pop. U.S. geol. Surv. 274-E 1955.

Hamineobulla kawamurai gen. et sp. nov. Amami-Oshima, Japan, T. Habe Illust. Catal. Jap. Shells. No. 3 1950 p. 22 pl. 3 f. 9.

Hydatinidae of Japan, T. Habe (1).

†Oligoptycha? popenoei sp. nov. Cretaceous, Mexico, E. C. Allison J. Paleont. 23 3 1955: p. 430 pl. 44 f. 14, 15.

Philinoglossa general anatomy, E. & E. Marcus (2).

Pyrunculus obesus p. 13 pl. 2 f. 12; P. tokyoensis p. 14 pl. 2 f. 20; spp. nov. Japan T. Habe Illust. Catal. Jap. Shells No. 2, 1950.

Retusa elegantissima p. 12 pl. 2 f. 22, 23; R. (Ventomnestia) kawamuriai p. 10 pl. 2 f. 13; spp. nov. Japan, T. Habe Illust. Catal. Jap. Shells No. 2 1950.

Retusidae of Japan, T. Habe.

Ringicula (Ringiculina) takeyamai p. 9 pl. 2 f. 5; R. (Ringiculina) taramachii p. 9 pl. 2 f. 6; R. (Ringiculina) pilula p. 10 pl. 2 f. 11; R. (Ringiculina tosaensis p. 10 pl. 2 f. 2, spp. nov. Japan T. Habe Illust. Catal. Jap. Shells No. 2 1950.

Ringiculidae of Japan, T. Habe.

Scaphander Montfort 1810 Validation, Opinion 287.

Toledonia succineaformis sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 114 pl. 3 f. 30.

APLYSIACEA

Aplysia from South Africa, W. Macnae; Aplysia, egg mass and figs. of animal in motion, J. Bernstein; Aplysia depilans L. aspects of ovogenesis, A. Bolognari; Aplysia vaccaria sp. nov. deep waters off San Pedro and Palos Verdes, Southern California, L. H. Winkler Bull. S. Calif. Acad. Sci. 54 1 1955 p. 5 text-fig.

Chelidonura evelinae sp. nov. São Sebastião Island, Brazil, E. Mareus Bol. Fac. Fil. Cienc. S. Paul (Zool.) 20 1955: p. 95 f. 8-19.

Laplysia [recte Aplysia] fasciata Poiret 1789 addition of specific name to Official List, Opinion 354; Laplysia [recte Aplysia] punctata Cuvier 1803 validation of specific name, Opinion 355.

Notarchus (Stylocheilus) notes on Japanese species, K. Baba.

Phyllaplysia engeli sp. nov. Şão Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 105 f. 29–37.

PTEROPODA

Adula falcatoides sp. nov. p. 4 pl. 7 f. 7 Akkeshi Bay, Japan; A. nipponica sp. nov. Japan = A. californiensis auctt. (not the American species of

this name) p. 5 pl. 3 f. 5 Akita Prefecture, northern Honshu, Japan, T. Habe Publ. Akkeshi mar. biol. Sta. No. 4 1955.

CONULARIACEA

†Charruia annulata n. gen. n. sp. Gastropoda Cambrian, Argentina, C. Rusconi Rev. Mus. Hist. nat. Mendoza 8 1-4 1955: p. 11 pl. 1 f. 14.

†Conularia leonardensis sp. nov. Permian, West Texas, North America, R. M. Finks J. Paleont. 29 5 1955: p. 834 pl. 82 f. 1-13; C. pentagonala sp. nov. Carboniferous, Poland, S. Kwiatowski Ann. Soc. géol. Pologne 23 (1953) 1955: p. 158 pl. xi.

†Hyolithellus quadrangularis sp. nov. Cambrian, Argentina, C. Rusconi Rev. Mus. Hist. nat. Mendoza 8 1-4 1955 : p. 12 pl. 1 f. 11.

†Tentaculites mogotensis sp. nov. Ordovician, Argentina, C. Rusconi Rev. Mus. Hist. nat. Mendoza 8 1-4 1955: p. 44 pl. 5 f. 6.

Nudibranchiata

SACOGLOSSA

ELYSIACEA

Alderia modesta range extended to Washington, C. Hand; A. modesta (Lovén) occurrence on the Central California coast, figs., Hand & Steinberg.

Elysia canguzua p. 111 f. 45–48, 60–45; E. serca p. 113 f. 49–52, 59; E. chitwa p. 115 f. 53–56, 58 spp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955; E. elsieae p. 122 f. 9 pl. 1.; E. degeneri p. 124 f. 10, pl. 1—E. nealae p. 125 f. 11 pl. 1 spp. nov. Hawaii, J. M. Ostergaard Pacif. Sci. 9 2 1955.

Hermaea coirala sp. nov. São Se bastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 108 f. 38-44.

Limapontia depressa pellucida Kevan in Germany, figs., Jaeckel (5).

DORIDACEA

Archidoris montereyensis reproductive system and embryology, McGowan & Pratt.

Awuka gen. nov. p. 155 Centrodoridinae: Genotype: Awuka spazzola sp. nov. p. 156 f. 180-192 São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (zool.) 20 1955.

Cadlina rumia sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 119 f. 78-89.

Chromodoris macfarlandi emendation, Opinion 221.

Corambella carambola sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 163 f. 209-217.

Discodoris evelinae p. 143 f. 141-150; D. pusae p. 147 f. 151-165 spp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955.

Doridopsis macfarlandi sp. nov. Hawaii, J. M. Ostergaard Pacif. Sci. 9 2 1955: p. 128 f. 13.

Doris bovena sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 131 f. 109-118.

Doris (?) ignota nom. nov. p. 285 for "D. grandiflora Rapp" Cantraine, 1840, non Pease, 1861, non Rapp, 1827; D. (?) dalyelli nom. nov. p. 289, for D. argo Dalyell, 1853 p.p., non Linné, non tuberculata Cuvier, A. Pruvot-Fol Faune de France 58 1954.

Glossodoris neona sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 124 f. 90-101.

Goniodoris minula sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 161 f. 203-208.

Hexabranchus tinkeri p. 128 f. 14 pl. 2; H. aureomarginatus p. 132 f. 15, pl. 2 spp. nov. Hawaii, J. M. Ostergaard Pacif. Sci. 9 2 1955.

Peltodoris greeleyi MacFarl., description and figures, E. Marcus.

Polycera odhneri sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Dac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 155 f. 193–202.

† Pelagiella staufferi p. 14, pl. 16 f. 1, 2, Kasota sandstone, Ordovician, Minnesota, U.S.A., L. Powell Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935. Siraius gen. nov. p. 134 Dorididae, genotype: Siraius ilo sp. nov. p. 135 f. 119-125 São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955.

Taringa gen. nov. p. 151 Discordoridinae: Genotype: Taringa telopia sp. nov. p. 152 f. 166–179 São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955.

Thordisa diuda sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 140 f. 133-140.

Tritoniopsis aurantia sp. nov. near Santa Catalina Island, S. California, N. T. Mattox Bull. S. Calif. Acad. Sci. 54 1 1955 p. 8 pls. 4 and 5.

AEOLIDIACEA

Aeolidia edmonsoni sp. nov. Hawaii, J. M. Ostergaard Pacif. Sci. 9 2 1955: p. 133 f. 16, pl. 1.

Doto uva p. 167 f. 218–222; D. pita p. 169 f. 223–225 sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955.

Favorinus auritulus sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 181 f. 259-26.

Janolus comis sp. nov. São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955: p. 170 f. 226–236.

Microhedyle general anatomy, E. Marcus.

Piseinotecus gen. nov. p. 176 Cuthonidae: Genotype: Piseinotecus divae sp. nov. p. 177 f. 244–248 São Sebastião Island, Brazil, E. Marcus Bol. Fac. Fil. Cienc. S. Paulo (Zool.) 20 1955.

Pseudovermis schulzi p. 235 pl. 36 f. 9-4 Arcachon; P. axi p. 236 pl. 37 f. 15-22 Banyuls-sur-Mer spp. nov. France, E. & E. Marcus Kieler Meeresforsch. 11 1955.

RHODOPACEA

Rhodope veranyi general anatomy, E. & E. Marcus.

Pulmonata ELLOBIACEA

Maripythia gen. nov. Ellobiidae for xanthostoma Iredale Rec. Austr. Mus. 19 1936 in text p. 328.

Ovatella myosotis Drap. anatomy, radula fig., K. O. Meyer.

SIPHONARIACEA

Kerguelenella flemingi sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 121 pl. 5 f. 44.

LYMNAEACEA

Amerianna pesigani sp. nov. Philippines, B. Hubendick Philipp. J. Sci. 83 3 1954: p. 319 pl. 1 f. 1-8.

Ancylastrum addition of name to official list with type Ancylus (Ancylastrum) cumingianus Bourgt. 1853, Opinion 364.

Ancylus, On type species of Ancylus Müller 1774, Opinion 363.

Anisancylus lagunarum sp. nov. Laguna Jejane, F. Haas Tr. Linn. Soc. London (3) 1 1955 p. 284 f. 7a-c.; A. crequii (Bavay), in Lake Langui, F. Haas (6).

Australorbis glabratus temperature effects on tissue hydration, T. v. Brand.

Biomphalaria elegans sp. nov. Lake Albert, Uganda, Mandahl-Barth Ann. Mus. Congo belg. Svo 32 1954: p. 82 f. 36c; B. boissyi anatomy relations to infection with Schistosoma, E. T. Abdel-Malek; Biomphalaria relations to other Planorbidae, H. Watson (2).

Bulimus corneus Lea, B. columbianus Lea, B. lacteus Lea, B. pralianus Lea, holotypdes fig., R. T. Abbott.

Bulinus reticulatus sp. nov. p. 109 f. 51—B. globosus ugandae subsp. nov. p. 114 f. 55 a—h Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

Camptoceras hirasei Walker anatomy figs. and notes on the genus, B. Hubendick.

Carinorbis subgen. nov. of Segmentorbis Planorbidae. Subgenotype: Segmentorbis (Carinorbis) kanisaensis (Preston). From the Nile at Kanisa, immediately north of Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

Choanomphalus anatomy, B. Hubendick.

Ferrissia parallelus North American snail found in Germany, S. Jaeckel (2); F. toroensis p. 119 f. 59a; F. kavirondica p. 120 f. 59e spp. nov. Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

Glyptanisus ordessus [? gen. or subgen. nov.] sp. nov. Planorbidae, Iredale in text p. 224.

Gundlachia general morphology, M. I. Scott (7).

Gyraulus costulatus subtilis p. 87 f. 39b—G. c. exilis p. 88 f. 39c; G. kigeziensis nyanzae p. 91 (foot-note) subspp. nov.; G. concavus p. 89 f. 40c; G. crassus p. 90 f. 40b; G. bicarinatus p. 91 f. 41b spp. nov. Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

Hippeutis riparius occurrence, ∇ . Lozek.

Lentorbis gen. nov. p. 92 Planorbidae, sub-family Planorbinae. Genotype: L. benguelensis (Dunker) from Angola and Lower Congo. Genus comprises 2 species—L. junodi (Connolly, 1922)—hitherto referred to Hippeutis Charpentier, 1837, Mandahl-Barth Ann. Mus. Congo belge 8vo 32 1954.

Limnaea plicata sp. nov. Argentina, M. I. H. Scott Physis 20 59 1954: p. 406 f. 4-6.

Littoridina stiphra Lake Titicaca at Molinopampa p. 285 f. 8; L. lacustris Paton, Lake Titicaca p. 287 f. 10; L. languiensis Langua, L. Titicaca p. 287 f. 11; L. profunda Taman, Lake Titicaca p. 289 f. 13; L. saracockae Laguinilla Saracockae p. 289 f. 14; L. vestita Lagunilla Saracockae p. 290 f. 15; L. aperta Molinopampa Bay p. 296 f. 21; spp. nov. Lake Titicaca, F. Haas Trans. Lim. Soc. London (3) 1 1955.

Lymnaea (Pseudosuccinea) columella Say in the Basle Botanic Gardens, fig., Bruggen (1); Lymnaea stagnalis L. physiology etc., Horstmann.

Oppletora name in text (Physopsislike jukesii, H. & A. Ad. in text under Leishkardtian theocifaunula p. 54. Physa fontinalis (L.) its life cycle and other biological details graphs and tables, W. F. de Wit (1); Physa gyrina Say, intrinsic rate and natural increase R. M. DeWitt.

†Physa skinneri sp. nov. Pleistocene, Berends fauna, Beaver Co., Oklahoma, U.S.A., D. Taylor Occ. Pap. Mus. zool. Univ. Mich. 557 1954: p. 4, p. 9.

Planorbarius corneus and its range in the British Isles, K. F. McMillan.

Planorbidae, Parasitism, E. Dias; Planorbidae, nomenclature of African forms, J. Schwetz; Planorbidae, Phylogeny, figs., B. Hubendick.

Planorbis metidjensis—radula and genitalia, Fraga de Azevedo & Medeiros; P. bosqueti (G. Vincent mss.) nov. nom. = P. depressus Nyst, 1843; non Faure et Sionest, 1808; = P. lovanensis mss. in Glibert, M. and de Heinzelin, J. 1952, p. 12 (non. G. Vincent, mss.), Glibert & Heirzelin Bull. Mus. Hist. nat. Belg. 30 1 1954: p. 10 pl. 2 f. 8; P. (Gyraulus) laevis Alder in Dulwich Park and associated species, M. P. Kerney.

†Planorbis (Anisus) brunnensis p. 53 pl. 2 f. 3a-e; P. (Gyraulus) cryptornatus p. 56 pl. 7 f. 4a-d; P. (G.) götzendorfensis p. 56 pl. 7 f. 3a-c—P. (G.) rotundistomus p. 58 pl. 6 f. 3a-c; P. (G.) edlaurei p. 58 pl. 6 f. 4a-c; P. (G.) trolli p. 58 pl. 7 f. la-c; P. (G.) pseudhomalosomus p. 58 pl. 9 f. 4 4a-e; P. (G.) symmetricus p. 60 pl. 1 f. 2a-c; P. (G.) primiformis p. 62 pl. 10 f. la-c; P. (G.) seibersdorfensis p. 62 pl. 10 f. 3a-c; P. (G.) zemendorfensis p. 64 pl. 10 f. 2a-c; P. (Bathyomphalis) mödlingensis p. 66 pl. 4 f. 4 a-c spp. nov.; P. (Odontogyrorbis) krambergeri angustigyratus p. 54 pl. 3 f. 2a-e; P. (Gyraulus) verticillus planitesta p. 58 pl. 6 f. 2a-c; P. (G.) turislavicus posterior p. 60 pl. 9 f. 2a-c; P. (G.) popovici complanatus p. 61 pl. 3 f. 4a-e; P. (G.) globulus coristomus p. 61 pl. 8 f. 2a-c; P. (G.) globosus globosus p. 62 pl. 8 f. 1a-e; P. (G.) fuchi asymmetricus pl. 8 f. 4a-c; P. (Segmentina) loczy anterior p. 66 pl. 4 f. 3a-c subspp. nov. Pannon (Tertiary), eastern Europe, F. Sauerzopf Burgenland Heimat. Bl. 15 2 1953; P. (Gyraulus) cordatus vinciennei var. nov. p. 539 pl. xxiii b, f. 3, 4.—Planorbis (Paraspira) elegans

minor [var.] nov. p. 540 pl. xxiii b f. 2, Oligocene, Jura, P. Jodot Bull Soc. géol. Fr. 6 4 7-9 1954.

Plananisus for series of Planorbidae [? gen. or subgen. nov.] typified by tasmanicus Ten.-Woods in text p. 224, Iredale.

Pygmanicus [? gen. or subgen. nov. for scottiana Johnston, in text p. 224, Iredale.

Segmentorbis gen. nov. p. 94 Planorbidae, sub-family Planorbinae. New genus for the African species previously referred to Segmentina Fleming, 1922—Segmentorbis (Carinorbis) subgen. nov. genotype S. (C.) kanisaënsis (Preston) p. 58 f. 45 Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

Stagnicola reflexa studies on Toxicity, and Mollucicide, R. A. Wood.

Taphius spp. in Lake Titicaca, F. Haas (6).

Tropicorbis nordestensis sp. nov. Pernambuco N.-E. Brazil, D. T. de Lucene J. Conchyliol. Paris 95 1 1955 p. 20 f. la-c.

VAGINULACEA

Veronicella (sens lat.) tolana sp. nov. Mt. Tola, Gughé Highlands, southern Ethiopia, H. Quick J. Linn. Soc. (Zool.) 42 1954: p. 385 f. 12-17.

Veronicellidae from West Africa, L. Forcart (1).

SUCCINEACEA

Austrosuccinea archeyi from Mt. Maunganui, New Zealand, W. Thomson (1); Austrosuccinea archeyi further records for New Zealand, W. Thomson & A. Jones.

Oxyloma verrili (Bland) note and fig. of radula, Venmans.

Succinea arenaria Beach-Chant and S. oblonga Drap. in Belgium, figs., A. Anteunis; Succinea, studies on chromosome numbers, figs., Kayama; Succinea hirasei—S. kwansae—S. horticola chromosome number of the Japanese species, T. Koyama; S. avara (Say) notes on the species, L. Venmans (1).

ATHORACOPHORACEA

Pseudaneitea sorenseni sp. nov. New Zealand A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 128 pl. 4 f. 40.

ACHATINELLACEA

Achatinella viridans Mighels Lectotype fig., R. I. Johnson.

Lamellidea solomonensis sp. nov. Solomon Islands, R. K. Dell Pacif. Sci. 9 3 1955: p. 357 f. l.

VERTIGINACEA

†Acanthinula lamellata (Jeff.) in Holocene tufa at Totland, Isle of Wight, A. G. Davis.

Albinaria (Albinaria) brevicollis theodori subsp. nov. S. Theodoro Is. p. 120. pl. 8 f. 9; A. (A.) b. castellensis C. Capria W. Coast Rhodes p. 125 pl. 9 f. 14-15; A. (A.) b. telensis Piscopi Id. p. 139 pl. 11, f. 11-32; A. (A.) b. simicensis Id. of Simi p. 140 pl. 11 f. 33-34; A. (A.) maculata kosensis Rastel Kos, p. 150, pl. 11 f. 37-38; subspp. nov., K. L. Pfeiffer Arch. Mollusk. 84 4/6 1955.

Charpentieria (Dilateria) marcki pumila subsp. nov. Balkan peninsula, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 60 f. 5.

Clausilia dubia in the Malham area, Yorkshire, L. W. Stratton.

Clausiidae of Estonia, C. Krausp (1).

Gastrocopta additional localities of species in Oklahoma, D. S. Dundee; Gastrocopta crucifera p. 245 f. 1; G. pulvinata p. 246 f. 2-3 spp. nov. Northwest Argentina, M. Scott Acta zool. Lilloana 6 1948.

†Gastrocopta (s.s.) lavernensis p. 11 Pliocene, Laverne formation, Beaver Co., Oklahoma—G. (s.s.) chauliodonta p. 12 Pleistocene, late Nebraskan age, Brown Co., Nebraska, spp. nov. U.S.A., D. Taylor Occ. Pap. Mus. zool. Univ. Mich. 557, 1954.

Gyrodaria gen. nov. Gastrocoptidae, for the strangeana series, Iredale Aust. Naturalist 10 1940 p. 234.

Herilla excedens minor subsp. nov. Balkans, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 61 f. 6.

Iphigena riloensis pygmaea subsp. nov. North-west Macedonia, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 58 f. 3.

Laciniaria (Strigilecula) vetusta gracilior subsp. nov. Macedonia, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 59 f. 4; L. (Denticularia) denticulata sprattiana nom. nov. = Clausilia denticulata spratti Boettger 1883 non Pfeiffer 1846, K. L. Pfeiffer Arch. Mollusk. 84 4/6 1955 p. 153.

Lauria anglica at Meathop Fell, Westmorland, G. W. Pitchford.

Macedonia macedonia pirinensis p. 63 f. 8; M. (Serbia) frauenfelli occidentalis p. 64 f. 11, 12 subspp. nov. Macedonia, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954.

Nenia (Neniatracta) adusta tumens road between Huancayo and Mejorada, Huancavelica, Peru p. 305 f. 57: N. (?) angrandi kalinowskii Polanco, Tambo, San Miguel, Agauscho, Peru, p. 306 f. 58, F. Haas Fieldiana Zool. 37 1955; Nenia argentina sp. nov. Las Capillas, Jujuy, Argentina, M. I. Scott Neotropica 1 No. 1 1954: p. 2 f. 1-7.

Papualbinula subgen. from Australbinula macrodon Pils., Iredale Aust. Zool. 10 1 1941, in text p. 63.

Pupillidae from Northern Venezuela, S. Arias; Pupillidae from Estonia, C. Krausp.

Truncatellina cylindrica britannica Pils. in Dorset and the Isle of Wight, A. G. Davis.

Valloniidae of Estonia, C. Krausp (2).

Vertigo parcidentata (Braun) its former occurrence in Hertfordshire, M. P. Kerney; V. moulinsiana in Brandenburg, T. Schmierer (1); V. ronnebyensis in Berlin, F. Zimmerman; V. arctica figs. of shells, vicinity of Stockholm, H. W. Walden.

ACHATINACEA

Achatina fulica Bowdich, and its control by Gonaxis kebweziensis E. A. Smith, A. R. Mead.

Acicula Hartmann 1821 addition to official list, Opinion 344.

Cocklicopa figs. of shells in the vicinity of Stockholm, H. W. Walden.

Megaspira ruschenbergiana Lea, holotype figured, R. T. Abbott.

ZONITACEA

Aegopis verticillus ljubetinensis subsp. nov. Ljubotin, Balkans (Macedonia), S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 66 f. 13. Agriolimax gughensis p. 382 f. 1-5; A. scotti p. 386 f. 6-11 spp. nov. Gughé Highlands, southern Ethiopia, H. Quick J. Linn. Soc. (Zool.) 42 1954.

Allodiscus austrodimorphus p. 1137 Text-fig. 1 f. 1; A. fectoloides p. 1139. Text-fig. 1 f. 2, 5 spp. nov. New Zealand, R. K. Dell Trans. roy. Soc. N.Z. 82 5 1955.

Arion lusitanicus in France, C. v. Regteren Altena (1); A. lusitanicus Mabille its occurrence in France, anatomy, figs., C. O. v. R. Altena & others; A. rufus L. experiment on regeneration, P. Laviolette; Arion and other British slugs, J. N. Savory.

Charopidae—studies on the Australian forms, R. C. Kershaw (2).

Daudebardia fallax sp. nov. Lillafüred, Hungary, L. Soós Acta Zool., Budapest, 1 1955 pp. 358-359 f. 5-8; D. rufa hydrophila var. nov. Hungary, L. Soós Acta Zool. Budapest 1 1955 p. 355; D. brevipes occurrence W. R. Schlichum.

Deroceras reticulatum (Muller) Spermatogenesis and effect of light, D. & N. E. Pelluet.

Discus (Gonyodiscus) brunsoni sp., nov. north side of McDonald Lake, Mission Range, Mission Range Lake County, Montana, S. S. Berry Bull. S. Calif. Acad. Sci. 54 1 1955 p. 17 pl. 7; D. macclintocki, F. C. Baker in Bexby State Park, Iowa, L. Hubricht (6).

Drepanostomella circumscripta p. 262 f. 10, 10a; D. tucma p. 264 f. 11 spp. nov. Northwest Argentina, M. Scott Acta zool. Lilloana 6 1948.

Entodina lundi (Mörch, 1871) "comb. n." [Helix (Polygyra) lundii Mörch, 1871; H. lundii Pfeiffer, 1876; H. lundii Tryon, 1887], H. de S. Lopes (1).

Flammulina lateaperta sp. nov. New Zealand, R. K. Dell Trans. roy. Soc. N.Z. 82 5 1955 p. 1140 Text-f. 1 f. 3, 6, 10, 15; F. sorenseni sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 122 pl. 4 f. 38, 39.

Gigantomilax (Turcomilax) iliensis sp. nov. Kouldja, China, C. van. R. Altena Beaufortia 4 36 1954: p. 47 f. 1-3.

Gyropena gen. nov. of Charopidae genotype G. verans sp. nov. Lord Howe Id., Iredale p. 321. Aust. Zool. 10 1944. Huonodon gen. nov. type P[tychodon] psiertes Charopidae, Iredale Aust. Zool. 11 1 1945, in text f. 56 under Norfolcioconcha.

Insullaoma subgen. nov. for Paralaoma riddlei, Iredale S. Aust. Natst. 18 1937, in text p. 19.

Kermodon gen. nov. for Ptychodon royanus Charopidae, Iredale Aust. Zool. 11 l 1945, in text p. 56 under Norfolcioconcha.

Limax carbonarius albanicus subsp. nov. Berisha, Albania, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 68 f. 14, 15.

Magnipelta mycophaga Pils. (see Nautilus 67: 37 pl. 5 f. 1-3] further notes and figures now given, Pilsbry & Brunson Notul. Nat. Acad. Philad. No. 202 pp. 1-6, figs.

Nipponarion gen. nov. Arionidae genotype N. carinatus sp. nov. Hisamatsu-yama, Tottori City, Japan, Yamagushi & Habe Venus 18 4 1955 235 237 f. 1, 3, 6-9.

Nipponolimax gen. nov. Philomycidae genotype N. monticola sp. nov. Amabiki patti, Takigan, Hyogo Pref. Japan, Yamaguchi & Habe Venus 18 4 1955. 235 238 f. 2, 4, 5,

Oxychilus tasmanicus sp. nov. Tasmania, C. McLauchlan Aust. zool. 12 1954: p. 40 f. 1-4; Oxychilus draparnaldi occurrence, N. B. Schmidt.

Pallifera fosteri, F. C. Baker extension of range to north-eastern Ohio, R. W. Dexter.

Penescosta gen. nov. for Charopa mathewsi Preston, Iredale Aust. Zool., Vol. 10 1944, in text p. 320 under genus Charopella nov.

Phrixgnathus viridula caswelli subsp. nov. New Zealand, R. K. Dell Trans. roy. Soc. N.Z. 82 5 1955: p. 1145 Text-f. 1 f. 11; Phrixgnathus (Taguahelix) hirsutus p. 126 pl. 4 f. 32; P. (T.) delicatulus p. 127 pl. 4 f. 33-35; P. (T.) antipodensis p. 127 pl. 4 f. 36, 37 spp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955.

Polygyra additional localities for species in Oklahoma, D. S. Dundee.

Ptychodon gadus obscurus p. 1141 Text-fig. 1 f. 7; P. monoplax takahea p. 1142 subspp. nov.; P. blacki p. 1142 Text-f. 1 f. 9, 12, 13; P. fiordlandica p. 1143 Text-f. 1 f. 4, 14 spp. nov. New Zealand, R. K. Dell Trans. roy. Soc. N.Z. 82 5 1955.

Radiodiscus katiae sp. nov. Northwest Argentina, M. Scott Acta zool. Lilloana 6 1948: p. 253 f. 5.

Semilaoma subgen. nov. with P[archaoma] abjeta as type, Iredale Aust. Zool. 10 3 1944, in text p. 311 under Paralaoma lidgbirdensis.

Serpho matthewsi in New Zealand, N. W. Gardner (2).

Slugs of Japan, Gamaguchi & Habe; Slug, effect of light; speed and movement, M. J. Brown.

Stephanoda jujuyensis sp. nov. Northwest Argentina, M. Scott Acta zool. Lilloana 6 1948: p. 251 f. 4.

Systrophia stenogyra zischkai subsp. nov. Bolivia, W. Blume Arch. Mollusk. 84 4/6 1955 179, fig. 1; S. (Systrophia) siolii sp. nov. Serra Formosa Melata, Brazil, F. Haas Arch. Mollusk. 84 1/3 1955 103 f. 4-6.

Taguahelix subgen. nov. of Phrixgnathus Charopidae: type: Thermia expeditionis Suter (see P. (T.) campbellicus (Filhol)) New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 125.

Triodopsis cragini Call. 1886, type locality, E. J. & L. D. Roscoe.

Vitrina pellucida islandica subsp. nov. N-W-Island Hafjordan, L. Forcart Arch. Mollusk. 84 4/6 1955, 156, 160 pl. 12 f. 3, 6; Vitrina kochi Andreae and V. nivalis Dumont and Mortillet, a revision, L. Forcart; Vitrina East African forms and taxonomy, B. Hubendick (4).

Zonitoides (Zonitellus) arboreus (Say) in the Botanic Gardens, Basle, figs., **Bruggen** (1).

ARIOPHANTACEA

Alienitor lyndhurstoides sp. nov. Sydney, N.S.W. Australia, C. McLauchlan Aust. 2001. 12 1954: p. 40 f. 1-4.

Buffetia subgen. nov. for Allenoconcha retinacutum Preston, Iredale Aust. Zoologist 11 1945, in text p. 60 pl. iii f. 18-20.

Guppya lilloana p. 267 f. 12; G. aenea p. 270 f. 13 spp. nov. Northwest Argentina, M. Scott Acta zool. lilloana 6 1948.

Kermarion gen. nov. from H. kermadecensis Smith, Iredale, in text p. 329 under Howarion gen. nov.

Mussomula gen. nov. Thalassia delta Pfeiffer P.L.S. N.S.W. (3) vi 1891 p. 553, Iredale Aust. Zool. 9 1 1937, in text p. 13 under Sodaleta scandans Cox.

Modonitor subgen. nov. for Nitor puribundus Cox, Iredale Aust. Zoologist 9 1937, in text p. 3 under the species above named.

Orpiella (Owaraha) nissani sp. nov. Solomon Islands, R. K. Dell Pacif. Sci. 9 3 1955 : p. 326 f. la-e.

Parmavitrina subgen. for Helicarion planilabris Cox, Iredale in text p. 8.

Pravonitor subgen. proposed for Nitor kreffti insertaram subsp. nov. Iredale in text p. 3 under Nitor kreffti Cox. Aust. Zool. 9 1937.

Peloparion subgen. nov. for Helicarion hyalinus Pfr., Iredale, in text p. 8.

Videnoida subgen. nov. of Videna for the Japanese species of V. (V.) gouldiana (Pilsbry) Japan, T. Habe Venus 18 4 1955: p. 231 [English p. 234] f. 2.

ACAVACEA

Caryodes dufresnii extra, C. d. superior, C. d. dextra subspp. nov., Iredale Aust. Zool. 9 1 1937 p. 15.

Gyropena nissani sp. nov. Solomon Islands, R. K. Dell Pacif. Sci. 9 3 1955; p. 328 f. 1e-q.

Strophocheilus (Microborus) tenuis sp nov. Peru, F. Haas Fieldiana Zool. 37 1955 p. 330 f. 70.

BULIMULACEA

Austroborus nom. nov. for Microborus Pilsbry, 1926, subgen. of Strophocheilus, J. J. Parodiz Physis 20 57 1949: p. 189.

Bostryx (Platybostryx) doelloi sp. nov. Argentina, M. I. Scott Physis 20 59 1954: p. 142 f. 1-4.

Bulimulus (Scutalus) haenkei p. 372 f. 76 Alto Plano, Bolivia; B. (Lissoacme) extraneus p. 382 f. 82 Chimantamassif, Venezuela spp. nov., F. Haas Fieldiana, zool. 34 1955; B. (Scutalus) phaeocheilus sp. nov. Peru, F. Haas Fieldiana Zool. 37 1955 p. 334 f. 74; B. (Lissoacme) rudistriatus p. 318 f;

64; B. (Peronaeus) pyrgidium p. 321 f. 66; B. (P.) extensus p. 323 f. 67; B. (P.) acme p. 325 f. 68; spp. nov. Peru F. Haas Fieldiana Zool. 37 1955; B. (Scutalus) longitudinalis sp. nov. Ayacucho, Peru, F. Haas, Fieldiana Zool. 37 1955 p. 316 f. 63; B. sporadicus gracilis subsp. nov. Cerro Colorado, Argentina, M. Scott Acta Zool. Lilloana 6 1948: p. 238 pl. 2 f. 3.

Cyclodontia (Spixia) pyriformis (Pilsbry) p. 230. The anatomy; C. (S) dubia p. 235 pl. 1 f. 9, 10 sp. nov.; C. (Plagiodontes) multiplicatus crassus p. 236 pl. 2 f. 1, 2 subsp. nov. Cerro Colorado, Argentina, M. Scott Act. Zool. Lilloana 6 1948.

Drymaeus (Drymaeus) semistriatus p. 374 f. 77 Fazenda Ipanema, São Paulo, Brazil; D. (D.) griffini p. 383 f. 83 Chimantá massif, Venezuela, F. Haas Fieldiana, zool. 34 1955; D. (Drymaeus) eucosmatus p. 331 f. 71-72; D. (D.) zilchi p. 333 f. 73 spp. nov. Peru, F. Haas Fieldiana Zool. 37 1955; D. (Drymanus) schmidti sp. nov. Cozco, Peru, F. Haas Fieldiana Zool. 37 1955 314 f. 62; D. droueti deletus subsp. nov. p. 13 pl. 1 f. 12.; D. neaxanus sp. nov. p. 14 pl. 1 f. 9: Mexico A. Solem Occ. Pprs. Mus. Zool. Univ. Mich. 566 1955.

Liguus, additional names to Catalogue by Clench (Occ. Pprs. No. 10), W. J. Clench (2); Liguus catalogue of Cuban species, M. Jaume; Liguus fasciatus nodali p. 1375; L. f. goenagai p. 1376; L. f. guajaibonense p. 1377; L. f. guirensis p. 1378; L. f. pridai p. 1379; L. f. caibarienense p. 1381; L. f. fuentesi p. 1384; L. f. jibacoaense p. 1387; L. f. ramsdeni p. 1386; L. f. cinnamomum p. 1387 all from Cuba, West Indies; L. f. caballosense p. 1381; L. f. pharius p. 1384 from Isla de Pinos, Cuba, West Indies; L. blainianus guanensis p. 1382, Cuba, W.I. subspp. nov., M. L. Jaume Circ. Mus. biblio. zool. Habana, 1954.

Neopetraeus platycheilus sp. nov. Aparimac, Peru, F. Haas Fieldiana Zool. 37 1955 p. 311 f. 60, 61; N. stelzneri apertus subsp. nov. Cerro Colorado, Argentina, M. Scott Acta zool. Lilloano 6 1948: p. 238 pl. 2 f. 4, 5.

†Palaeobulimulus gen. nov. Bulimulidae genotype Palaeobulimulus eoce-

nicus sp. nov. Eccene, Patagonia, J. J. Parodiz Physis 20 57 1949: p. 177 f. 2.

Partula cedista f. 2A, p. 1; P. candida p. 4 f. 2 B; P. remota p. 5 f. 2 C, D; P. levilineata p. 6 f. 2E; P. levistriata p. 7 f. 2F; P. cuneata p. 8 g. 3A; P, dolichostoma p. 11 f. 3C; P. leptochila p. 10 f. 3B; P. dolichostoma p. 11 f. 3C; P. protracta p. 12 f. 3D; P. atilis p. 13 f. 3E, F; spp. nov. Raiatea, Society Islands, H. E. Crampton Amer. Mus. Novit. No. 1761 1956.

Peronaeus (Lissoacme) birabeni sp. nov. North-west Argentina, M. Scott Acta zool. Lilloana 6 1948: p. 272, pl. f. 4, 10.

Placostylus strattoni New Caledonia sp. nov. holotype f. 9 T. Pain J. Conchyliol. Paris 95 1 1955 p. 18; Placostylus fibratus powelli p. 12, subsp. nov. holotype fig. 2 also sub-fossil; P. fibratus mosesi p. 14 subsp. nov. holotype f. 4; P. scarabus verdis p. 14 subsp. nov. holotype f. 5. New Caledonia, T. Pain J. Conchyliol. Paris, 95 1 1955; Placostylus fibratus pecularius Kobelt, P. f. infundibulum, figd., T. Pain (1).

Plecocheilus (Eurytus) juliani p. 375 f. 78; P. (E.) steyermarki p. 377 f. 79; P. (E.) mundi-perditi p. 378 f. 80 spp. nov.; P. (Plecocheilus) fulminans alticola p. 381 f. 81 subsp. nov. Chimantámassif, Venezuela; P. (Eurytus) tatei p. 385 f. 84 sp. nov. Mount Duida, Venezuela, F. Haas Fieldiana, zool. 34 1955; P. (Eurytus) ortizianus sp. nov. Chancay, Peru, F. Haas Fieldiana, zool. 34 1955: p. 366 f. 73.

Plectostylus argentinensis sp. nov. Argentina, J. Parodiz Physis 20 58 1951: p. 334 1 fig.

Strephocheilus (Austroborus) nom nov. for Strophocheilus (Microborus) Pilsbry, 1926, J. J. Parodiz Physis 20 57 1949: p. 189.

†Strophocheilus (Megalobulimus) avus sp. nov. Tertiary?, Argentina, J. J. Parodiz Physis 20 57 1949: p. 181 f. 1b.

Thaumastus (Scholvienia) schmidti sp. nov. Hacieuda Piso, Locroja Huancavelica Peru, F. Haas Fieldiana Zool. 37 1955 p. 309 f. 59. †Urocoptis antiquus sp. nov. Miocene Dominica, Pilsbry & Olsson Notul. Nat. Acad. Philad. No. 266 1954 p. 4 f. 2.

HELICACEA

Amphidromus (Syndromus) contrarius jaeckeli nom. nov. for A. contrarius floresi Haltenorth & Jaeckel 1940, L. J. M. Butot Penggemar Alam 34 3-4 1954: p. 106.

Amphidromus banksi sp. nov. Mt. Kadam region Pulan Panaitan, L. J. M. Butot Treubia 23 1955 p. 127 f. 29a-e, pl. v.

Amphidromus webbi babiensis subsp. nov. Pecloc Babi, Sumatra p. 76 f. 1; A. (Syndromus) sumatramus jacobsoni forma nov. pp. 77, 80 f. 2; A. webbi simalurensis subsp. nov. Simatang p. 78; A. enganoensis frunstorferi nom nov. for A. e. gracilior Fruhstorfer 1905 now Fulton 1896 p. 80; F. F. Laidlaw Zool. Meded. 33 1954.

Baccalena subgen. nom. for Pleuroxia squamulosa (Yates), Iredale S. Aust. Natst. 18 1937 p. 50 in text.

Bradybaena similaris breeding experiments and populations genetics figs. and tables, Komai & Emura.

Cepaea distribution and variation, C. Boettger (3).

Camaenidae of America systematics and anatomical study, figures, C. B. Wurtz.

†Cepolis (Coryda) propappus p. 2 f. 1, 1a; C. (Coryda) miocenica p. 3 f. 3, 3a spp. nov. Miocene Republic of Dominica, Pilsbry & Olsson Notul. Nat. Acad. Philad. No. 266 1954.

Dirutrachia subgen. nov. for Semotrachia sublevata, Iredale S. Aust. Natst. 18 1937, in text p. 36.

Disteustoma gen. nov. type Helix dinodeomorpha Tap-Can, Iredale Aust. Zool. 10 1941 p. 85.

Eobania vermiculata biological influences on the postembryonic development, E. Frömming.

Eobania vermiculata insular dwarf population on islands in Salerno Bay, C. F. Sacchi (3).

Epiphragmophora puella p. 252 f. 1, 2 31; E. semiclausa p. 257 f. 32; E. proseni p. 258 f. 33 spp. nov. Argentina, M. I. H. Scott Physis 20 58 1951.

Euhadra herklotsi hesperia Pils., with abnormal tentacles, Kishimoto.

Fatulabia subgen. nov. for Pleuroxia elderi, Iredale S. Aust. Natst. 18 1937, in text p. 51.

Findomelon subgen, nov. for Merocomelon interfusca Cox, Iredale S. Aust. Natst. 18 1937, in text p. 32.

Helicella eriaetorum, H. obvia. Distribution and Variation, H. Melchinger; H. (Candidula?) renschi sp. nov. Pirin-Gebirge, Macedonia, p. 70 f. 16—H. candicans L. Pfeiffer (= obvia Hartmann) p. 73, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954; H. delermai sp. nov. Caprara, Tremiti islands, Adriatic, C. Sacchi Boll. Zool. 22 1955: p. 99 pl. 1 f. 7-9; H. (Cernuella), H. (Xeroamanda), H. (Xeroamanda), H. (Xeromanda) and H. (Trochoidea) systematics on the basis of genital characters, C. Sacchi; H. claudia sp. nov. environs of Sorrento, C. F. Sacchi (9); Helicella—distribution in the Mediterranean region, C. Sacchi (2).

Helicigona stenomphala albanica subsp. nov. Albania, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 87 f. 23.

Helix pomatia, adaptation to temperature, P. Blazka; H. pomatia structure and physiology of the digestive system, A. Guardabassi; H. aspersa figarolae subsp. nov. Insel La Figarola Grande, near Rovigno, Balkan peninsula, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 88 f. 24; H. (Cantareus) aperta general note, W. Vielhauer; H. hortensis Müll., fluorescence of, P. H. Firscher (3); H. nemoralis L. in a calcium deficient habitat, G. W. Pitchford (1).

†Helix (Pseudotachea) tersannensis minor var. nov. Miocene, Spain, P. Jodot Mem. Comun. Inst. geol., Barcelona 10 1953: p. 44 pl. 1 f. 1-4; H. (Archelix) tectiformis p. 127 pl. 1 f. 2, 4-7, 11; H. (A.) betieri p. 129 pl. 1 f. 9, 10, 13 spp. nov.; H. (A.) betieri conica var. nov. p. 129 pl. 1 f. 8 Miocene, Algeria, P. Jodot Bull. Soc. Hist. nat. Afr. N. 46 3-4 1955; H. (Plebecula) cf. ramondi race minor nov. Oligocene, Jura, P. Jodot Bull. Soc. géol. Fr. 6 4 7-9 1954: p. 541 pl. xxiii b f. 12, 13a, b.

Humboldtiana queretaroana Dall, fig., H. durangoensis Solem, figs.; Humboldtiana pergranulosa sp. nov. San José Range Mts., Durango, Mexico, A. Solem Nautilus, Philad. 69 2 Oct. 1955 p. 42 pl. 3 f. 1-3.

Karlschmidtia gen. nov. Fruticolidae genotype Karlschmidtia leutiformis sp. nov. Peru, F. Haas Fieldiana Zool. 37 1955 pp. 327-328 f. 69.

Lacustrelix subgen. nov. for Semotrachia eyrei H. and A. Adams, Iredale S. Aust. Natst. 18 1937, in text p. 39.

Lysinoe sebastiana Dall, figd., A. Solem.

Macroceramus loeryi sp. nov. Puerte Rico on tree trunks, M. K. Jacobson Arch. Mollusk. 84 1/3 1955 97 f. 1-3.

Meridolum bowdenae p. 39 f. 1 Springwood; M. middenense p. 39 f. 1 Mona Vale spp. nov. N.S.W. Australia, C. McLauchlan Aust. zool. 12 1954.

Minacispira gen. nov. for "Planispira" loxotropis Pfr. Iredale Aust. Zool., Vol. 10 1941, in text p. 91 under Setogibba enigma.

Melcstrachia subgen. nov. for Forresitrachia glomerans sp. nov., Iredale Aust. Zool. 9 2 1938, in text p. 111 under Terresi trachia.

Monacha microtricha sp. nov. between Tetovo and Gostivar, Macedonia, S. Jaeckel Mitt. zool. Mus. Berl. 30 1954: p. 84 f. 20, 21.

Monadenia (Corynadenia) tuolumneana sp. nov. Crystal Cave Tuolumne Co., California, S. S. Berry Bull. S. Calif. Acad. Sci. 54 l 1955 p. 14 pl. 6 f. 1-3.

Mocella cogitata nov. New Zealand Iredale Aust. Zool. 10 1 1941 10, in text p. 91 under Setogibba enigma.

Nannochloritis subgen. nov. type Austrochloritis layardi Jude, Iredale Aust. Zool. 9 1938, in text p. 94 under Austrochloritis blighiana.

Oligomita subgen. nov. of Polymita type Helix versicolor Born., C. de la Torre Mem. Sci. Cubana, Hist. mat. 20 1 1950 p. 18.

Papuina (Pinnadena) periwonensis sp. nov. Solomon Islands, R. K. Dell Pacif. Sci. 9 3 1955: p. 327 f. 2b-d.

Patrubella subgen. nov. for Austrochloritis buxtoni Bray, Iredale Aust. Zool. 9 2 1938, in text p. 95 under above specific names. Phrenegibba gen. nov. for "Planispira" scheepmakeri Pfr. Iredale Aust. Zool., Vol. 10 1941, in text p. 91 under Setogibba enigma.

Polymita picta iolimbata p. 11 pl. 2; P. p. fuscolimbata p. 11 pl. 3, 4; P. p. nigrolimbata p. 13 pl. 5; P. p. roseolimbata p. 13 pl. 6 subspp. nov. Cuba, C. de la Torre Mem. Soc. Cubana Hist. Nat. 20 1 1950; P. versicolor reticulata var. nov. pl. x; P. brocheri cuestana, P. b. ovandoi subsp. nov. pl. x: Cuba, C. de la Torre Mem. Soc. Cubana Hist. nat. 20 1 1950.

Pseudopartula arborascens sp. nov. Harashar region Pulau Panaiten, Butot Palae. Panaiten Treubia 23 1955 p. 129 f. 30–32.

Qualigibba subgen. nov. for Sotogibba enigma sp. nov. Iredale, Iredale Aust. Zool. 10 1941 p. 91, in text under the above species.

Solaropsis (Psadara) angulifera sp. nov. Chapare, Bolivia, F. Haas Fieldiana zool. 34 1955: p. 370 f. 75; S. (Psadara) rugifera Dohrn. description and figure, F. Haas (3).

Strangesta sanguinolenta sp. nov. Sugarloaf Range, N.S.W. Australia, C. McLauchlan Aust. zool. 12 1954: p. 39 f. l.

Trichia sericea in Brandenburg, I. Pfitzner.

Trichochloritis (Yakuchloritis) subgen.
nov. subgenotype: Chloritis albolabris
(Pilsbry et Hirase) together with T.
hoshiyamai Kuroda (MS.) and T.
obscurus (Pilsbry et Hirase) Japan,
T. Habe Venus 18 4 1955: p. 224
[p. 232, in English] f. 11.

Yakuchloritis subgen. nov. of Trichochloritis subgenotype albolabris Pilsbry and Hirase, Habe Venus 18 4 1955 p. 224 f. 11.

STREPTAXACEA

Cymatona tomlini "n. sp." nom. nov. for Nassaria kampyla Tomlin, 1948 B.A.N.Z.A.R.E. Ser. B, 5 (5), p. 228, pl. 2 f. 5 (not of Watson, 1883).

Discartemon stenostomus p. 83, f.5; D. platymorphus p. 88, f. 6; D. leptoglyphus p. 90, f. 7; D. hypocrites p. 92, f. 8 spp. nov. Malaya, W. S. S. v. Benthem Jutting Bull. Raffles Mus. No. 25 [1954] 1955. Gonaxis kibuszsensis E. A. Smith with introduction as a control of Achatina fulica Bowdich, A. R. Mead.

Oophana (Haploptychius) eutropha p. 96, f. 9; O. (H.) thamnophila p. 98, f. 10—O. (H.) atopospira p. 102, f. 12— O. (H.) diaphanopepla p. 104, f. 13 spp. nov. Malaya W. S. S. v. Benthem Jutting Bull. Raffles Mus. No. 25 [1954] 1955.

Paryphanta watti an extension of range in New Zealand, N. W. Gardner.

†Rhytida yaldwyni sp. nov. Holocene New Zealand, R. K. Dell Rec. Dom. Mus., Wellington 2 3 1956: p. 135 f. 2, 3, 4.

Streptaxis (Streptartemon) extraneus, p. 102 f. 1-3; sp. nov. Serra Formosa Mulata Brazil, F. Haas Arch. Mollusk. 84 1/3 1955.

Streptostele horei a further note of rectification and addition, Venmans (2); Streptostele (Raffraya) horei E. A. Smith, Notes, Venmans; Anatomy, H. Watson.

SCAPHOPODA

Antalis H. & A. Adams 1854, validation of name, Opinion 361.

Dentalium tiruhana pp. 37 48, text figs. 17-22. D. (Antalis) glaucarena pp. 37 48 text figs. 16, 18, 21, spp. nov. Chatham Rise, New Zealand, R. K. Dell Dom. Mus. Rec. 2 1 1953.

†Dentalium yotsukurensis sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo, Kyoiku Daig. 4 1955: p. 109 pl. 4 f. 26-28; D. acutoides nom, nov, for D. acutum Richardson 1906 non Hébert 1849: D. boussaci for D. martini 1911 non Whitfield 1882; D. insuranum for D. (Fustiaria) martini Dollfus 1915 non Whitfield 1882; D. invalidum for D. laquestum regulare Henderson 1920 non D. regulare Ahlburg 1906; D. suteri for D. (Episiphon) arenarium Suter 1907 non Römer 1855, nomm. nov., W. K. Emerson Proc. biol. Soc. Wash. 67 1954 pp. 183-185; D. readi for D. simile Read 1907 non Wissman 1841; D. richardsoni for D. subquadratum Richardson 1906 non Meek 1860; D. sangicasii for Articlaria de la disconi giorgii for Antale dentale alternans Sangiorgi 1926 non Dentalium alternans Chenn 1842 nec Müller 1849 nec Ryckholt 1852 D. waihoraensis for D.

(Lavideritaliua) filam Marwich 1931; D. subgracilis for D. gracilis Grönwall & Harder 1907 non Hall & Meek 1854 nec Jeffreys 1870 nec Philippi 1887 D. subparvulum for D. parvulum for D. parvulum Richardson 1906 non Fustiaria parvula Stoliczka 1886, nec D. parvulum Philippi 1887; D. subquadrangulare for D. quadrangulare Deshayes 1833 non Broderip & Sowerby 1832; nomm. nov., W. K. Emerson Proc. biol. Soc. Wash. 67 1954 pp. 183–187.

Pelecypoda or Lamellibranchia

Bivalvia in Bronns Klassen monograph, F. Haas (4).

Lamellibranch of the Miocene of Portugal, Ferreira (1).

Pelecypoda Acrosome reaction on spermatozoa, Dan & Wada; Pelecypod fauna of the Arctic marine analysed, D. Nicol; Pelecypoda of the Gulf of Mexico, H. A. Rehder.

PRIONODESMACEA

†Casterella gen. nov. C. gratiosa sp. nov. Permian Brazil, J. C. Mendes Bol. Soc. Lilor Ciênc. S. Paulo 145 8 1952 pp. 4 100 pl. i f. 6a, b, pl. iii f. 9, 10.

†Casterella (?) camargoi sp. nov. Tertiary, Brazil, K. Beurlen Bol. Geol. Min. Caracao 152 1954 : p. 16 fig. f.

†Edmondia salamunii sp. nov Tertiary, Brazil, K. Beurlen Bol. Geol. Min. Caracao 152 1954: p. 14 f. a, b.

NUCULACEA

†Acila (Truncacila) hidakensis p. 124 pl. 29 f. 20, 20a sp. nov.; A. (T.) gottschei ôsugii p. 125 pl. 29 f. 18, 19 var. nov.; A. (Acila) kusiroensis p. 135 pl. 31 f. 10-11b sp. nov.; A. (A.) vigilia brevis p. 132 pl. 30 f. 1-4 var. nov. Miocene; A. (A.) praedivaricata p. 137 pl. 31 f. 5-8 sp. nov.; A. (A.) vigilia elongata p. 133 pl. 31 f. 1-4 var. nov. Oligocene—Miocene, Japan, T. Nagao & K. Huzioka J. Fac. Sci. Hokkaido Univ. (4) 6 1941; A. (Truncacila) oyamedensis sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955 : p. 75 pl. 1 f. 1-7.

† ?Ctenodonta arroyana "n. sp." Ordovician, Argentina, C. Rusconi Rev. Mus. Hist. nat. Mendoza 8 1-4 1955: p. 46 pl. 5 f. 8.

Exacholeda names for group represented by L[eda] dasea Hedley, **T.**Iredale Mollusca Part I. Great Barrier Reef Expedition 1928–29 Sci. Report **5** No. 6 1939 p. 241.

Kamaleda names for new group represented by L[eda] macleriformis Smith, T. Iredale Mollusca Part I Great Barrier Reef Expedition 1928–29 Sci. Report vol. V No. 6 1939 p. 341.

Nucula schmidti nom. nov. p. 1 (Nucula compressa Philippi, 1846—van Straelen, 1923—Görges, 1952 (non Sowerby, 1834); N. orbignyi (Nyst mss. 1868) sp. nov. p. 2 Oligocene, Belgium, M. Glibert Bull. Inst. Sci. nat. Belg. 31 86 1955.

†Nucula renouardi sp. nov., Portlandian, Liban [Abyssinia], D. Mongin Bull. Soc. géol. Fr. 6 4 7-9 1954 p. 501 pl. xxiiia f. 1, 2; N. ventricosa sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 77 p. 1 f. 12-15.

†Nuculites pusillus sp. nov. Devonian, Germany, A. Rösler Notizbl. Hess. Landesmt. Bodenforsch, 81 1953: p. 143 pl. 7, f. 6, 7.

Palaeoneilo emendation, Opinion 215.

† Paleoneilo iwaiensis sp. nov. Triassic. Tokyo Prefecture, Japan, K. Ichikawa Jap. J. Geol. Geogr. 25 1–2 1954: p. 181 pl. 17 f. 1–3.

†Solemya (Acharax) prashadi nom. nov. for S. (A.) grandis Prashad 1932 (non Verrill and Bush 1898); S. stantoni nom. nov. for S. occidentalis Stanton 1895, (non S. (Petrasma) occidentalis Deshayes 1857), H. E. Vokes J. Paleont. 29 3 1955: p. 538; S. tokunagai elongata subsp. nov. Miocene, Fukushima Prefecture, Japan, S. Aoki Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 30 pl. 1 f. 15.

†Straba gen. et subgen. nov. Strabinae (subfam. nov. of Ctenodontidae) p. 10: genotype Straba bohemica sp. nov.—
Straba (Straba) subgen. nov. p. 11: subgenotype Straba (Straba) bohemica p. 12 pl. 2 f. 1, 2, 5; pl. 3 f. 3, 4; S. (Straba) barrandei p. 14 pl. 1 f. 1-6; pl. 2 f. 3-4; pl. 3 f. 1, 2; pl. 4 f. 2; S. (Straba) kettneri p. 18 pl. 2 f. 6, 8; pl. 4 f. 4 spp. nov.; Straba (Strabina) subgen. nov. p. 20 subgenotype: Straba (Strabina) formosa p. 21 pl. 2 f. 7; pl. 4 f. 3 sp. nov.—Straba (Strabiella)

subgen. nov. p. 23 subgenotype: Straba (Strabiella) holynensis p. 24 pl. 1 f. 7; pl. 4 f. 1 sp. nov. Devonian, Bohemia, F. Prantl & B. Růžička Acta Mus. nat. Prag. Series B 10 3 1954.

†Strabiella subgen, nov. of Straba gen. nov. q. v. F. Prantl & B. Růžička.

†Strabina subgen, nov. of Straba gen. nov. q. v. F. Prantl & B. Růžička.

†Strabinae subfam. nov. of the family Ctenodontidae, type-genus: Straba gen. nov. Devonian, Bohemia, F. Prantl & B. Růžička Acta Mus. nat. Prag. Series B 10 3 1954: p. 10.

†Stutchburia? argentinensis sp. nov. Permian, Argentina, H. J. Harrington J. Paleont. 29 1 1955: p. 117 pl. 24 f. 1-3.

†Trigonucula sakawana inequilatera var. nov. Triassic, Tokyo Prefecture, Japan, K. Ichikawa Jap. J. Geol. Geogr. 25 1-2 1954: p. 183 pl. 17 f. 7.

† Yoldia omorii sp. nov. Miocene, Fukushima Prefecture, Japan, S. Aoki Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 30 pl. 2 f. 18; Y. kawadai sp. nov. Miocene, Tochigi Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 48 pl. 3 f. 1.

ARCACEA

Arcacea, light perception, R. Braun.

Arca novaezelandiae in captivity, G. Williams (1); Arca (Senilia) senilis Lamarck anatomy and habits, figs., C. M. Yonge.

Arcidae, A report on the family, figs., H. Rost.

†Barbatia? cuniculana sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264-B 1954: p. 27 pl. 6 f. 2-5.

Cosatova gen. nov. type Philobrya recapitula Hedley, T. Iredale Mollusca Part I, Great Barrier Reef Expedition 1928-29 Sci. Report 5 No. 6 1939 p. 304.

Epideisis name proposed for Arca semitorta Lmk., T. Iredale Mollusca Part 1, Great Barrier Reef Expedition, 1928-29 Sci. Report 5 No. 6 1939 p. 270. †Glycymeris convexa; G. crassicosta 'n. sp.' name without description. Pliocene, Tyôsi City, Japan, H. Ozaki Bull. nat. Sci. Mus. Tokyo N.S. 1 2 1954: p. 53; Glycymeris subovata (Say) fig., H. E. Vokes.

Indacar gen. nov. for "species dissected by Ghosh under the identity of Arca pterossa Smith," (See Inst. J. Morph. 1921, 457-459), T. Iredale Mollusca Part I Great Barrier Reef Expedition 1928-29 Sci. Report vol. V No. 6 1939 p. 261.

†Mnataia gen. nov. Pelecypoda (pseudocyrtodont type?) p. 4: genotype, by monotype: Mnataia přibyli sp. nov. p. 5 pl. 1 text-f. 1–9 Devonian, Bohemia, F. Prantl & B. Růžička Acta Mus. nat. Prag. Series B 11 5 1955.

†Nemodon obesus sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264-B 1954: p. 28 pl. 6 f. 10-15.

Opularca gen. nov. Arcidae type O. tenalla egenora subsp. nov., T. Iredale Mollusca Part I, Great Barrier Reef Expedition 1928–29 Sci. Report vol. V No. 6 1939 269 pl. iii p. 2, 2a.

†Parallelodontidae of Peru, R. Rivera.

Pseudocucullaea rentemiensis sp. nov. Senonian Pougo de Rentema R. Marañon, Peru, R. Revera Bol. Soc. geol. Peru 25 2 1949 Fasc. 17 pp. 7, 9 pl. i f. 1, 2.

Tucetopsis subgen. nov. of Tucatona type Cardium amboinensis Gmelin, T. Iredale Mollusca Part 1 Great Barrier Reef Expedition 1928–29 p. 303 Sci. Report vol. V No. 6 1939.

MYTILACEA

Amygdalum americanum sp. nov. Isla Gorgoma, Colombia 20 fms. T. Soot Ryen Allan Hancock Pacif. Exped. 20 1 1955 p. 70 pl. 8 f. 37.

†Brachidontes obessus sp. nov. Miocene, Tochigi Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 50 pl. 3 f. 9, 10.

Choromytilus Soot-Ryen 1952, Soot Ryen.

Crenella tamurai sp. nov. Akkeshi Bay, Japan, T. Habe Publ. Akkeshi mar. biol. Sta. No. 4 1955: p. 26 pl. 7 f. 1, 2.

†Crenella microstriata sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274E 1955: p. 114 pl. 17 f. 7-10.

Crenomytilus gen. nov. = Mytiloconcha auct. non Conrad. Mytilidae genotype Mytilus grayanus Dunker 1853, T. Soot Ryen Allan Hancock Pacif. Exped. 20 1 1955 p. 23, pl. 2 f. 9-10, text fig. 7.

Dacrydium (Quendreda) elegantulum sp. nov. Bahia de Gardner, Galapagos Islands, T. Soot-Ryen Allan Hancock Pacif. Exped. 20 1 1955 p. 87 pl. 8 f. 41.

Doliolabis sec "nominated" of Lithophaga, T. Iredale Mollusca Part 1 Great Barrier Reef Expedition 1928–29 Sci. Report 5 No. 6 1939 p. 417.

Exoliberus sec of Lithophaga "nominated," T. Iredale Mollusca Part 1 Great Barrier Reef Expedition 1928–29 Sci. Report 5 No. 6 1939 p. 417.

Lithodomus boring habits, E. Seibold.

†Lithodomus mitzopoulosi sp. nov. Neogene, Greece, S. Charalambakis Ann. géol. Hellen. 1 4 1952 : p. 105 pl. viii f. 2.

Lithophaga (Leiosolenus) hancocki sp. nov. Isla Onslow, N. of Isla Floreana (Charles Island) Galapagos Islands, T. Soot-Ryen Allan Hancock Pacif. Exped. 20 1 1955 p. 102 pl. 10 f. 60.

Modiolus neglectus sp. nov. syn. Volsella recta auct. non Conrad; V. glabellata auct. non Gould., San Diego, California, T. Soot Ryen Allan Hancock Pacif. Exped. 20 1 1955 p. 64 pl. 7 f. 31-32 text-fig. 49, 51, 59.

Myapalmula sec of Lithophaga 'nominated', T. Iredale Mollusca Part 1 Great Barrier Reef Expedition 1928–29 Sci. Report 5 No. 6 1939 p. 417.

Mytella gen. nov. Mytilidae genotype Modiola guyanensis Lamarck 1819, T. Soot Ryen Allan Hancock Pacif. Exped. 20 1 1955 p. 47 figs. 36, 41-44, 47.

Mytilidae, A report on the family, figs., T. Soot-Ryen.

Mytilus californianus habitat, S. Berry (3); M. galloprovincialis, Heldt; Mytilus Observation of straining of suspensions, Tammes & Deal; M. galloprovincialis Parasitism and Commensalism, Genovese; M. californianus

tidal rhythmicity of water propulsion rate, K. Rao; M. edulis the heart beat, C. Schlieper.

†Pelecypodichnus amygdaloides gen. nov., sp. nov. Jurassic, Central Europe, A. Seilacher Neues Jb. Min. Geol. Pälaont. 98 1 1953 p. 105 pl. 10 f. 1, pl. 12.

Salebrolabis sec of Lithophaga for L. divaricalx, T. Iredale Mollusca Part I Great Barrier Reef Expedition 1928– 29 Sci. Report Vol. V No. 6 1939 p. 417, 418.

Semimytilus gen. nov. Mytilidae genotype Mytilus algosus Gould 1850, T. Soot-Ryen Allan Hancock Pacif. Exped. 20 1 1955 p. 25 pl. 4 f. 17, text figs. 8, 9, 14-16.

Stumpiella subgen. nov. of Lithophaga subgenotype Lithophaga (Stumpiella) calyculata (Carpenter) 1856, T. Soot-Ryen Allan Hancock Pacif. Exped. 20 1 1955 p. 93 pl. 10 f. 61–63.

Volsella difficilis sp. nov. for V. modiolus auctt. non Linne, Kuroda & Habe Illust. Catal. Jap. Shells. No. 4 1950 p. 30.

PTERIACEA

Anisoperna subgen. nov. types Perna australida Reeve, Iredale Mollusca Part 1 Great Barrier Reef Exped. 1928-9 Sci. Report 5 No. 6 1939 p. 122.

Aucella Keyserling 1946 proposal to validate and Avicula mosquensis as type, Jeletzky; Aucella preservation of name and the type species Avicula mosquensis Buck, J. A. Jeletzky.

†Avicula revision of J. Barrand's figured specimens of 1881, B. Ružička (1).

†Bakevellia matsushitai p. 214 pl. 3 f. 1-5, 8; B. oyogiensis p. 216 pl. 3 f. 9; pl. 4 f. 1-4; B. monobensis p. 217 pl. 4 f. 5-9; pl. 5 f. 1a-d, 2; B. subhekiensis p. 219 pl. 6 f. 4-7 spp. nov. Triassic, Kyoto Prefecture, Japan, K. Nakazawa Mem. Coll. Sci. Kyoto 2-B 2 1954.

Buchia Rouillier 1845 proposal to suppress, Jeletzky.

†Conocardiidae—nomenclatural review, D. Nicol (2).

†Eumorphotis multiformis shionosawensis subsp. nov. Shionosawa limestone, Japan, K. Ichikawa & Y. Yabe Trans. Palaeont. Soc. Japan N.S. 17 1955: p. 6 pl. 2 f. 1-15. Isognomon maxillata (Desh.) fig., H. E. Vokes.

Inoceramus on retention of name and type to be accepted, L. R. Cox.

†Liopteria bonaerensis p. 120 pl. 24 f. 10—L. ? dutoiti p. 120 p. 24 f. 5, 11 spp. nov. Permian, Argentina, H. J. Harrington J. Paleont. 29 1 1955.

Pinctada martensii biochemical studies, Horiguchi & Miyake; Pinctada revision of the Australian pearl-shells, J. S. Hynd; P. martensii—metabolism. Carbonic anhydrase, K. Ikinaga; P. martensii death due to bacteria, Kotoke & Miyawaki.

Pteria pinctada matensii influence of diluted sea water on heart beat, Kawamoto & Motoki; P. (Pinctada) martensi influence of diluted sea water, Kawamoto & Motoki (1).

Pteralectroma subgen. nov. of Electroma for Avicula zebra Reeve, T. Iredale Mollusca Part 1 Great Barrier Reef Expedition 1928–29 Sci. Report Vol. V No. 6 1939 p. 332.

†Tenuipteria gen. nov. Pteriacea p. 110 type species: Inoceramus argenteus Conrad 1858; T. argentea p. 111 pl. 16 f. 4-9 sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955.

PECTINACEA

†Amussiopecten hyugaensis sp. nov. Tertiary, Japan, T. Shuto Trans. Palaeont. Soc. Japan N.S. 20 1955: p. 105 pl. 16 f. 1-5, pl. 17 f. 2, 4, 5, text-f. 3.

†Aviculopecten rotundatus—description of figured species of J. Barrande 1881, B. Ružička.

†Chlamys rapovensis sp. nov. Miocene, Eastern Europe, V. Hano & J. Senes Geol. sborn. Slovensk. Akad. 3 3-4 1953: p. 323 Pl. xliii f. la-d; Pl. xliv f. 1, 2; C. (Chlamys) polyaktinos sp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 30 pl. 4 f. 16 [nom. nud. loc. cit. 77 1954: p. 56].

†Chlamys kakisakiensis tokawaensis 'subsp. nov.' name without description. Pliocene, Tyôsi City Japan, H. Ozaki Bull. nat. Sci. Mus. Tokyo N.S. 1 2 1954: p. 53. †Coralichlamys shigemai sp. nov. Miocene, Tochigi Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig 3 C 1954: p. 51 pl. 3 f. 2.

†Cryptopecten oyamadensis sp. nov. Miocene, Tochigi Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 54 pl. 3 f. 3.

†Ctenamussium (Bathyamussium) amakusaensis p. 14 pl. 1 f. 8, 9—C. (B.) inouei p. 15 pl. 1 f. 7—C. (B.?) cancellata p. 16 pl. 1 f. 13 spp. nov. Oligocene, Miocene, Japan, M. Ômori Sci. Rep. Tokyo Kyoiku Daig. 4 1955.

Edentiplica "name" for Pecten strangie Resoe, T. Iredale Mollusca Part 1 Great Barrier Reef Expedition 1928– 29 Sci. Report 5 No. 6 1939 p. 362.

Eltopera subgen, nov, of Spondylus type S. pacificus fortior subsp. nov., Iredale Mollusca Part 1 Great Barrier Reef Expedition 1928–29 Sci. Report 5 No. 6 1939: 372.

Escalima regularis 'n. sp.' nom. nov. for Lima angulata Suter, 1913, Manual N.Z. Moll., p. 885 (not of Sowerby: N.Z. Recent records only), A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 27.

Lima angolensis sp. nov. South Atlantic, West African coastal waters, W. Adam & J. Knudsen Bull. Inst. Sci. nat. Belg. 31 61 1955: p. 19 pl. 2 f. 1-4; L. (Acesta) mori sp. nov. deep water off Central California, L. G. Hertlein Proc. Calif. Acad. Sci. (4) 27 12 1952: p. 379 pl. 20 f. 12, 13.

†Lima robini sp. nov. Tertiary, Campbell Island, New Zealand, C. A. Fleming Cape Exped. Ser. Bull. No. 3 1950: p. 50 pl. 9 f. l, 6; L. becki sp. nov. Miocene, New Zealand, C. A. Fleming Trans. roy. Soc. N.Z. 82 5 1955: p. 1052 pl. 40 f. 3.

†Limatula craigensis sp. nov. Miccene, New Zealand, C. A. Fleming Trans. roy. Soc. N.Z. 82 5 1955 : p. 1052 pl. 40 f. 5.

Lanilda subgen. type Sprodylus ducalis Bolten T. Iredale Mollusca, Part 1 Great Barrier Reef Expedition 1928-29 Sci. Report vol. V No. 6 1939 pp. 372, 373.

Lentipecten adelaidensis sp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 32 pl. 1 f. 13a, b, c [nom. nud. loc. cit. 77 1954: p. 56].

†Parvamussium takasensis 'n. sp.' [nom. nov.] for Propeamussium transnipponica Suzuki, 1952 Geology in the north-eastern part of Yamagata City Res. Fac. of Arts Yamagata Univ., vol. 2 No. 1 p. 50 (not Propeamussium transnipponica Mochizuki) Yamagata Prefecture, Japan, Miocene, M. Omori Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 12 pl. 2 f. 1.

Pecten irradians water filtration, W. Chipman & J. G. Hopkins; P. jacobaeus byronensis subsp. nov. Byron Bay, N.S.W., C. A. Fleming Aust. Zool. 12 1955 p. 108 fig. 1; P. tereinus Dall a synonym of Pecten (Euvola) chazaliei Dautzenberg a rectification, G. Grau; P. maximus L. marking experiment, Mason & Colman.

Pectinidae of the Miocene of the Azores, Ferreira.

Placopecten magellanicus abundance in Bay of Funda, L. Dichie.

† Pleuronectites hirabarensis sp. nov. Triassic, Japan, M. Amano Trans. Palaeont. Soc. Japan N.S. 17 1955 p. 25 pl. 5 f. 1-7.

†Plicatula ferrata sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264-B 1954: p. 31 pl. 7 f. 6.

†Propeamussium circularis sp. nov. upper Oligocene or lower Miocene, Nagano Prefecture, Japan, M. Ômori Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 16 pl. 1 f. 10, 11.

Sponvola subgen. nov. of Spondylus type S. wrightianus ella subsp. nov., T. Iredale Mollusca Part 1, Great Barrier Reef Expedition 1928–29 Sci. Report vol. V No. 6 1939 pp. 372, 379 pl. vi f. 1.

†*Vertumnia volitans*—description of figured specimens of J. Barrande 1881, B. Ružička.

ANOMIACEA

†Anomia asagaiensis sp. nov. Tertiary Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 83 pl. 1 f. 23–26.

OSTREACEA

Crassostrea virginica—changes in shell structure indicative of drought conditions, R. H. Parker; Crassostrea virginica uptake of phosphate ions from sea water, L. Pomeroy & Haskim.

†*Lopha karrenbergi* **sp. nov.** Jurassic, Yémen, **E. Basse** Bull. Soc. géol. Fr. **6 4** 7–9 1954 p. 664 pl. 27 f. 1a–d.

Gryphaea Lmk. 1801, see Opinion 356; Gryphaea Lamarck 1801 as nom. conserv. and type G. arcuata Lamarck, Opinion 338; Gryphaea problems of oyster taxonomy, J. Hedgpeth; G. gigas from Akheshi Lake (Japan). Notes on Shell structure, M. Miyawaki.

Ostrea knorrii Voltz 1828 validation of name, Opinion 360; Ostrea L. 1758, see Opinion 356; Ostrea summary of knowledge of the genus in the Gulf of Mexico, P. A. Butler; Ostraea and the parasite Dermocystidium marinum, fig., Ray & Chandler; Ostrea. Purification in simple pits, figs., H. A. Cole; O. lurida mortality at low temperatures, H. Davis; Ostrea a study of the biology and hydrography at Crystal R., Florida, C. E. Dawson.; O. laperousi Schrenck, nucleic acids and nucleoproteins contents, H. Kobayashi; Ostrea growth and weight, J. L. McHugh; O. equestris Say phases of biology and comparison with Cressontrea virginica, R. W. Menzel; Ostrea-Reefs in the G. of Mexico. W. A. Price.

†Ostrea iphigeniae sp. nov. Neogene, Greece, S. Charalambakis Ann. géol. Hellen. 1 4 1952: p. 93 pl. 5, 6; O. (Alectryonia) jerseiana sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264-B 1954: p. 29 pl. 6 f. 20.

Placenta placenta in the G. of Cutch, Gokhale.

Pretostrea subgen. nov. of Ostrea for Ostrea baesia sp. nov., T. Iredale Mollusca Part 1, Great Barrier Reef Expedition 1928-29 Sci. Report vol. V No. 6 1939 397 pl. vii f. 4.

TRIGONIACEA

†Geratrigonia gen. nov. Trigoniinae, p. 171, Type species: Trigonia hosourensis Yokoyama, 1904; G. lata sp. nov. p. 173, pl. 15 f. 6-7; G. hosourensis convexa var. nov. p. 172 pl. 16 f. 9a, b. Jurassic, Japan, Kobayashi Jap. J. Geol. Geogr. 25 1954.

[1955]

†Hijitrigonia subgen. nov. of Vaugonia Jurassic, cosmopolitan distribution, Kobayashi Jap. J. Geol. Geogr. 26 1955: p. 84.

Hoffetrigonia and Trigonioides belong to the Unionidae [q.v.] and not to the Trigoniidae, L. R. Cox (1).

Lyriomyophoria gen. nov. "to comprise konzentrisch skulptierte Myophorien;" type species: Myophoria elegans Dunker, 1849, Kobayashi Jap. J. Geol. Geogr. 25 1954: p. 66.

†Myophorella (Myophorella) dekaiboda sp. nov. p. 95 pl. 6 f. 6-9; M. (Promyophorella) subgen. nov. subgenotype: M. (P.) sigmoidalis p. 96 pl. 5 f. 1-3; M. (P.) sugayensis p. 97 pl. 5 f. 4; M. (P.) orientalis p. 98 pl. 5 f. 6, a, b; M. (P.) obsoleta p. 99 pl. 5 f. 7 spp. nov.; M. (P.) sugayensis geniculata var. nov. p. 97 pl. 5 f. 5; M. (Haidaia) crenulata p. 100 pl. 5 f. 8-10; M. (H.) subcircularis p. 101 pl. 6 f. 3 spp. nov.; M. (H.) crenulata lunulata p. 101 pl. 6 f. 1a, b, 2 var. nov. Jurassic, north Japan, Kobayashi & Tamura Jap. J. Geol. Geog. 26 1955.

†*Oistotrigonia prima* sp. nov. Jurassic, Fukushima Prefecture, Japan, Kobayashi & Tamura Jap. J. Geol. Geogr. 26 1955: p. 102 pl. vi f. 4a-b, 5.

†Orthotrigonia midareta p. 87 pl. 4 f. 11a, b; O. corrugata p. 87 pl. 6 f. 10 spp. nov. Jurassic, Kitakami Mts., North Japan, Kobayashi & Mori Jap. J. Geol. Geogr. 26 1955.

†Promyophorella subgen. nov. of Myophorella middle Lias to Neocomian, cosmopolitan distribution, Kobayashi & Tamura Jap. J. Geol. Geogr. 26 1955: p. 96.

†Trigomyophoria gen. nov. "for cancellate Myophorians on the basis of Myophoria decussata Munster, 1938", Kobayashi Jap. J. Geol. Geogr. 25 1954: p. 66.

†Trigonia senex p. nov. Jurassic, Miyagi Prefecture, Japan, Kobayashi & Mori Jap. J. Geol. Geogr. 25 1954: p. 167, pl. 16, f. 8a, b; T. longa costatissima pl. 1 f. 1, 2; T. corbierensis "sp. var. nov. Cretaceous, France p. 62 (6) nov." = T. corbierensis Villatte 1954.

Cretaceous, France p. 63 (7) pl. 1 f. 4, 4a, 4b, 5; T. padernensis "n. sp." [nom. nov.] = T. jacobi Villatte 1954; p. 67 (11), G. Petter-Receveur Ann. Paléont. 41 1955.

Trigonioides and Hoffetrigonia belong to the Unionidae [q.v.] and not to the Trigoniidae, L. R. Cox (1).

†Vaugonia niranohamensis p. 80 pl. 3 f. 1-6; pl. 4 f. 1; V. kodaijimensis p. 82 pl. 3 f. 12-15; V. namigashira p. 83 pl. 3 f. 10, 11 spp. nov.; V. yokoyamai p. 81 pl. 3 f. 7-9 sp. nov. [For Trigonia v-costata (Lycett) Yokoyama, 1904, non Lycett); †Vaugonia (Hijitrigonia) subgen. nov. Type species: V. (H.) geniculata p. 84 pl. 4 f. 4-9; V. (H.) kojiwa p. 85 pl. 4 f. 2, 3 spp. nov. Jurassic, Kitakami Mts., North Japan, Kobayashi & Mori Jap. J. Geol. Geogr. 26 1955.

UNIONACEA

Anodonia cygnea action of acetylcholine and eserine on heart and intestine, J. Ten Cate; A. cygnea on the heart nerves, J. T. Cate; Anodonia experimental studies, R. Fange.

Bartlettia stefanensis Moricand, apparent first record from Peru, F. Haas (3).

Caelatura (Zairia?) cridlandi sp. nov. p. 135 f. 70; C. hauttecoeuri kyogae subsp. nov. p. 134 f. 68a Uganda, Mandahl-Barth, Ann. Mus. Congo belg. 8vo 32 1954; Caelatura aegyptiaca Caillaud 1827 f. calathus Bourguignat 1885 and its anatomy, figs., J. Bouillon.

Cardinia as from Agassiz, J. L. R., 1841, Validation, Opinion 292.

Elliptio spinosus in the Altamaha River, I. R. Tomkins.

Hoffetrigonia belongs to the Unionidae and not to the Trigoniidae, L. R. Cox (1).

Hyriella australis (Lmk.) 1819 its identity and validity, D. F. McMichael.

Hyriopsis schlegelii studies on the pearl-sac, Ojima & Ikenaga.

Jardinella? nov. for Petterdiana thaanumi Pils., Iredale & Whitley S. Austl. Natst. 18 1938, in text under Jardinean fluvifaunulae p. 67.

Lamellidens lamellidens oxygen carrier in the blood, P. Rangneker (1).

Leptodea laevissima abundant in Lake Springfield, Illinois, Parmalee. Margaritifer margaritifera growth, C. R. Boettger (3).

†Naiadites of the Carboniferous of Gt. Britain, A. Trueman & Weir.

†Naritra gen. nov. Pollicidae, p. 31 type species: N. polliciformis sp. nov. p. 32 pl. 7 f. 7-9. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264-B 1954.

†Psilunio stegersbachensis sp. nov. Pannon (Tertiary) Austria, F. Sauerzopf Burgenland. Heimat Bl. 14 l 1952: p. 9 pl. 1 f. 1-2.

Quadrula quadrula abundant in Lake Springfield, Illinois, Parmalee.

Trigonioides belongs to the Unionidae and not to the Trigoniidae, L. R. Cox (1)

Unionetta nom. nov. = Unionella Haus 1913 non Etheridge 1888, F. Haas, Archiv. f. Molluskenk 84 4/6 1955 p. 212.

Unionid fauna of Lake Springfield, Illinois and some ecological aspects, P. W. Parmalee.

Unionidae studies on the Natural History, M. R. Matteson.

ASTARTACEA

†Astartella? pusilla sp. nov. Permian, Argentina, H. J. Harrington J. Paleont. 29 1 1955: p. 118 pl. 25 f. 4; Astartella morphology, D. Nicol (1).

†Cuna caerulea sp. nov. Miocene, New Zealand, C. A. Fleming Trans. roy. Soc. N.Z. 82 5 1955: p. 1053 pl. 40 f. 6, 7.

†Opis? elevata biangulata var. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264-B 1954: p. 32 pl. 7 f. 15, 16.

CARDITACEA

Cardita subdeceptiva sp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 40 pl. 4 f. 14 [nom. nud. loc. cit. 77 1954: p. 56].

†Cyclocardia (Arcturellina) hindmarshensis p. 44 pl. 2 f. 9; C. (A.) peridonea p. 44 pl. 2 f. 7 spp. nov. Pliocene, Australia, N. H. Ludbrook, Trans. roy. Soc. S. Aust. 78 1955 [nom. nud. loc. cit. 77 1954 : p. 57].

Fulvia mutica Reeve food in the Selonaikai inland sea of Japan, Chiba, Tsuruta & Inque.

Pleuromeris subpecten sp. nov. Pliocene, Australis, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 42 pl. 2 f. 3 [nom. nud. loc. cit. 77 1954: p. 56].

†Venericardia southlandica sp. nov. Miocene, New Zealand, C. A. Fleming Trans. roy. Soc. N.Z. 82 5 1955: p. 1053 pl. 40 f. 11, 12; V. haskelli sp. nov. Tertiary, Campbell Island, New Zealand, C. A. Fleming Cape. Exped. Ser. Bull. No. 3 1950: p. 52 pl. 9 f.2-5; V. (Cyclocardia) otatsumei sp. nov. Tertiary, Japan, S. Uozumi Trans. Palaeont. Soc. Japan, N.S. 19 1955: p. 76 pl. 12 f. 2.

SPHAERIACEA

Byssanodonta ovata p. 176 f. 94; B. crassa p. 178 f. 96 spp. nov. Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

Corbicula japonica habitats of this group and some ecological aspects, A. Fuji; Corbicula africana (Krauss, 1848) = C. radiata auct. p. 158; C. africana albertiana subsp. nov. p. 160 f. 86b Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

† Filosina gen. nov. Corbiculidae type F. gregaria sp. nov. Lower Cretaceous Southern England, R. Casey J. Wash. Acad. Sci. 45 12 1955 p. 369 f. 1-36 c-d.

†Hemicorbicula subgen. nov. of Eocallista Corbiculidae subgenotype Cyclas parva J. de C. Sowerby, Upper Jurassic Purbeck beds = Astarte socialis Orbigny, R. Casey J. Wash. Acad. Sci. 45 12 1955 p. 367 f. 4-5, 6b.

Lasaea rubra (Mont.) anatomy and mode of life, figs., E. Oldfield.

Pseudochama and distinctions from genera of the Chamidae, figs., N. H. Odhker.

†Litigiella adelaidensis sp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 57 pl. 3 f. 11 [nom. nud. loc. cit. 77 1954: p. 57].

†Nemetia gen. nov. Corbiculidae type Platopis triangularis Whitfield Lower Cretaceous (Aptian) Syria, R. Casey J. Wash. Acad. Sci. 45 12 1955 p. 371.

Pisidium victoriae p. 173 f. 92—P. fistulosum p. 174 f. 93 spp. nov. Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

Sphaerium victoriae albertianum p. 166 f. 87c, 88; S. victoriae lacuum p. 166 f. 87b, 88 d-e; S. stuhlmanni regularis p. 169 f. 87g, 89d-e subspp. nov. Uganda, Mandahl-Barth Ann. Mus. Congo belg. 8vo 32 1954.

Turtonia minuta (Fab.) Anatomy and mode of life, figs., E. Oldfield.

CYPRINACEA

Arctica islandica, as subject of a new food industry, H. D. Russell.

†Cypricardinia biumbonata; C. pseudomira; C. consobrina—description of figured specimens of J. Barrande 1881, B. Ružička (1).

Cyprina islandica. The ocean quahaug (fig.) in the maritime provinces of Canada, Medcof & MacPhail.

Cyrenopsis Etheridge 1902 placed in Neomiodontidae fam. nov. [q.v.], R. Casey (1).

†Ecmiodon Cox placed in Neomiodontidae fam. nov. [q.v.], R. Casey (1).

†Musculiopsis MacNeil 1939 placed in Neomiodontidae fam. nov. [q.v.], R. Casey (1).

†Myrene gen. nov. (feminine) Arcticacea Neomiodontidae [q.v.] genotype Myrene fittoni sp. nov. Upper Jurassic, S.-E. England, R. Casey Proc. Malac. Soc. London 31 5/6 Dec. 1955, 209 218 pl. 11 f. 1-2 text-fig. 9.

Neomiodon Fischer type genus of Neomiodontidae fam. nov. [q.v.], R. Casev.

†Neomiodontidae fam. nov. of Arcticacea Pelecypoda (= Cyprinacea) including Neomiodon Fischer, Eomiodon Cox, Myrene gen. nov. Casey, Cyrenopsis Etheridge, and Musculiopsis MacNeil, R. Casey Proc. Malac. Soc. London 31 5/6 Dec. 1955: 208.

GAIMARDIACEA

Gaimardia trapesina flemingi subsp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 28 pl. 1 f. 4.

DREISSENACEA

Ciboticola gen. nov. type? Congeria lunata Hedley, T. Iredale Mollusca Part 1, Great Barrier Reef Expedition 1928-29 Sci. Report 5 No. 6 1939 p. 425. †Congeria spathulata praebalatonica p. 9 pl. 1 f. 7-8; C. gitneri majoritesta p. 11 pl. 2 f. 6-7; C. partschi subpannonica p. 11 pl. 1 f. 5-6; C. pancici longiconcha p. 11 pl. 1 f. 3-4 subsp. nov. Pannon, (Tertiary) Austria, F. Sauerzopf Burgenland. Heimat Bl. 14 1 1952.

Dreissena acceptance of name as emendation of Driessena, v. Beneden 1835, Opinion 351; Dreissena polymorpha (Pallas) in a closed pond, G. W. Pitchford (2); Dreissena polymorpha, D. Ludemann.

† Dreissena turislavica stegersbachensis subsp. nov. Pannon (Tertiary) Austria, F. Sauerzopf Burgenland. Heimat. Bl. 14 1 1952: p. 11 pl. 2 f. 8-9.

Miridas gen. nov. for Myrina coppingeri Smith, T. Iredale Mollusca Part 1, Great Barrier Reef Expedition 1928–29 Sci. Report 5 No. 6 1939 p. 425.

LUCINACEA

†Claibornites (Saxolucina) quinquangulus sp. nov. Tertiary, Japan, S. Uozumi Trans. Palaeont. Soc. Japan, N.S. 19 1955: p. 77 pl. 12 f. 6a, b.

†Conchocele compacta minor subsp. nov. Miocene, Tochigi Prefecture, Japan, M. Ômori Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 90 pl. 7 f. 1a, b, 2.

†Eomiltha (Gibbolucina) confirmans sp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 49 pl. 2 f. 11 [nom. nud. loc. cit. 77 1954: p. 57].

†Gonimyrtea salisburyensis p. 50 pl. 2 f. 12; G. crassior p. 51 pl. 3 f. 3, 4; G. validior p. 52 pl. 3 f. 5, 6—G. notabilior p. 53 pl. 3 f. 7, 8 spp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955 [nom. nud. loc. cit. 77 1954 : p. 57].

†Lucina shimogoensis sp. nov. Miocene, Tochigi Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 59 pl. v f. 1.

†Lucinoma katayosensis sp. nov. Miocene, Fukushima Prefecture, Japan, S. Aoki Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 34 pl. 1 f. 13, 14; L. gracilistriata p. 104 pl. x f. 1a-c, 2. Idegawa, Fukushima Prefecture, probably Upper Miocene; L. shinokii p. 105 pl. x f. 3a-c, 5a-b, 4, Kanagawa Prefec-

ture, Lower Miocene spp. nov. Japan, K. Hirayama Jap. J. Geol. Geogr. 25 1-2 1954.

Maorithyas flemingi sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 34 pl. 1 f. 1.

†Miltha harrisi nom. nov. for Lucina cf. landanensis, K. E. Caster 1938? (non Lucina landanensis, E. Vincent 1913) Africa, E. Dartevelle & J. Roger Comun. Serv. geol. Portug. 35 1954: p. 285.

†Monitilora (Monitilora) idonea p. 47 pl. 3 f. 1, 2; M. (Prophetilora) chavani p. 47 pl. 2 f. 13 spp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955 [nom. nud. loc. cit. 77 1954: p. 57].

ERYCINACEA

Aclistothyra gen. nov. Galeomidae genotype A. atlantica sp. nov. off Palm Beach, Florida, T. L. McGinty Proc. Acad. nat. sci. Philad. 107 1955 p. 84 pl. 2 f. 11, 11a.

Erycina type species, Opinion 217.

† Properycina torrensensis sp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. pl. 3 f. 13 [nom. nud. loc. cit. 77 1954: p. 57].

CHAMACEA

Agria in the Cretaceous, Emberger (1).

†Barrettia gigas nom. nov. for B. monilifera Whitfield 1897 non Woodward 1862, L. J. Chubb Amer. Mus. Novit. No. 1713 1955 p. 9

†Durania coahuilensis sp. nov. Cretaceous, Mexico, F. Mullerried Ciencia, Mex. 12 1-2 1952: p. 36 f. 1-4.

Praeradiolites in the Cretaceous of Algiers, Emberger (1).

Radiolitis and their stratigraphy in the Pyrenees, G. Astre; Radiolites and Whitfield's Rudist types from the Cretaceous Jamaica, L. J. Chubb.

CARDIACEA

†Cardium (Granocardium) lowei sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955: p. 119 pl. 19 f. 1-8; C. (Acanthocardia) andrussovi latecostatum subsp. nov. p. 150 pl. iv f. 1-3; C.

(Limnocardium) solense p. 151 pl. iv f. 5, 6; C. (L.) lamellosum p. 151 pl. iv f. 9, 10; C. (L.) plicatoforme p. 152 pl. iv f. 7; C. (L.) subplicatofittoni p. 152 pl. iv f. 13; C. (L.) rotundum p. 153 pl. iv f. 8; C. (L.) kolesnikovi p. 153 pl. iv f. 11 spp. nov. Miocene, Czechoslovakia, J. Senes Geol. Práce Slovensk. Akad. 40 1955; C. (Replidacna) krambergeri sp. nov. Pannon (Tertiary) Austria, F. Sauerzopf Burgenland. Heimat Bl. 14 1 1952: p. 11 pl. 2 f. 10-11.

Clinocardium uchidai sp. nov. Akkeshi Bay, Japan, T. Habe Publ. Akkeshi mar. biol. Sta. No. 4 1955: p. 11 pl. 2 f. 5, 6.

†Laevicardium fiski sp. nov. from well off mouth of the Mississippi River, H. G. Richards Natul. Nat. Acad. Nat. Sci. Philad. No. 263 1954 p. 2 f. 1, 2.

†Limnocardium fittoniforme p. 231 pl. xxxiii f. 1-8; L. gilleti p. 231 pl. xxxiii f. 9-16 spp. nov. Miocene, Czechoslovakia, J. Seneš Geol. sbornik. Slovensk. Akad. 3 3-4 1953.

†Nemocardium (Pratulum) proterothetidis sp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 64 pl. 3 f. 16, 17 [nom. nud. loc. cit. 77 1954: p. 57].

†Serripes expansus sp. nov. Miocene, Tochigi Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 66 pl. 4 f. 1, 2.

Tridacna crocea biological note, fig., R. D. Purchon; T. gigas produces giant pearls, Lay Yoe Soey.

†Vasticardium tochigiensis sp. nov. Miocene, Tochigi Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 64 pl. 3 f. 13-15; V. (Vasticardium) submaculosum p. 60 pl. 4 f. 18; V. (Regozara) praecygnorum p. 61 pl. 4 f. 12 spp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955 [nom. nud. loc. cit. 77 1954: p. 57].

VENERACEA

Callista brevisiphonata chemical contents, M. Asano.

†Callista pseudobrevisiphonata sp. nov. Miocene, Tochigi Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 68 pl. 4 f. 4-6; C. chinensis takagii subsp. nov. Miocene, Japan, K. Masuda Trans. Palaeont. Soc. Japan N.S. 20 1955 p. 121 pl. 19 f. 7.

†Clementia japonica sp. nov. Miocene, Japan, K. Masuda Trans. Palaeont. Soc. Japan N.S. 20 1955 p. 121 pl. 19 f. 8.

† Dosinia ovata sp. nov. p. 81, pl. 13 f. 6a-c; D. (Kaneharaia) kaneharai ouchiensis subsp. nov. p. 82, pl. 13 f. 1-5. Tertiary, Japan, S. Kanno Trans. Palaeont. Soc. Japan N.S. 19 1955.

†Lamelliconcha kawadai sp. nov. Miocene, Fukushima Prefecture, Japan, S. Aoki Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 36 pl. 2 f. 1-15, 22.

Meretrix lamarckii biology, K. Shigeno.

Sunetta americana sp. nov. Buenos Aires Province, M. D. Jurado Physis 20 58 1951: p. 249 f. 1-6.

T. semidecussata as food, J. Krenmayr.

Tapes decussatus, spermatogenesis, Heldt.

Tawera rosa sp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 41 pl. 5 f. 41, 42.

†Tawera incurvilamellata sp. nov. Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 68 pl. 3 f. 18, 19 [nom. nud. loc. cit. 77 1954: p. 57].

Venerupis chemical contents, Y. Kado; Venerupis philippinarum spawning season, J. Yasuda.

Venus mercenaria population, Dow & Wallace; V. mercenaria the quahaug, fig., Medcof & MacPhail.

MACTRACEA

†Ervilia (?) minima sp. nov. Miocene, Czechoslovakia, J. Seneš Geol. sbornik Slovensk. Akad. 3 3-4 1953: p. 233 pl. xxxiii f. 19-20.

†Lutraria sanna longa subsp. nov. Miocene, Czechoslovakia, J. Senes Geol. Práce Slovensk. Akad. 40 1955: p. 154 pl. v f. 5-6.

Schizothaerus keenae n. sp. for S. nuttalli [not Conrad], Kuroda & Habe Illust. Catal. Jap. Shells No. 4 1950 p. 30.

Spirula solidissima the bar clam (fig.) in the maritime Provinces of Canada, Medcof & MacPhail.

TELLINACEA

Amphidesma ventricosum Gray, studies on the species, R. M. Cassie.

Leptomya retiaria aucklandica subsp. nov. New Zealand, A. W. B. Powell Cape Exped. Ser. Bull. No. 15 1955: p. 38 pl. 1 f. 2.

†Linearia lirulifera sp. nov. Oretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 264–B 1954: p. 34 pl. 7 f. 22–25.

†" Macoma" [?] prima sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 106 pl. iv f. 24, 25.

†Macoma ishimoriensis sp. nov. Miocene, Fukushima Prefecture, Japan, S. Aoki Sci. Rep. Tokyo Kyoiku Daig. 3 C 1954: p. 37 pl. 2 f. 21.

Nuttallia ezonis sp. nov. Akkeshi Bay, Japan, Kuroda et Habe Publ. Akkeshi mar. biol. Sta. No. 4 1955: p. 17 pl. 1 f. 12, 13.

Tellina perrieri, gibba and parvula of Bertin 1878 notes, T. Habe (7); Tellina (Eurytellina) eburnea askoyana subsp. nov. Champerico, Guatemala to Pitas Bay, Panama, Hertlin & Strong Bull. Amer. Mus. nat. Hist. 107 2 1955 p. 197 pl. 3 f. 3 13-15, 20, 21, 23; Tellina (Moerella) tabogensis Salisbury, type loc. dredged in 30 fms. Panama Bay st. 2799, San Filepe, G. of Calif. to Santa Elena, Ecuador, Hertlein & Strong.

†Tellina hamadai sp. nov. p. 122 pl. 19 f. 10, 11.—T. notoensis sp. nov. p. 122 pl. 19 f. 12, Miocene, Japan, K. Masuda Trans. Palaeont. Soc. Japan N.S. 20 1955; T. buboana sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-E 1955: p. 121 pl. 20 f. 12-14.

SOLENACEA

Ensis tropicalis sp. nov. valve from South Passage, Pearl Islands, Panama, Hertlein & Strong Bull. Amer. Mus. nat. Hist. 107 2 1955 p. 203 pl. 3 f. 34, 35; Ensis directus captured by Polinices duplicatus, H. J. Turner, jr.

Panope floridanae Heilprin, fig., H. E. Vokes.

Solecurtus leone sp. nov. off Brush Island, 15 miles S. of Ulladulla, N.S.W. Australia; 40–45 fathoms, L. Woolacott Proc. R. zool. Soc. N.S.W. 1952–53 1954: p. 39 pl. 3 f. 3–5.

SAXICAVACEA

†Panope rhomboidea sp. nov. Tertiary, Fukushima Prefecture, Japan, K. Hirayama Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 108 pl. 3 f. 12, 13.

MYACEA

†Aphanaia? orbirugata sp. nov., Posidonia orbirugata Harrington, 1947 (nomen nudum), p. 27. Permian, Argentina, H. J. Harrington J. Paleont. 29 1 1955: p. 121 pl. 24 f. 7.

†Corbula depressa minima var. nov. Upper Noric Trias Val Solda, Italy, S. Conti Mem. Carte geol. ital. 30 1954 p. 114 pl. iii f. 10a-b, text-fig. 39; C. adelaidensis nom nov. for C. equivalis N. H. Woods, 1931 (non Philippi, 1836) Pliocene, Australia, N. H. Ludbrook Trans. roy. Soc. S. Aust. 78 1955: p. 78.

†Cowperesia gen. nov. genotype C. anceps (Reed) Permian, J. C. Mendes Bol. Fac. Filos Ciênc. S. Paulo 145 8 1952 pp. 4 86 pl. ii f. 3a, b, pl. iv f. 2a-5b.

Mya arenaria and Molgula manhattensis, a case of symbiotic relationship, fig., F. A. Aldrich; M. arenaria, Pismo clam census, J. E. Fitch (1); M. arenaria, a note on Orkney specimens, R. Rendall (1); M. arenaria, on the mechanism of the apparatus for opening the valves, E. R. Trueman.

Myalina magna an emended description, G. M. Bennison.

†Myophoria orbicularis nana p. 94 pl. ii f. 10a, b text-fig. 21; M. ovata minuta p. 95 pl. ii f. 11 text-fig. 22; vars. nov. Upper Norico Val Solda, S. Conti Mem. Carte geol. ital. 30 1954.

†Myophoriopis (Pseudocorbula) sancti mametis p. 100 pl. ii f. 18 text-fig. 27; M. (P.) subpyramidata p. 102 pl. ii f. 20, 21 text-figs. 28-30; M. (P.) repossii p. 103 pl. 11 f. 22 text-fig. 31; spp. nov. Upper Norico, S. Conti Mem. Carte geol. ital. 30 1954. †Parmicorbula moodiana sp. nov. Pepper Shale (Cretaceous) Lemmon Co., Texas, L. W. Stephenson Prof. Pprs. U.S. geol. Surv. 243E 1953 pp. 57 63 pl. 13 f. 21-23.

Patamocorbula gen. nov. genotype Corbula amurensis Schrenck, T. Habe Zool. Mag. Tokyo 64 1955 p. 271.

GASTROCHAENACEA

Rocellaria (Gastrochaena) cuneiformis general structure and biology, R. D. Purchon (1).

ADESMACEA

†Bankia (Bankiella) minima hemicalix subsp. nov. Tortonian, Austria, A. F. Tauber Wiss. Arb. Burgenland No. 3 [1954]: p. 25 pl. 1 f. 1a, b, c.

Martesia striata tolerance to low salinity, R. Nagabhusham.

Pholadidea (Penitella) chishimana sp. nov. Akkeshi Bay, Japan, T. Habe Publ. Akkeshi mar. biol. Sta. No. 4 1955: p. 23 pl. 7 f. 8, 9.

Teredo and cellulase concentration, L. J. Greenfield; T. tritubulata description of valves, P. Rancurel; T. thomsoni and T. lieberkindi changes in plates due to growth, P. Rancurel (1); T. arenaria systematic standing, G. F. Roch; T. (Coeloteredo) bayeri p. 130 fig. 4; T. (Dactyloteredo) juttingae p. 135 fig. 6 spp. nov. Rhiouw Archipelago, East Indies, F. Roch (1) Zool. Meded. 34 1955.

PANDORACEA

†Allorisma inflectoventris sp. nov. Permian, Argentina, H. J. Harrington J. Paleont. 29 1 1955: p. 126 pl. 24 f. 8.

Myochama anomioides Stutchbury 1830 in S. Australia, Pt. Adelaide, B. C. Cotton (5).

Myodora soyae p. 25 pl. 4 f. 18; M. teramachii p. 26 pl. 4 f. 14-16; M. japonica p. 27 pl. 4 f. 4-6; M. yokoyamai p. 28 pl. 4 f. 7-9; spp. nov. Japan, T. Habe Illust. Catal. Jap. Shells No. 4 1950.

†Myodora ikebei sp. nov. Pleistocene Japan, T. Habe Illust. Catal. Jap. Shells No. 4 1950 p. 30 pl. 4 f. 17. †Oliveiraia gen. nov. genotype Thracia pristina Reed 1929, status of Paresia J. C. Mendes Bol. Fac. Filos. Cienc. S. Paulo 1954 175 (10) pp. 104 pl. 3 f. 8

Pandora inaequivalvis (L.) on the habits, figs., M. F. and J. A. Allen.

†Pholadomya quenstedti 'n. nom.' [sp. nov.] Retia Val Solda, Lugano, Italy, S. Conti Mem. Carte geol. ital. 30 1954 p. 153 pl. v f. 10 text-fig. 46; P. japonica Yokoyama subsp. of Cyrtopleura dilatata (Soulayet), Hatai & Nisiyama J. geol. Soc. Japan 56 654 1950 p. 100 text-fig.; P. (Procardia) ziergiebeli sp. nov. Callovian [Germany], A. Zeiss Neues Jb. Geol. Paläont. 11 1955: p. 499 f. la-c.

CLAVOGELLACEA

†Brechites (Warnea) yokoyamai sp. nov. Pliocene, Japan, T. Shikama Sci. Rep. Yokohama Univ. Sect. II No. 3 1954: p. 63 pl. 3 f. 1a-d.

†Gresslya agassizi pseudopolonica subsp. nov. W. Wetzel Palaeontographica A105 3-6 1954 p. 149 pl. 22 f. 10.

Cephalopoda

RECENT

Abralia redfieldi sp. nov. at surfaces off Gun Cay, Bahamas, G. L. Voss Bull. mar. Sci. Gulf and Caribbean 5 2 1955 pp. 81 99 f. 4A-H.

Acanthosepion pageorum sp. nov. Queensland, Australia, T. Iredale Aust. 2001. 12 1954: p. 76 pl. 4 f. 7-9.

Amplisepia parysatis sp. nov. Western Australia, T. Iredale Aust. zool. 12 1954: p. 71 pl. 4 f. 1, 2.

Arctosepia rhoda sp. nov. Point Cloates, Mid-western Australia, T. Iredale Aust. zool. 12 1954: p. 75 pl. 4 f. 10-12.

Blandosepia bartletti sp. nov. Louisiade Archipelago, Australia, T. Iredale Aust. zool. 12 1954: p. 67 pl. 5 f. 15, 16.

Cephalopoda of the Eastern Mediterranean, Distribution and list of species, W. J. Rees.

Eledone affinities with Octopus, A. da F. Sacarrão.

Fiscisepia subgen. nov. for Acanthosepion ellipticum adjacens Iredale, Iredale Aust. Zool. 4 1926, in text p. 239 pl. xxxv f. 56.

Glyptosepia cultrata [Hoyle] = G. macilenta Iredale a synonym of the former, T. Iredale (4).

Heteroteuthis (Stephanoteuthis) atlantis sp. nov. off Puerto Tanamo, Cuba, in 640 fms. G. L. Voss Bull. mar. Sci. Gulf and Caribbean 5 2E 1955 pp. 81 93 figs. 3A-F.

Illex coindeti secretion in the brain, H. Haefelfinger.

'Kraken' the legendary giant cephalopod, J. Allan.

Lligo opalescens sexual behaviour and spawning, J. McGowan; Loligo pealii experiments on muscular activity, G. de Villafranca.

Mesembrisepia ostanes sp. nov. Stanley, Tasmania, T. Iredale Aust. zool. 12 1954: p. 69 pl. 4 f. 5, 6.

†Nautilus and Nautiloid coiling, figs., R. H. Flower.

Octopodia Schneider 1784, suppression, Opinion 233.

Octopus chemical contents, Asano & Sato; Octopus vulgaris learning capacity, B. Boycott; Octopus vulgaris—memories controlling attacks on food, B. Boycott & J. Z. Young; Octopus reactions, Boycott & Young; O. macropus Risso on the larvae and late larval stages, figs., W. J. Rees (1); Octopus affinities with Eledone, G. da F. Sacarrão

Ocythoe tuberculata Raf., occurrences in California, measurements, etc., S. S. Berry.

Ommastrephes sloani pacificus studies on the tropomyosin, K. Yoshimura.

Opisthoteuthis californiana Berry, a male taken off Humboldt Co., California, described and figured, S. S. Berry.

Rossia (Austrorossia) antillensis sp. nov. off Sagua la Grande, Cuba, 280–300 f. pp. 81 86 f. 1A–E; Rossia brachyura Verrill, figured, G. L. Voss Bull. mar. Sci., Gulf and Caribbean 5 2 1955.

Sepiella melwardi sp. nov. Melville Island, Northern Territory, Australia, T. Iredale Aust. zool. 12 1954: p. 78 pl. 5 f. 1-6. Solitosepia genista sp. nov. Broome, Western Australia, T. Iredale Aust. zool. 12 1954: p. 66 pl. 5 f. 17, 18.

Spirula live specimens taken and new light on biology of the species, figs., A. F. Braun.

Spirula spirula L. animal cast up on Boynton Beach, Florida, T. L. McGinty (1).

Spisula solidissima isolation of blood anticoagulant, L. Thomas.

Tetracheledone type spinicirrus gen. et sp. nov. 145-190 fms. off Matanzas, G. L. Voss Bull mar. sci. Gulf and Caribbean 5 2 1955 pp. 81, 107 f. 5A-H.

FOSSIL

†Acanthoceras athabascense sp. nov. Cretaceous, Canada, P. S. Warren & C. R. Stelek Rep. Res. Counc. Alberta No. 70 1955: p. 71 pl. 6 f. 5, 6 pl. 7 f. 1, 3, 4; pl. 8 f. 1-3; pl. 9 f. 2; A. johnsonanum sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-C 1955: p. 58 pl. 4 f. 14-17.

†Aktubonautilidae fam. nov. p. 100 Aktubonautilus gen. nov. genotype: A. eruciformis sp. nov. p. 101 pl. x f. 1, Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50

†Aktubonautilus gen. nov. Aktubonautilidae fam. nov. [q. v.] V. E. Ruzhentzev & V. N. Shimanskii.

†Allotrioceratidae fam. nov. p. 356 Allotrioceras gen. nov. p. 357 genotype Allotrioceras bifurcatum sp. nov. p. 358 pl. 34 f. 3-19, text-f. 6A-H. Palaeozoic, America, R. H. Flower, J. Paleont. 29 3 1955.

†Ammonites published in Vol. 2 Bull. Zool. Nomen. Additions and rejections, **Direction 14**.

Ammonites cordatus J. Sowerby Designation of lectotype, Opinion 235; A. cornuclianus, proposal to place on the Official List of Specific Names in Zoology the specific name cornuclianus d'Orbigny, 1841 as published in the above combination, R. Casey & C. W. Wright, Bull. zool. Nom. 9 1954: p. 280; A. mammillatus Schlotheim, 1813, proposal to designate a neotype, R. Casey (2); Ammonites rogoznikensis Zeuschner, 1868, (specific name of type

species of Aspidoceras Zittel, 1868) proposal to place the specific name on the Official List of Specific Names in Zoology, W. J. Arkell (4); Ammonites royerianus proposal to suppress for the purposes of the Law of Priority, but not for those of the Law of Homonymy, the specific name royerianus d'Orbigny, 1841 as published in the above combination, R. Casey & C. W. Wright, Bull. l. zooNom. 9 1954: 279; 280.

†Ammonoidea Zittel 1884 an addition of names to Official list and type genus, W. J. Arkell; †Ammonoidea, suppression of five early and obsolete generic names, Opinion 304.

†Apogonoceras gen. nov. p. 133 Triboloceratidae, genotype: A. remotum sp. nov. p. 134 pl. xi f. 8, Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

†Aptychus Meyer, 1831—Aptychus Deshayes, 1845, in Lamarck Hist. Anim. sans Vertèbr. (ed. 2) 11: 228 (an emendation of Aptychus Meyer, 1831) proposed suppression by the Plenary Powers, W. J. Arkell (4).

†Argonauta kagana p. 341 pl. 1 f. 1a, b, c—A. kasataniensis p. 343 pl. 1 f. 4, 5, 6 spp. nov. Late Miocene or early Pliocene, Ishikawa Prefecture, Japan, Y. Kaseno Sci. Rep. Kanazawa Univ. 3 2 1955.

†Arieticeras Quenstedt 1883 rejection of names and addition to list of Arieticeras Seguenza 1885, Opinion 337.

†Articheilus gen. nov. p. 89 Mosquoceratidae genotype: A. luxuriosum sp. nov. p. 90 pl. viii f. 7; pl. ix f. 1. Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954.

†Aspidoceras Zittel, 1868 proposal to place the generic name on the Official List of Generic Names in Zoology, W. J. Arkell (4); A. (Euaspidoceras) rotundatum sp. nov. p. 187 pl. 30 f. 5-16, O. Haas Bull. Amer. Mus. nat. Hist. 103 1 1955; A. (Euaspidoceras) hermonis nom. nov. for A. perarmatum Sow. sp. Noetting, O. Haas Bull. Amer. Mus. nat. Hist. 108 1 1955 179.

†Aturia c.f. alabamensis (Morton) Eocene Cyrenaica, figs., O. Haas; Aturia in the Miocene of Japan, Kobayashi & Mastani; A. minoensis sp. nov. Miocene, Hazamagahora, central Japan, T. Kobayashi Jap. J. Geol. Geogr. 25 1-2 1954: p. 36 pl. 5 f. a-d; text-f. 1; A. peruviana—first record from Costa Rica, A. K. Miller, Furnish & Paguaga; A. stansburiensis sp. nov. Oligocene, South Australia, M. F. Glaessner Rec. S. Aust. Mus. 11 4 1955: p. 357 pl. 34 f. 1a, b.

†Aturoidea brunnschweileri sp. nov. Mid Eocene, Giralia Range, North West Cape area, Western Australia, Glenister, Miller & Furnish, J. Paleont. 30 3 1956: p. 500 pl. 55 f. 1, 2, 5, 6 pl. 56 f. 1, 2.

†Aulacostephanus Tornquist 1896, Designation of type species, Opinion 302.

†Bactrites sempiternus p. 72 pl. viii f. 9, 10—B. ? ultuganensis p. 74 pl. VIII f. 8 spp. nov. Permian, southern Ural, U.S.S.R., V. N. Shimanskii Trav. Inst. Paléont. Acad. Sci. URSS 44 1954.

†Balearites 'gen. nov.' Crioceras baleare Nolan 1894 taken as type Cretaceous, Majorca, S. S. Sarkar C.R. Soc. géol. Fr. 4 1954 p. 98.

†Baltoceras minor sp. nov. Chazyan, North America, R. H. Flower J. Paleont. 29 5 1955: p. 810 pl. 80 f. 5-6.

†Barroisiceras haberfellneri hemitissotiformis subsp. nov. Cretaceous, Austria, B. Plöchinger S.B. öst. Akad. Wiss. Abt. I 164 4-5 1955: p. 203 pl. 1.

†Basleonautilus gen. nov. Aktubonautilidae genotype: Pleuronautilus dyadicus Haniel, 1915. Permian, Southern Ural., V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954: p. 102.

†Belemnitomimus gen. nov. Bactritoidea p. 91 type species B. palaeozoicus sp. nov. p. 92 pl. xii f. 1, 2. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. Paléont. Acad. Sci. URSS 44 1954.

†Belennitella praecursor media p. 497 Pl. 57 f. 4a-b, pl. 56 f. 2—B. praecursor mucronatiformis p. 497 pl. 56 f. 5a-d vars. nov. Cretaceous, U.S.S.R. J. A. Jeletzky J. Paleont. 25 3 1955.

†Belemnites listeri proposal to suppress the specific name listeri Mantell, 1822, as published in the above combination, and to place it on the Official Index of Rejected and Invalid Specific Names in Zoology—B. minimus proposed to place the specific name minimus, Miller (J. S.), 1826, as published in the above combination, on the Official List of Specific Names in Zoology, H. Swinnerton Bull. zool. Nom. 9 1954: 284-285.

† Belemnopsis (or Hibolites) compressa sp. nov. p. 163 pl. xv f. 28—B. kuntkotensis Waagen puenensis var. nov. p. 164 pl. xiv f. 4, 9, 12, 13, 15, pl. xv f. 24, 30, 31, 34, 36, 37, 38, 39 pl. xvi f. 5, 11, 17 Jurassic, Central New Caledonia, J. Avias Sciences de la Terre, Nancy, 11-21953.

†Belosepiella De Alessandri 1905 "Type species here designated" Belosepiella cossmanni De Alessandri, D. Cassey Proc. Malac Soc. London 31 116.

†Belosepiella parisiensis De Alessendri and its occurrence in the English Eocene, fig., D. Curry.

†Burenoceras cornucopiaforme sp. nov. Oneota dolomite, Ordovician, Minnesota, U.S.A., L. Powell Sci. Bull. Sci. Mus. St. Paul Inst. 1 1935: p. 70 pl.8. f. 1-3.

†Beudanticeras revoili elegans nom. nud. Cretaceous, Egypt, I. G. E. Mahmoud Bull. Inst. Egypt 36 1 1955: p. 224.

†Borissiakoceras orbiculatum sp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-C 1955: p. 64 pl. 6 f. 1-4.

†Campbelloceras overmani sp. nov. Ordovician, North America, R. Heller Bienn. Rep. Mo. Bur. Geol. Min. (2) 35 1954: p. 41 pl. 17 f. 6-8.

†Canavarella helvetica sp. nov. Jurassic, Switzerland, P. L. Maubeuge Schweiz. palaeont. Abh. 71 1955: p. 26 pl. 11 f. 1, a, b, c.

†Cephalopoda. Trails and tentacular impressions of cephalopods in the Corryville of Cincinnati, America, R. H. Flower (3).

†Ceratites nodosus optimus p. 13 pl. 6 f. 37; C. nodosus subpostspinosus p. 14 pl. 6 f. 38 subspp. nov.; C. meissnerianus p. 16 pl. 3 f. 18 sp. nov. Triassic, Germany, **H. Penndorf** Abh. senckenb. Naturf. Ges. No. 484 1951.

†Cimonia tenuicosta sp. nov. Maestrichtian Miria marl, Cardabia Range, North West Cape area, Western Australia, Glenister, Miller & Furnish, J. Paleont. 30 3 1956: p. 495 pl. 53 f. 1-9.

†Collignoniceras gen. nov. Acanthoceratidae p. 361, genotype: C. hammersleyi p. 361 f. 7-9; C. peregrinator p. 364 f. 10, 11 spp. nov. C. vinassai Venzo = Mortoniceras vinassai Venzo 1936?—Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955.

†Cheloniceras Hyatt, 1903, proposal to designate as type-species Ammonites cornuclianus d'Orbigny, 1841 and to place the generic name Cheloniceras on the Official List of Generic Names in Zoology, R. Casey & C. W. Wright Bull zool. Nom. 9 1954: p. 280.

†Caenisites Buckman, 1925, proposed suppression of the name, W. J. Arkell & D. T. Donovan.

†Clymenia Taxonomy and Nomenclature, O. H. Schindewolf.

†Condraoceras ellipsoidale sp. nov. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954: p. 129 pl. xii f. 4.

†Ctenobactrites mirus sp. nov. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. Paléont. Acad. Sci. URSS 44 1954: p. 82 pl. ix f. 1, 2.

† Dentoceras gen. nov. Dentoceratidae fam. nom. q. v., V. E. Ruzhentzev & V. N. Shimanskii.

†Dentoceratidae fam. nov. p. 141; Dentoceras gen. nov. p. 141; genotype: D. magnum p. 142 pl. xv f. 1; D. ultuganense p. 142 pl. xiv f. 1; D. latum p. 143 pl. xiv f. 2 spp. nov. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

† Docidoceras liebi sp. nov. Jurassic, Switzerland, P. L. Maubeuge Schweiz. palaeont. Abh. 71 1955: p. 42 pl. 9 f. 1a-d, 2.

† Dolorthoceras sociale—from the Maquoketa at Graf, Iowa, P. Tasch (1).

† Douvilleiceras de Grossauvre, 1893, proposal to designate a type species, R. Casey (2); Douvilleiceras coronatum proposal to place on the Official List of Specific Names in Zoology the specific name coronatum Rouchadzé, 1932 as published in the above combination, R. Casey & C. W. Wright Bull. zool. Nom. 9 1954: p. 280.

†Dunveganoceras clowi p. 66 pl. 9 f. 1, 3; D. hagei p. 69 pl. 5 f. 1 spp. nov. Cretaceous, Canada, P. S. Warren & C. R. Stelck Rep. Res. Counc. Alberta No. 70 1955.

†Endolobus clorensis sp. nov. Clore limestone, Carboniferous, Kentucky, U.S.A., C. Collinson J. Palaeont. 29 1 1955: p. 178 text-f. 1-3.

†Eomadrasites gen. nov. Kossmaticeratidae p. 132. Type species: Eomadrasites nipponicus p. 135 pl. x f. 3; Text-f. 7a-e; E. subnipponicus p. 137 pl. x f. 2a-b; Text-f. 9a-e spp. nov. Cretaceous, Hokkaido, Japan, T. Matsumoto Jap. J. Geol. Geogr. 26 1-2 1955.

†Eotetragonites breistrofferi nom. nud. Cretaceous, Egypt, I. G. E. Mahmoud Bull. Inst. Egypt. 36 1 1955: p. 223.

†Erioliceras gen. nov. Erioliceratidae fam. nov. q.v., E.. van Hoepen (2).

†Erioliceratidae fam. nov. Ammonoidea type genus: Erioliceras gen. nov. p. 378 genotype: E. tenuis p. 378 f. 32; E. elegans p. 379 f. 33; E. humilis p. 380 f. 34 spp. nov. Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955.

†Foodiceras whnneiforme [sic] sp. nov. p. 184, as Foordiceras whyneiforme pl. 1 f. 1 (= Foordiceras wynneiforme p. 185) Permian, Japan, Hayasaka & Ozaki Sci. Rep. Kanazawa Univ. 3 1 1955.

†Gardeniceras subgen. nov. of Hauericeras de Grossouvre, subgenotype: Ammonites gardeni Baily, 1855, Cretaceous, Japan, T. Matsumoto & I. Obata Mem. Fac. Sci. Kyushu Univ. 5D 3 1955: p. 134.

†Gattendorfia stummi sp. nov. Carboniferous, U.S.A., A. K. Miller & H. F. Garner Contr. Mus. Geol. Univ., Michigan 12 8 1955: p. 125 pl. 6 f. 6, 7.

†Gauthiericeras de Grossouvre, type genus of Gauthiericeratidae fam. nov. q. vide, E. van Hoepen (1).

†Gauthiericeratidae fam. nov. p. 367 Ammonoidea type genus: Gautheiriceras de Grossouvre, emend. v. Hoepen genotype: G. margae de Gros.; G. libertae p. 372 f. 23-25; G. obesum p. 374 f. 26-28; G. planulatum p. 374 f. 29-31 spp. nov. Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955.

†Glyptophiceras japonicum sp. nov. K. Nakazawa and D. Shimizu Trans. Palaeont. Soc. Japan N.S. 17 1955 p. 16 pl. 3 f. 1, 2.

†Gzheloceras gen. nov. Gzheloceratidae fam. nov. q. vide, V. E. Ruzhentzev & V. N. Shimanskii.

†Gzheloceratidae fam. nov. Nautiloidea p. 74; Gzheloceras gen. nov. p. 75 type species G. uralense p. 78 pl. vii f. 4; G. sholakense p. 76 pl. vii f. 3; G. ellipsoidale p. 79 pl. vii f. 5-8; G. biangulare p. 81 Pl. viii f. 1, 2 spp. nov. Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954.

†Hammatoceras tuberculata p. 86, pl. 8 f. 2 holotype; pl. 7 f. 1; pl. 9 f. 1; text-f. 3-5; H. subtile p. 89, pl. 7 f. 3; pl. 8 f. 4; pl. 9 f. 3; H. hosourense p. 91, pl. 9 f. 5, 6 (holotype); pl. 7 f. 2, 5, 6; text-f. 8-14 spp. nov. Jurassic, Japan, T. Sato Jap. J. Geol. Geogr. 25 1954.

†Harpoceras Waagen 1869. Designation of type species, Opinion 303.

†Hauericeras (Gardeniceras) subgen. nov. Cretaceous, Japan, T. Matsumoto & I. Obata Mem. Fac. Sci. Kyushu Univ. 5D 3 1955: p. 134; Hauericeras "is not a typical member of the Desmoceratidae (s.s.) nor of the Puzosiidae and should be placed in the subfamily Hauericeratinae", Matsumoto & Obata.

†Hecticoceras (Sublinuloceras) socium sp. nov. p. 42 pl. 6 f. 26-38 pl. 7, f. 1-10; H. (Brightia) syriacum p. 56 pl. 8 f. 36-54 pl. 9 f. 1-14; H. (Putealiceras) solare p. 73 pl. 11 f. 33-57 pl. 12 f. 1-12; spp. nov. 0. Haas Bull. Amer. Mus. nat. Hist. 108 l 1955.

†Hemibactrites gen. nov. Bactritoidea p. 75 type species: H. ellipsoidalis p. 77 pl. viii f. 1-5; H. primus spp. nov. p. 76 pl. VIII f. 6, 7. Permian, Southern Ural, U.S.S.R., V. N. Shimanskii Trav. Inst. Paléont. Acad. Sci. URSS 44 1954.

†Hemiliroceras gen. nov. p. 125. Liroceratidae, genotype H. inflatum p. 126 pl. xii f. 2; H. zhiltauense p. 127 pl. xii f. 3 spp. nov. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

†Heurekoceras gen. nov. Gzheloceratidae, p. 82 type species: H. notabile sp. nov. p. 83 pl. viii f. 3. Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954.

†Hoplites Neumayr 1875, Validation of name, Opinion 353.

†Hudsonoceras gen. nov. p. 351 Manchuroceratidae, genotype Hudsonoceras aristos sp. nov. Palaeozoic, America, R. H. Flower J. Paleont. 29 3, 1955: p. 352, pl. 32 f. 1-4, pl. 33 f. 1-13, text-f. 3.

†Hulenites gen. nov. Kossmaticeratidae. Type species: Puzosia reesidei Anderson, 1938, also includes Puzosia diadema Anderson, 1938, P. jimboi Anderson, 1938, P. onona Anderson, 1938, T. Matsumoto Jap. J. Geol. Geogr. 26 1955.

†Hypophylloceras (Hyporbulites) aegyptiacum nom. nud. Cretaceous, Egypt, I. G. E. Mahmoud Bull. Inst. Egypt. 36 1 1955: p. 223.

†Izumonauta gen. nov. Argonautinae, genotype: Izumonauta lata sp. nov. Miocene, Shimone Prefecture, Japan, T. Kobayashi, Jap. J. Geol. Geogr. 25 1-2 1954: p. 31 pl. 3 f. 1-4.

†Kabylites 'gen. nov.' Ammonitacea Bochianites superstes Peru 1910, taken as type, M. D. Delga C.R. Soc. géol. Fr. 7 1954: p. 136 text-f. B.

†Kaiparaites gen. nov. Kossmaticeratidae, Type species: Madrasites multicostatus Marshall, 1926; Kossmaticeras multicostatus (Marshall), K. regularis (Marshall), K. sulcatus (Marshall), K. natalensis (Crick MS) (Spath), K. acuticostatus (Crick MS) (Spath) and probably K. buddhaicus (Kossmat) and (?) K. faku (van Hoepen) are referred to this genus. Cretaceous, T. Matsumoto Jap. J. Geol. Geogr. 26 1–2 1955.

†Karapadites gen. nov. Kossmaticeratidae, Type species: Holcodiscus karapadensis Kossmat, 1897, mono-

typic. Arrialoor group in India, Upper Campanian (?), T. Matsumoto Jap. J. Geol. Geogr. 26 1-2 1955.

†Kazakhstania americana sp. nov. Carboniferous, U.S.A., A. K. Miller & H. F. Garner Contr. Mus. Geol. Univ. Michigan 12 8 1955: p. 129 pl. 1 f. 3; pl. 6 f. 2, 3.

†Kingites shimizui sp. nov. Triassie, Kwanto Massif, Japan, S. Sakagami Sei. Rep. Tokyo Kyoiku Daig. 4 1955: p. 138 pl. 2 f. 2a-c.

†Kinoceras serenum sp. nov. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 44 1954: p. 132 pl. 111 f. 10-12.

†Koninckioceras from the Japanese Permian, I. Hayasaka.

†Kosmoceras Waagen 1869. Designation of type species, Opinion 303.

†Kossmaticeras japonicum (ex. Yabe MS.), "sp. nov." Cretaceous, Hokkaido and Saghalien, Japan, T. Matsumoto Jap. J. Geol. Geog. 26 1-2 1955: p. 150 pl. ix, f. 3a, b.

†Lamottoceras gen. nov. Endoceratidae p. 813 genotype: Lamottoceras ruedemanni sp. nov. p. 815 pl. 77 f. 3-13; pl. 78 f. 1-9, 11-13; L. nodosum sp. nov. p. 817 pl. 77 f. 2, 14, 15; Pl. 78 f. 10 Chazyan, North America, R. H. Flower J. Paleont. 29 5 1955.

†Leonardocheilus gen. nov. Mosquoceratidae, genotype: Temnocheilus inaequilaterale Miller et Youngquist, 1949. Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954: p. 92 pl. iii f. 8.

†Lepthoplites topolianus sp. nov. Cretaceous, Serbia, M. Z. Andelković Ann. géol. Pén. Balkan 22 1954: p. 65 pl. vii f. 4.

†Lewisiceras australis sp. nov. Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955: p. 364 f. 12-14.

†Ludwigia pseudocrassa p. 27 pl. 4 f. 4a, b; L. reicheli p. 28 pl. 5 f. 1a-c; L. betznauensis p. 28 pl. 4 f. 3a-c spp. nov. Jurassic, Switzerland, P. L. Maubeuge Schweiz. palaeont. Abh. 71 1955.

†Mariceras gen. nov. p. 138 Seyphoceratidae, genotype: M. ferum sp. nov. p. 139 pl. xiii f. 6, 7. Permian, South-

ern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

†Marshallites gen. nov. p. 119 Kossmaticeratidae. Type species: Marshallites compressus 'sp. nov.' p. 123, pl. 8 f. 1a, b, 2; text-f. 1, 2. "Listed by Matsumoto, 1942 p. 193 and elsewhere as Maorites compressus (nom. nud.)", Matsumoto Jap. J. Geol. Geogr. 26 1-2 1955.

†Marshallitinae subfam. nov. Kossmaticeratidae, for Marshallites, nov. Holcodiscoides Spath, 1922, Eogunnarites Wright & Matsumoto, 1954, Hulenites nov., Eomadrasites nov., T. Matsumoto Jap. J. Geol. Geogr. 26 1955: p. 119.

†Mayaitidae Spath on proposed protection of family-group, W. S. Arkell (1).

†Megatyloceras Humphrey, 1949 proposal to place the generic name on the Official List of Generic Names in Zoology, R. Casey & C. W. Wright Bull. zool. Nom. 9 1954: p. 280.

†Menuites japonicus p. 158 pl. 31 f. 1a-c, 2a-d, 3a-b; pl. 33 f. 2a-d, 3a-c; text-f. 4a-b, 5a-c; M. naibutiensis p. 164 pl. 33 f. 1a-d; M. pusillus p. 165 pl. 32 f. 1a-d, 3a-c, 4a, b spp. nov. Cretaceous, Hokkaido and Saghalien, Japan, T. Matsumoto Mem. Fac. Sci. Kyushu Univ. 5D 3 1955.

†Merocanites drostei sp. nov. Carboniferous, America, C. W. Collinson J. Paleont. 29 3 1055: p. 434 pl. 45 f. 1, 2.

†Mesnaquaceras gen. nov. Michelinoceratidae p. 820 genotype: Mesnaquaceras curviseptatum sp. nov. p. 821 pl. 79 f. 7-8 Chazyan, North America, R. H. Flower J. Paleont. 29 5 1955.

†Metacoceras subpiszovi p. 57 pl. iv f. 1; M. parartiense p. 61 pl. iv f. 5, pl. v f. 1-3; M. kruglovi p. 63 pl. v f. 4; M. orthogonium p. 64 pl. vi f. 1, 2—M. subquadratum p. 66 pl. vi f. 3, 4; M. altilobatum p. 67 pl. vi f. 5 spp. nov. Permian, southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954.

†Microbactrites gen. nov. Bactritoidea p. 93 type species Parabactrites scorobogatovae Shimansky, 1948; M. scorobogatovae (Shimansky) p. 94 pl. xi f. 1, 2—M. parvus sp. nov. p. 94 pl. xi f. 3. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. Paléont. Acad. Sci. URSS 44 1954.

†Mirabiloceras gen. nov. p. 360 Allotrioceratidae genotype Mirabiloceras multitubulatum sp. nov. p. 364, pl. 35 f. 1-9 text-f. 5. Palaeozoic, America, R. H. Flower J. Paleont. 29 3 1955.

†Mooreoceratidae fam. nov., V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 44 1954 : p. 119.

†Morphoceras multiforme nom. nov = Am. polymorphus d'Orb. 1846, p. 379 pl. 124 f. 1-4 only, non Quenstedt 1845, W. J. Arkell Palaeontogr. Soc. [Monogr.] 5 1955: p. 132 pl. xvi f. 1, 2 text-fig. 47, 50.

†Mosquoceras gen. nov. Mosquoceratidae fam. nov. q. v. V. E. Ruzhentzev & V. N. Shimanskii.

†Mosquoceratidae fam. nov. p. 84—Mosquoceras gen. nov. p. 85 type species Nautilus tschernyschewi Tzwetaev, 1888; M. simense p. 86 pl. viii f. 4; M. jakowlewi p. 87 pl. viii f. 5, 6 spp. nov. Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954.

†Muensteria Eudes-Deslongchamps, 1835, proposed suppression by the Plenary Powers, W. J. Arkell (4).

†Münsteroceras pergibbosum sp. nov. Carboniferous, U.S.A., A. K. Miller & H. F. Garner Contr. Mus. Geol. Univ. Michigan 12 8 1955: p. 140 pl. 2 f. 12, 13; pl. 7 f. 1, 2.

†Murrayoceras primum sp. nov. lower Chazyan, North America, R. H. Flower J. Paleont. 29 5, 1955: p. 811 pl. 77 f. 1 pl. 79 f. 1.

†Nanno n.sp. description without name. Chazyan, North America, R. H. Flower J. Paleont. 29 5 1955: p. 812 pl. 81 f. 6-7.

†Nautilus in the Miocene of Japan, T. Kobayashi & Mastani.

†Neodomatoceras gen. nov. p. 97 Domatoceratidae, genotype: N. rarum sp. nov. p. 98 pl. ix f. 3 Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954. Neograhamites (?) paulckei nom. nov. for Kossmaticeras theoboldianum Paulcke Ber. Naturf. Ges. Freiburg B 15 1907: p. 54 pl. 7 (xvi) f.1, 1a not Stoliczka, Matsumoto Jap. J. Geol. Geogr. 26 1–2 1955: p. 149.

†Neohibolites minimus submedius var. nov. Gault of Folkestone. Type in Brit. Mus., H. H. Swinnerton Palaeontogr. Soc. [Monogr.] 5 1955: p. 77 pl. xvii f. 19–21.

†Neolobites medeninensis sp. nov. Cretaceous, Tunisia, E. Basse Bull. Soc. Sci. nat. Tunis, 6 1954: p. 199 pl. 23 f. 1-11.

†Neothrincoceras gen. nov. Thrincoceratidae fam. nov. q. vide, V. E. Ruzhentzev & V. N. Shimanskii.

†Neposiella gen. nov. Ammonoidea, Cretaceous of Gebel Manzour (Moghara-Sinai-Egypt), I. G. El-din Mahmoud, C.R. Soc. géol. Fr. 14 1953 p. 286 [see also C.R. Soc. géol. Fr. 1952 p. 119, name without description].

†*Nybyoceras cryptum* **sp. nov.** Chazyan, North America, **R. H. Flower** J. Paleont. **29** 5 1955: p. 818 pl. 79 f. 2–3, 5–6; pl. 81 f. 3–4.

†Ochetoceras (Campylites) freboldi p. 85 pl. 14 f. 1–47 text figs. 1–4; O. (C.) evolutum p. 93 pl. 14 f. 48–51 pl. 15 f. 1–29, O. Haas Bull. Amer. Mus. nat. Hist. 108 l 1955.

†*Oiophyllites* should be placed in the subfamily Hauerceratinae with *Hauericeras*, Matsumoto & Obata.

† Omocyrtoceras gen. nov. Erioliceratidae fam. nov. Ammonoidea: genotype: O. errabundum p. 380 f. 35; O. mutabilis p. 381 f. 36 spp. nov. Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955.

†*Ophiceras iwaiense* sp. nov. Triassic, Kwanto Massif, Japan, S. Sakagami Sci. Rep. Tokyo Kyoiku Daig. 4 1955: p. 135 pl. 1 f. 1–9.

†Orthocerotoidea superfam. nov. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 44 1954: p. 95.

†Parammatoceras (?) liebi sp. nov. Jurassic, Switzerland, P. L. Maubeuge Schweiz. palaeont. Abh. 71 1955: p. 32 pl. 6 f. 1a-c. †Pararhiphaeoceras gen. nov. p. 107 Rhiphaeoceratidae, genotype: Temnocheilus multituberculatum Waagen var. tastubensis Kruglov, 1928; P. tastubense (Kruglov) nom. nov. p. 107 pl. x f. 5; P. aktastense p. 109 pl. x f. 6; P. incallidum p. 110 pl. x f. 7 spp. nov. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

†Parastenopoceras gen. nov. p. 98 Domatoceratidae, genotype: P. khvorovae sp. nov. p. 99 pl. ix f. 4, 5. Permian, Southern Ural, V. E. Ruzhnetzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954.

†Peripetoceras asselense sp. nov. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954: p. 132 pl. xii f. 5.

†Perisphinctes Waagen 1869. Designation of type species, Opinion 303; Perisphinctes (Properisphinctes) vicinus p. 127 pl. 19 f. 22-23, pl. 20 f. 1-18 P. (Prop.) filocostatus p. 130 pl. 20 f. 19-36; P. (Prop.) radiocostatus f. 137 pl. 21 f. 32-46; P. (Prop.) hermonis p. 146 pl. 23 f. 16-42; P. (Prop.) delicatulus p. 149 pl. 23 f. 43-60 pl. 24 f. 1-18 spp. nov., O. Haas Bull Amer. Mus. nat. Hist. 108 1955; Perisphinctes (Paraboliceras?) japonicus sp. nov. Jurassic, Japan, H. Yabe Proc. Imp. Acad. Tokyo 19 9 1943: p. 583 f. 1.

†Permodomatoceras gen. nov. Domatoceratidae, genotype: P. trapezoidale sp. nov. Permian, South Urals., V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954: p. 95 pl. ix f. 2.

†Peroniceras de Grossouvre, emend. v. Hoepen, now included in Gauthiericeratidae fam. nov. Ammonoidea; P. platycostatum p. 367 f. 15-17; P. undulato-carinatum p. 369 f. 18-22 spp. nov. Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955.

†Peroniceratidae. Evolution, T. Matsumoto (1).

†Platyceratidae—association with echinoderms, A. L. Bowsher.

†Poecilomorphus arisdorfensis p. 35 pl. 6 f. 6a-c; P. lauwilbergensis p. 35 pl. 7 f. la-d; P. buckmani p. 36 pl. 7 f. 3a-c spp. nov. Jurassic, Switzerland, P. L. Maubeuge Schweiz. palaeont. Abh. 71 1955. †Polysphinctites ebrayoides sp. nov. Bathonian [Jurassic] Dorset, W. J. Arkell Palaeontogr. Soc. (Monogr.] 5 1955 p. 138 pl. xvi f. 3.

† Probillingsites kessleri sp. nov. Ordovician, Colorado, W. C. Sweet (1) J. Palaeont. 29 1 1955: p. 80 pl. 16 f. 1, 2.

†Procerites Siemiradzhi 1898. Designation of type species, Opinion 301.

†Proendoceras "gen. nov." Proterocameroceratidae, genotype: Cameroceras annuliferum Flower, 1941, Paleontographica Americana, 3 12 p. 25 pl. 1 f. 9-10. Roubidoux, North America, R. H. Flower J. Paleont. 29 3 1955: p. 350.

†Proteoceras gen. nov. Stereoplasmoceratidae genotype: Oonoceras perkinsi Ruedemann, 1906—P. pulchrum sp. nov. p. 825 pl. 80 f. 8-9 pl. 81 f. 8 Chazyan, North America, R. H. Flower J. Paleont. 29 5 1955.

†Protocanites lyoni from Mississippian of Nevada, W. M. Furnish (1).

†Pseudomuites gen. nov. Pachydiscidae. Type species: Pachydiscus ambiguus Grossouvre Upper Senonian of Europe and India (?), T. Matsumoto Mem. Fac. Sci. Kyushu Univ. 5D 3 1955; p. 169.

†Pseudorthoceras simense p. 100 pl. vi f. 6-8; P. neumannae p. 101 pl. vi f. 5 spp. nov. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 44 1954.

†Pseudotemnocheilus gen. nov. p. 69 Tainoceratidae, type species Nautilus postuberculatus Karpinsky, 1874 (= N. tuberculatus Verneuil, 1845); P. posttuberculatum (Karpinsky) p. 70 pl. vii f. 1 Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 50 1954.

†Puzosia willemventeri sp. nov. Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955: p. 357 f. 4, 5.

†Rhiphaeoceras gen. nov. Rhiphaeoceratidae fam. nov. q. v., V. E. Ruzhentzev & V. N. Shimanskii.

†Rhiphaeoceratidae fam. nov. p. 103 Rhiphaeoceras gen. nov. p. 103, genotype: R. venustum p. 105 pl. x f. 3, 4; R. humile p. 104 pl. x f. 2 spp. nov. Permian, Southern Ural, V. E. Ruzhentzev et V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954. †Rhiphaeonautilus gen. nov. p. 116 Rhiphaeoceratidae, genotype: R. curticostatus sp. nov. p. 117 pl. x f. 8. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

†Rhynchoteuthis kaibabensis sp. nov. Permian, Arizona [U.S.A.] L. F. Brady J. Palaeont. 29 1 1955 p. 104 pl. 21 f. 2-8.

†Salfeldiella (Goretophylloceras) moreti nom. nud.; S. (Aphroditiceras) aphrodite venus nom. nud. Cretaceous, Egypt, I. G. E. Mahmoud Bull. Inst. Egypt 36 1 1955: p. 223.

†Scyphoceras gen. nov. Scyphoceratidae fam. nov. q. v., Ruzhentzev & Shimanskii.

†Scyphoceratidae fam. nov. p. 134; Scyphoceras gen. nov. p. 135 genotype: S. dionysi p. 136 pl. xii f. 6; S. ellipticum p. 137 pl. xiii f. 1-4, pl. xv f. 2; S. angulatum p. 137 pl. xiii f. 5 spp. nov. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

†Seguenziceratidae—proposed suppression of this family-group name under the Plenary Powers, W. J. Arkell (2).

†Shikhanoceras gen. nov. Orthocerotoidea p. 116—S. sphaerophorum sp. nov. p. 118 pl. 111 f. 1-9. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 44 1954.

†Sholakoceras gen. nov. p. 111 Rhiphaeoceratidae genotype: S. bisulcatum p. 112 pl. xi f. 1; S. transforme p. 114 pl. xi f. 2; S. privum p. 115 pl. xi f. 3 spp. nov. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

†Simorthoceras gen. nov. Orthocerotoidea; S. gracile sp. nov. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 44 1954: p. 108 pl. 11 f. 7.

†Solenites Schlotheim, 1820 (an emendation of Solennites Schlotheim, 1813) proposed suppression of this genus by the Plenary Powers, W. J. Arkell (4).

†Solennites Schlotheim, 1813, proposed suppression of this genus by the Plenary Powers, W. J. Arkell (4)

†Sphaerocerus Bayle 1878. Validation, Opinion 300.

†Stereospyroceras gen. nov. Stereoplasmoceratidae: genotype: Stereospyroceras champlainense sp. nov. Chazyan, North America, R. H. Flower J. Paleont. 29 5 1955: p. 827 pl. 81 f. 1, 9.

†Storthoceras paucicostatum sp. nov. Jurassic, New Caledonia, J. Avias Sciences de la Terre, Nancy 1 1–2 1953: p. 132 pl. xvii f. 1a-b, fig. 72a, c.

†Strigites pseudostrigifer sp. nov. Jurassic, Switzerland, P. L. Maubeuge Schweiz. palaeont. Abh. 71 1955: p. 31 pl. 11 f. 5; pl. 2 f. 8; pl. 5 f. 3.

† Tabantaloceras gen. nov. Bactritoidea p. 86 type species T. planum sp. nov. p. 87 pl. xi f. 4. Permian, Southern Ural, V. N. Shimanskii Trav. Inst. Paléont. Acad. Sci. URSS 44 1954.

†Taramelliceras (Taramelliceras)
obumbrans p. 79 pl. 16 f. 1, f. 14d; T.
(Taramelliceras) pseudotrachinotum p. 89
Abb. 1/30; T. intersistens p. 116 pl.
19 f. 27; T. (Metahaploceras) semibarbarum p. 139 Abb. 19/167 spp. nov.;
Taramelliceras costatum rivale p. 100
pl. 17 f. 16; T. costatum nodoserratum
p. 101 pl. 17 f. 15; T. hauffianum rieberi p. 108 Abb. 8/87; T. pseudoflexuosum gracile p. 119 pl. 19 f. 24; T.
klettgovianum modeli pl. 19 f. 28; T.
kobyi quenstedti p. 129 Abb. 16/138—
subspp. nov. Jurassic, Germany, H.
Hölder Palaeontographica A 106 3-6

†Tarrantoceras gen. nov. Acanthoceratidae type species T. rotatile p. 59 pl. 5 f. 1-10; T. stantoni p. 60 pl. 5 f. 11-21; T. lillianense p. 60 pl. 5 f. 22-27; T. multicostatum p. 61 pl. 6 f. 21-23 spp. nov. Cretaceous, North America, L. W. Stephenson Prof. Pap. U.S. geol. Surv. 274-C 1955.

†Teichertia gen. nov. p. 496-497 Nautiloidea. Type species: Teichertia prora sp. nov. p. 497, pl. 54 f. 1-8, this species is related to typical Deltoidonautilus Mid Eocene, Giralia Range, North West Cape area, Western Australia, Glenister, Miller & Furnish J. Paleont. 30 3 1956.

†*Teshioites* gen. nov. Pachydiscidae, p. 173 Type species: *Teshiotes ryugasensis* p. 174 pl. 36 f. 1a, b, 2a, b; pl.

37 f. 3a b; text-f. 8a, b; T. teshioensis p. 177 pl. 3 f. 2a, b; pl. 35 f. 3a-c spp. nov. Upper Campanian (Cretaceous), Hokkaido & Saghalien, Japan, T. Matsumoto Mem. Fac. Sci. Kyushu Univ. 5D 3 1955.

†Tetragonoceras gen. nov. Erioliceratidae fam. nov. Ammonoidea: genotype: T. robustum sp. nov. Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955: p. 382 f. 37.

†Thrincoceratidae fam. nov. p. 118; Neothrincoceras gen. nov.; N. uralicum (Fredericks) p. 119 pl. xi f. 4, 5 nom. nov.; N. soshkinae p. 121 pl. xi f. 6 sp. nov. Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954.

†Trigonellites Parkinson, 1811, proposed suppression under the Plenary Powers of this genus and the species T. lamellosus and T. latus, W. J. Arkell (4).

†Tylonautilus nodosocarinatus (Roemer)—from Carboniferous of Oklahoma, Miller & Furnish (1).

†Uhligella toucasiformis 'nov. sp.' name without description. Cretaceous, Rumania, **D. Iacob** Bul. sti. Acad. Republ. rom. **5** 3 1953 : p. 665.

†Urakawites gen. nov. Pachydiscidae. Type species: Pachydiscus rotalinioides Yabe. Cretaceous, Japan, T. Matsumoto Mem. Fac. Sci. Kyushu Univ. 5D 3 1955: p. 167.

†Uralorthoceras tzwetaëvae sp. nov. Permian, Southern Ural, V. N. Shimanskii in litt: in Devingtal, V. V. Doklad. An. S.S.S.R. 81 6 1951: p. 1140. See also Shimanskii Trav. Inst. paléont. Acad. Sci. URSS 44 1954: p. 103.

†*Valdedorsella gignouxi* nom. nud. Cretaceous, Egypt, **I. G. E. Mahmoud** Bull. Inst. Egypt. **36** 1 1955 : p. 223.

†Vasseuria occidentalis Munier-Chalmas and its occurrence in the English Eocene, figs., **D. Curry**.

†Venatoroceras gen. nov. p. 139. Scyphoceratidae, genotype: V. verae sp. nov. p. 140 pl. xv f. 3 Permian, Southern Ural, V. E. Ruzhentzev & V. N. Shimanskii Trav. Inst. paléont. Acad. Sci. U.S.S.R. 50 1954. †Witchellia helvetica p. 37 pl. 7 f. 4a, b; W. (Maceratites) hansenbodensis p. 38 pl. 1 f.4; pl. 2 f. 2; pl. 3 f. 4a, b spp. nov. Jurassic, Switzerland, P. L. Maubeuge Schweiz. palaeont. Abh. 71 1955.

†Yokoyamaoceras jimboi (Yabe MS) p. 153 pl. ix f. 4a, b, 5a, b; Text-f. 13a, b; Y. (?) mysticum p. 155, pl. x f. 1a, b spp. nov. Cretaceous, Hokkaido, Japan, T. Matsumoto Jap. J. Geol. Geogr. 26 1955.

† Zuluscaphites gen. nov. Palaehopli-

tidae p. 360 genotype: Z. orycteropusi sp. nov. p. 359 f. 6. Cretaceous, Zululand, E. van Hoepen S. Afr. J. Sci. 51 12 1955.

INCERTAE SEDIS

†Neklania nom. nov. for Dalila Barrande, 1881 (Lamellibranchiata) (non Dalila Thomson, 1858) Genotype: Dalila (recte Neklania) obtusa Barrande, 1881. Silurian, Bohemia, A. Přibyl & B. Růžička Přírod. Sborn. Ostravsk. Kraje 14 1–2 1953: p. 57.